

**Relationship between Parkin Function, Heavy Metal
Homeostasis and Oxidative Stress in a *Drosophila*
Model of Parkinson's Disease**

Dissertation

zur

**Erlangung der naturwissenschaftlichen Doktorwürde
(Dr. sc. nat.)**

vorgelegt der

Mathematisch-naturwissenschaftlichen Fakultät

der

Universität Zürich

von

Nidhi Saini

aus

Indien

Promotionskomitee

Prof. Dr. Walter Schaffner (Vorsitz und Leiter der Dissertation)

Prof. Dr. Christian Frei (ETH Zürich)

Prof. Dr. Gerd Multhaup (Freie University, Berlin, Germany)

Zürich, 2010

Dedicated to my Parents

Rajendra Kumar Saini

and

Kanchan Saini

INDEX

Zusammenfassung.....	4
Summary.....	6
Introduction.....	8
 Results as 3 published manuscripts	
 Paper 1: Extended lifespan of <i>Drosophila parkin</i> mutants through sequestration of redox-active metals and enhancement of anti-oxidative pathways.....	51
 Paper 2: <i>Parkin</i> mutant in the fly is largely rescued by metal-responsive transcription factor (MTF-1).....	88
 Paper 3: Zinc supplement greatly improves the condition of <i>parkin</i> mutant <i>Drosophila</i>.....	127
 Conclusion and outlook.....	134
 Acknowledgements.....	137
 Curriculum Vitae.....	139

ZUSAMMENFASSUNG

Die Parkinsonsche Krankheit ist die zweithäufigste neurodegenerative Erkrankung. Die Haupt-Symptome sind Tremor (unkontrolliertes Zittern) und maskenhafte Gesichtszüge. Auf zellulärer Ebene geht die Krankheit einher mit fortschreitendem Verlust von Neuronen der *Substantia nigra* im Zentralnervensystem. Die meisten Fälle sind idiopathisch, d.h. ohne erkennbaren Auslöser, doch kommt die Krankheit auch familiär gehäuft vor, wobei Mutationen in bisher sechs Genen als Ursache identifiziert werden konnten. Eine Untergruppe der autosomal-rezessiven form der Parkinsonkrankheit wird durch Mutation im Parkin-Gen, das für eine E3 Ubiquitinligase kodiert, bewirkt. Es gibt mehrere Hinweise darauf, dass oxidativer Stress die Parkinsonkrankheit begünstigt. Weil eine Störung im Schwermetallhaushalt der Zelle oxidativen Stress bewirken kann, haben wir bei einer *Parkin*-Mutante der Essigfliege *Drosophila melanogaster* den Effekt einer veränderten Schwermetall-Homöostase getestet. Homozygote *Parkin*-mutante Fliegen zeigen schwere motorische Defekte einschliesslich Flugunfähigkeit, männliche und weibliche Sterilität, und stark verkürzte Lebensdauer. Wir konnten zeigen, dass eine Doppelmutante für *Parkin* und MTF-1 (metal responsive transcription factor, auch metal response element binding transcription factor genannt) nicht lebensfähig ist, d.h. "synthetische Letalität" zeigt. Die hauptsächlichen Zielgene von MTF-1 kodieren für Metallothioneine, kleine cysteinhaltige Proteine welche in der Lage sind etliche Metalle, z.B. Kupfer, Zink, Kadmium und Quecksilber zu binden, oder Radikale zu neutralisieren. Interessanterweise zeigen transgene Fliegen mit erhöhtem MTF-1-Spiegel eine dramatische Verlängerung der Lebensdauer und eine verbesserte Motorik, welche sogar zu kurzen Flügen befähigt. Zudem sind die Weibchen (allerdings nicht die

Männchen) fruchtbar. Auf zellulärer Ebene ist die Struktur der Ovarien und der Flugmuskeln weitgehend normalisiert. Weitere Experimente mit Parkin-Mutanten ergaben starke Verbesserungen im Phänotyp auf Futter mit BPS und BCS, zwei Chelatoren mit Präferenz für die redox-aktiven Metalle Eisen und/oder Kupfer. Schliesslich wurden auch noch eine verlängerte Lebensspanne, verbesserte Motorik und teilweise Wiedergewinnung der weiblichen Fertilität beobachtet, wenn *Parkin*-Mutanten auf Futter mit Zink-Zusatz aufgezogen und gehalten wurden. Ein besonders günstiger Effekt wurde dabei mit einer hohen Konzentration (4 mM Zinkchlorid) beobachtet, welche bei Kontroll-Fliegen bereits Vermeidungsreaktionen und einen verschlechterten Gesamtzustand bewirkten. Zink ist, im Gegensatz zu Kupfer und Eisen, kein redox-aktives Metall und wirkt unter den meisten Bedingungen als Antioxidans, was zumindest teilweise den positiven Effekt auf *Parkin*-Mutanten erklären könnte. Zusammengefasst lässt sich sagen, dass unsere Studien eine Verbindung zwischen Parkin-Funktion und Schwermetall-Homöostase nahelegen, welche auch für Säuger einschliesslich des Menschen von Bedeutung sein könnte.

SUMMARY

Parkinson's disease (PD) is the second most common neurodegenerative disease. The symptoms are tremor and rigidity and a mask-like face expression. At the cellular level PD is caused by progressive loss of neurons in a brain region termed *Substantia nigra*. PD in most cases is idiopathic, but a fraction is caused by specific dominant or recessive mutations in either of six genes known. Some autosomal recessive PD are caused by mutations in the gene for Parkin, a E3 ubiquitin ligase. Several lines of evidence point to oxidative stress as a major cause of PD. Since a distortion of cellular heavy metal homeostasis has been associated with increased oxidative stress, we analyzed the response of a *parkin* mutation (*park*²⁵) in the vinegar fly *Drosophila melanogaster* to modulation of heavy metal homeostasis. *Drosophila* homozygous *parkin* mutants suffer from severe motoric defects including inability to fly, male and female sterility, and a very short life span. We found that a combination of a *parkin* mutation with a loss-of-function mutation of MTF-1 (metal responsive transcription factor, also referred to as metal response element binding transcription factor) is not viable, i.e., displays a "synthetic lethal phenotype". The major target genes of MTF-1 are encoding metallothioneins, small cysteine-rich proteins with the ability to bind a number of metals, including copper, zinc, cadmium and mercury, and also to scavenge radicals. Interestingly, transgenic flies with elevated MTF-1 expression show a dramatic prolongation of life span and improvement of motoric abilities which even includes short patches of flight. Furthermore the females (but not the males) are fertile. At the cellular level, muscle and ovary morphology are largely normalized. In an independent set of experiments with *parkin* mutants, we also find a beneficial effect upon food supplementation with BPS and BCS,

chelators with a preference for the redox-active metals iron and/or copper. Finally, improved life span, motoric abilities and partial recovery of female fertility is also observed when *parkin* mutants are raised and kept on food with a supplement of zinc ions. A particularly good effect is seen at a food zinc concentration (4 mM) which is already showing avoidance reactions and adverse effects in control flies. Zinc, unlike iron and copper, is not redox-active and under most circumstances acts as an anti-oxidant, which might at least in part explain its positive effect on the *parkin* mutant flies. Taken together, our findings point to an intriguing interplay between Parkin function and metal homeostasis which may well have relevance to mammals including humans.

INTRODUCTION

I Parkinson's disease

II MTF-1

III zinc homeostasis

I) Parkinson's disease

1.1 Clinical and pathological features of Parkinson's Disease

Parkinson's disease (PD) is the most common chronic movement disorder and the second most common neurodegenerative disorder which was first described by James Parkinson in 1817. The incidence of this disease, markedly increases with age (Twelves et al., 2003), typically manifesting itself after the age of 50, with an estimation of about four million people worldwide and 1% of the population above the age of 65 years being inflicted (Bushnell and Martin, 1999; Chung et al., 2003). Clinical manifestations of PD patients include a broad range of motor abnormalities including rigidity, bradykinesia (slowness of voluntary movement), resting tremor and postural instability. The movement disorder in PD can be largely credited to the selective loss of dopaminergic neurons in the nigrostriatal pathway of the brain (Dawson, 2000; Parkinson, 2002; Spacey and Wood, 1999). There are several secondary symptoms, such as constipation, difficulties in swallowing, loss of bladder and/or bowel control, abnormalities in olfactory and visual perception, bradyphrenia (slow response to questions) and hypophonia (soft voice) in addition to psychosocial disorders, such as, anxiety and depression as well as dementia and cognitive impairment (Gelb et al., 1999). In order to be surely diagnosed with PD, a patient must have at least two or more of the primary symptoms, one of which should be either resting tremor or bradykinesia. A caveat is that several other diseases display similar symptoms defined for PD making a definitive diagnosis possible *post mortem*. The motoric impairments in PD are usually responsive to levodopa (L-DOPA), resulting in the improvement of these clinical symptoms (Gelb et al., 1999).

Pathologically, PD is characterized by the presence of cellular inclusions called "Lewy bodies" and a progressive but selective loss of dopaminergic neurons in the substantia nigra pars compacta (SNc) which is necessary for normal motor function.

This selective loss of dopaminergic neurons ultimately leads to the loss of dopamine (DA), its metabolites, 3,4 dihydroxyphenylacetate (DOPAC) and homovanillic acid (HVA), the dopamine transporter (DAT) in the striatum and the SNc, and its biosynthetic enzyme, tyrosine hydroxylase (Zhang et al., 2000). The first PD symptoms develop when about 70% of the striatal dopamine neurons and about 50% of the nigral dopamine neurons are lost (Dunnett and Bjorklund, 1999; Lansbury and Brice, 2002). Friedrich Lewy first described the pathological hallmark of PD, Lewy bodies in 1912 as a hyaline inclusion in the neuronal cytoplasm. Classical Lewy bodies are 5-25 μ M fibrillar, proteinaceous cytoplasmic inclusions with a dense eosinophilic core surrounded by a clear halo and are usually found in the SNc and locus coeruleus of PD patients (Huang et al., 2004; Pollanen et al., 1993). Misfolded proteins can be generated due to genetic mutations (Goldberg et al., 2003) or oxidative modifications (Dalle-Donne et al., 2006) and accumulate in Lewy bodies, most likely due to a chronic imbalance in their generation and clearance. The main component of Lewy bodies is α -Synuclein (Goedert and Spillantini, 1998; Iwatsubo, 2003; Spillantini et al., 1997). In some Lewy bodies, several other proteins were shown to be present besides α -Synuclein. These include Ubiquitin (Manetto et al., 1988), Ubiquitin-carboxyterminal hydrolase (UCH-L1) (Lowe et al., 1990), Synphilin-1 (Wakabayashi et al., 2000), Tau (Ishizawa et al., 2003), Parkin-associated-endothelin receptor-like receptor (Pael-R) (Murakami et al., 2004), Parkin (Schlossmacher et al., 2002), neurofilaments (Schmidt et al., 1991), Torsin A (Shashidharan et al., 2000) and heat shock proteins such as Hsp70 and Hsp90, which act as molecular chaperones (Auluck and Bonini, 2002). Misfolded proteins are prone to aggregate formation and have the potential to impair cellular functions (Gregersen et al., 2006). Lewy bodies most likely have a protective function by sequestering these potentially cytotoxic proteins and preventing them from damaging cellular organelles present in the cytoplasm. In agreement with this idea, dopamine neurons containing Lewy bodies have been reported to escape from degeneration in PD, suggesting that Lewy-body positive neurons are less vulnerable than their Lewy-body negative counterparts (Tompkins and Hill, 1997).

Neurobiochemical implications of PD include mitochondrial complex I dysfunction, reduced glutathione, increased lipid peroxide and iron levels and decreased ferritin

concentrations which suggest that dopaminergic neurons of PD patients are under oxidative stress (Kosel et al., 1999; Munch et al., 1998). Furthermore, recent studies have implied altered proteasomal function as one of the contributing factors in the pathogenesis of PD (McNaught et al., 2003).

Evidence suggests that both environmental and genetic factors play contributing roles in the etiology of the disease. Both exogenous environmental toxins and endogenous proteotoxins have been implicated in PD pathogenesis (Yang et al., 2003). The molecular mechanism(s) underlying the selective dopaminergic neuronal loss in PD are not clearly outlined, but accumulating evidence indicate that mitochondrial dysfunction, oxidative stress, impaired ubiquitin-dependent proteolysis and protein aggregation are involved in the pathogenesis of PD. Mutations in a number of genes have been found to be linked to various inherited forms (familial) PD and several of them are associated with the ubiquitin-proteasome protein degradation pathway. Although inherited forms of PD account for only about 5-15% of known PD cases (de Silva et al., 2000; Gasser, 2001; Mizuno et al., 2001), the symptoms and neuropathology are essentially identical to idiopathic PD. However, most PD cases are still believed to be sporadic. Hence, the hunt for the possible environmental contaminants or other non-genetic causes that are linked to the development of PD still remains an active area of research.

1.2 Genetics of PD

1.2.1 Parkin (PARK2)

The gene *PARK2* also known as *parkin* was identified by positional cloning in Japanese families (Kitada et al., 1998). The locus *PARK2* responsible for causing autosomal-recessive juvenile PD (AR-JP), characterized by early-onset, slow disease progression and mostly absence of Lewy bodies, barring an exceptional case (Farrer et al., 2001; Mori et al., 1998; Takahashi et al., 1994) was mapped to chromosome 6q25.2-27 (Matsumine, 1999).

The *parkin* gene is one of the largest genes in the human genome (1.53 Mb), containing 12 exons and encoding a 465-amino acid protein of 52 kDa molecular weight. The *parkin* promoter functions as a bi-directional promoter. It regulates the

transcription of *parkin* on the one hand and a gene upstream of *parkin* in an antisense orientation, called *PACRG* which encodes Parkin co-regulated gene (PACRG) or Gene located upstream of *parkin* (Glup) on the other hand (West et al., 2003).




Figure 1. Schematic representation of the Parkin/PACRG locus.

The chromosomal location is shown (left). 12 individual exons of *parkin* are marked (not to scale) with intron sizes indicated (middle). Protein domain structure is indicated (right). Adapted from Mata et al., 2004.

PACRG is a component of Lewy bodies and forms a part of the chaperone complex with Hsp70 and Hsp90 and it suppresses unfolded Pael receptor-induced neuronal cell death (Imai et al., 2003), however, its exact function is unclear at present. This suggests a model of toxicity reduction of Pael-R in dopamine neurons by the joint

action of Parkin which degrades its substrate, Pael-R and PACRG which sequesters Pael-R into Lewy bodies.

Structure, expression and function of Parkin

Besides a ubiquitous expression in the brain, Parkin is found in several cellular compartments, including the cytosol, trans-Golgi network, synaptic membrane compartment, postsynaptic densities, lipid rafts, cytoplasmic vesicles, endoplasmic reticulum, outer nuclear and outer mitochondrial membranes (Fallon et al., 2002; Kubo et al., 2001; Shimura et al., 1999; Stichel et al., 2000). Functional Parkin protein has been suggested to be a requirement for Lewy body formation as it has been found in Lewy bodies of autosomal dominant PD patients (Schlossmacher et al., 2002; Shimura et al., 2001). However, several other studies failed to detect Parkin in Lewy bodies or Lewy neurites by Parkin antibodies (Pawlyk et al., 2003).

Parkin has a ubiquitin-like (UBL) domain at its N-terminus and two RING (Really Interesting New Gene) fingers, a type of zinc finger flanking the IBR (In-Between-RING fingers) domain at its C-terminus. The UBL domain binds to the Rpn10 subunit of the 26S proteasome (Sakata et al., 2003) and is also involved in substrate binding (Shimura et al., 2000). The segment present in between the UBL and RING1-IBR-RING2 domains shows no clear similarity to known proteins and was accordingly termed the unique parkin domain (UPD) (Kahle et al., 2000a). The PDZ-binding motif at the C-terminus of Parkin interacts with the calcium/calmodulin-dependent serine protein kinase (CASK) (Fallon et al., 2002) which co-localizes with Parkin at postsynaptic terminals and in lipid rafts in rat brain *in vivo* as well as in synapses in cultured cortical neurons. There is high evolutionary conservation in Parkin, with orthologs in *Drosophila melanogaster*, *Caenorhabditis elegans*, mouse, rat and other species (Bae et al., 2003; Culetto and Sattelle, 2000). *Drosophila* Parkin shows 42% amino-acid sequence identity to human Parkin (Bae et al., 2003).

Human	-----MIVFVRFNSSHGFPVEVDSITIFQLKEVA
Mouse	-----MIVFVRFNSSYGFPVEVDSITIFQLKEVA
Rat	-----MIVFVRFNSSYGFPVEVDSITIFQLKEVA
Drosophila	MSFTFKFIATFVRKMLELLQFGGKILIH LSTIVKINTGKTLTVNLEPQWDIKNVKELVA
Human	KRQGV PADQLRVIFAGKELNDWTVQNCDDQOSTVHIVQR--PWRKQEMNATGGEDER
Mouse	KRQGV PADQLRVIFAGKELNHLTVQNCDDQOSTVHIVQR--PRRSHEINASGGDEPQ
Rat	KRQGV PADQLRVIFAGKELNHLTVQNCDDQOSTVHIVQR--PQRKSHEINASGGDKPQ
Drosophila	PQLGLQDDDKLIFAGKELSDATTLEQCDLGGQSVLHAIRLRPVQRQKIQSATLEEEE-
Human	NAAGGCERE PQSLIRVDLSSSVLPDGSVGLAVILHIDSKRDSPPAGSPAGRSTYNSFYVY
Mouse	STSEGSIWESRSLIRVDLSSHLPVDSVGLAVILDIDSKRDSBAARG-IVKPTYNSEFTY
Rat	STPEGSIWESRSLIRVDLSSHLPADSVGLAVILDIDSKRDSBAARGPAKPTYHSFVY
Drosophila	--PSLS-----DEASKPLNETLLDLQLESEERLNTIDEER--VR-AKAHFVH
Human	CKGPGQRVQPGKLRVQCSTCRQATLTLTQGPSCWDDVLIPNRMSEGCQSPHCFGTS----
Mouse	CKGPGCHKVQPGKLRVQCGTCKQATLTLAQGPSCWDDVLIPNRMSEGCQSPDCFGIR----
Rat	CKGPGCHKVQPGKLRVQCGTCKQATLTLAQGPSCWDDVLIPNRMSEGCQSPDCFGIR----
Drosophila	CS-QCDKLNGKLRVRCALCKGGFTVHRDPEWDDVLKSRRIFGHCESLEVACVDNAG
Human	---AEFFFKCGAHPIS-DKETPVALHLIATNSRNTTCTCTCDVRSPVLVFCQNSRHVIC
Mouse	---AEFFFKCGAHPIS-DKDISVALNLTISNRSTIPCTACTIDVRSPVLVFCQNRHVIC
Rat	---AEFFFKCGAHPIS-DKDISVALNLTISNRSTIPCTACTIDVRSPVLVFCQNRHVIC
Drosophila	DPPFAEFFFFKCAEHVSGGEKDFAAFLNLIKNNIKNV CLACTIDVSDIVLVFPCASQHVTC
Human	LDCEHLYCVIRLNDROFVHDPQLGYSLPCVAGCPNSLIKEIHFRITLGEQYNRYQQYGA
Mouse	LDCEHLYCVIRLNDROFVHDAQLGYSLPCVAGCPNSLIKEIHFRITLGEQYTRYQQYGA
Rat	LDCEHLYCVIRLNDROFVHDAQLGYSLPCVAGCPNSLIKEIHFRITLGEQYNRYQQYGA
Drosophila	IDCFRHYCRSLRGERQFMHPDFGTYLPCPAGCH HSFTIEITHFKLLITREEDRYQRFAT
Human	EECVLQMGVLCPRPGCGAGLLPEPQQRKVTCEGGNGLGCGFAFCRCKEAYHEGDCSAV
Mouse	EECVLQMGVLCPRPGCGAGLLPEQQRKVTCEGGNGLGCGFVFCRCKEAYHEGDCDSL
Rat	EECVLQMGVLCPRPGCGAGLLPEQQRKVTCEGGNGLGCGFVFCRCKEAYHEGDCSM
Drosophila	EEYVLQAGGV LCPPPGCGMGLLVEPDCRKVTQNG---- CGYVFCRNCLOGYHIGELPE
Human	FE-ASGITTTQAYRVDERAABQARWEAASKETIKKTKPCPRCHVPVERKNGGOMHKCPQP
Mouse	LE-PSGATISQAYRVDKRAABQARWEEASKETIKKTKPCPRCNVPIERKNGGOMHKCPQP
Rat	FE-ASGATISQAYRVDQRAABQARWEEASKETIKKTKPCPRONVPIERKNGGOMHKCPQP
Drosophila	GTGASATNSCEYTVDENRAAEARWDEASNVTIKVSTKE CPKCRIPTERDGGOMHVCIRA
Human	QCRLEWCWNGCEANRVCMGHWFEDV
Mouse	QCKLEWCWNGCEANRAQCMGHWFEDV
Rat	QCKLEWCWNGCEANRAQCMGHWFEDV
Drosophila	QCGFEWCWVQTEWTRDQMAHWFEG-

Figure 2: Alignment of the amino acid sequence of *D. melanogaster* parkin with human, mouse and rat. The conserved domains are in bold and ubiquitin-like domain is in red. RING domains are shown in green and IBR domain is indicated in blue. According to Bae et al., 2003.

Parkin functions as an E3 ubiquitin ligase

A number of RING finger proteins function as E3 ubiquitin protein ligase which play an important part in the ubiquitin proteasome system (UPS). The E3 ligases are responsible for the covalent attachment of Ubiquitin to proteins that are targeted for degradation by the proteasome (Joazeiro and Weissman, 2000). The UPS is involved in the temporal degradation of short-lived proteins which have important roles in cell-cycle, signal transduction and metabolism. Additionally, the UPS ensures protein quality control, it removes the misfolded or aggregated proteins by recruiting the E3

ubiquitin ligase, which further links the protein folding (chaperone) and degradation (proteasome) machinery of the cell.

Extensive molecular, genetic and functional analyses have shown that Parkin is an E3 ubiquitin protein ligase which is involved in protein turnover via the UPS (Shimura et al., 2000). Mutants for the Parkin RING domain lose their E3 ligase activity and the ability to bind to E2 ubiquitin conjugating enzymes, such as UbcH7 (Shimura et al., 2000). This indicates that the RING domain is required for recruitment of E2 enzymes into the Parkin complex. α -Sp22, a 22-kDa glycosylated form of alpha-synuclein notwithstanding, all Parkin substrates bind to Parkin via its RING finger domain (Shimura et al., 2001). Hence, mutations in the RING-IBR-RING domain can affect both substrate recognition as well as E2 binding, either of which results in partial or total loss of substrate degradation ability. Parkin can ubiquitylate substrates by joining Ubiquitin moieties either via the classical lysine 48 (K48) (Pickart, 2001) or via the unclassical lysine 63 (K63), resulting in totally different outcomes, as displayed by the Parkin-binding protein Synphilin-1 which is an α -Synuclein-interacting protein (Lim et al., 2005). While K48-linked ubiquitination of Synphilin-1 promotes its proteasomal degradation, K63-linked ubiquitination by Parkin may be involved in the formation of Lewy body inclusions formed by the coexpression of α -Synuclein and Synphilin-1 associated with PD (Lim, et al. 2005) (Chung et al., 2001). The relative expression levels between Parkin and Synphilin-1 determine the Synphilin-1 degradation kinetics. Appreciable degradation of Synphilin-1 by Parkin occurs only at an unusually high Parkin to Synphilin-1 expression ratio or when the system is primed for lysine-48 (K48) ubiquitination. In contrast, parkin ubiquitinates Synphilin-1 via K63-linked ubiquitin chains when Synphilin-1 and Parkin are coexpressed at equivalent levels (Lim et al., 2005). This suggests that Parkin may act as a dual-function ubiquitin protein E3 ligase capable of mediating both proteasomal-linked (K48) and proteasomal-independent (K63) ubiquitination under different cellular conditions.

Mutations in *parkin* cause autosomal-recessive juvenile PD (AR-JP)

The most frequent form of familial PD is Parkinsonism due to *parkin* mutations. Half of the inherited PD cases are estimated to be caused by mutations in *parkin* (Lohmann

et al., 2003). (Huynh et al., 2003). Homozygous mutations usually result in the loss of Parkin's E3 ligase activity along with altered Parkin levels (Finney et al., 2003). Parkin mutations include exonic deletions, insertions and several missense mutations, with the hotspots for point mutations in the two RING finger motifs, the IBR as well as the UBL domain (von Coelln et al., 2004a). In addition to homozygous mutations, compound heterozygous mutations have been identified in patients that displayed PD symptoms clinically indistinguishable from idiopathic PD (Klein et al., 2000).

Substrates of Parkin

(A) Substrates forming components of Lewy bodies

A.1 α -Synuclein: Parkin interacts with a rare O-linked glycosylated form of α -Synuclein (α -Sp22) in Lewy bodies that were extracted from post-mortem brain tissue of AR-JP patients (Shimura et al., 2001), while mutant *parkin* fails to ubiquitinylate α -Sp22 *in vitro* (Chung et al., 2001). Parkin protects against α -Synuclein toxicity *in vitro* (Petrucelli et al., 2002) and prevents α -Synuclein induced dopaminergic neuron death in *Drosophila* and rat models of PD (Lo Bianco et al., 2004; Yang et al., 2003).

A.2 Synphilin-1

Synphilin-1 is a major component of Lewy bodies found in PD patients (Wakabayashi et al., 2000). Parkin ubiquitinates Synphilin-1, an α -Synuclein-interacting protein (Chung et al., 2001). Upon binding to synaptic vesicles Synphilin-1 possibly regulates the synaptic functions of α -Synuclein, by anchoring it to vesicle membranes (Ribeiro et al., 2002). Functional Parkin is essential for the formation of Lewy body inclusions. This is consistent with the absence of Lewy bodies in patients with AR-JP.

A.3 Pael-Receptor (Pael-R)

The Parkin-associated endothelin receptor-like receptor (Pael-R) is a putative G-protein-coupled multipass endoplasmic reticulum transmembrane protein which is abundantly expressed in dopamine neurons of the substantia nigra (Imai et al., 2001) and is found in the Lewy bodies of PD patients (Murakami et al., 2004). The folding of Pael-R is assisted by the 70-kDa heat-shock protein Hsp70. Overexpressed Pael-R

is insoluble and it triggers the so-called unfolded protein response which leads to ER stress. Insoluble Pael-R is ubiquitinated by Parkin with the help of ER-resident ubiquitin-conjugating enzymes (Ubc6 and Ubc7) and sent for proteasomal degradation before its accumulation (Imai et al., 2001).

A.4 p38 subunit of the aminoacyl-tRNA synthetase complex

The degradation of p38, a major structural component of the aminoacyl- tRNA synthetase complex is carried out by Parkin mediated ubiquitination (Corti et al., 2003). Overexpression of p38 induces cell death. p38 is found in Lewy bodies of patients with idiopathic PD.

A.5 Synaptotagmin XI

Synaptotagmin XI is localized to the core of Lewy bodies and its interaction with Parkin results in its ubiquitination and degradation (Huynh et al., 2003). The synaptotagmin family is crucial for vesicle formation and docking. The interaction with a member of the synaptotagmin family suggests an involvement of Parkin in the regulation of proteins involved in controlling neurotransmitter trafficking at the presynaptic terminal.

(B) Substrates regulating neurotransmitter secretion and uptake

B.1 Dopamine transporter (DAT)

DAT belongs to the catecholamine transporter protein family and is localized to the plasma membrane of dopamine neurons (Hersch et al., 1997). It regulates the uptake of dopamine from the synaptic cleft (Torres et al., 2003). Lack of glycosylation promotes the misfolding and aggregation of DAT which further oligomerizes with correctly folded DAT, affecting the targeting of normal DAT to the plasma membrane (Sorkina et al., 2003). Parkin significantly enhances the ubiquitination and subsequent degradation of misfolded DAT (Jiang et al., 2004). Results showing a significantly reduced DAT in the striatum of *parkin* knockout mice are in perfect agreement with the required role of Parkin in DAT targeting (Itier et al., 2003).

(C) Polyglutamine proteins

Polyglutamine proteins are aggregation-prone cytosolic proteins. Cytosolic protein-misfolding stress results in an upregulation of Hsp70 which further recruits Parkin

through its interaction with the RING-IBR-RING domain (Tsai 2003). The polyglutamine protein is then ubiquitinated and targeted for proteasomal degradation. This process might be facilitated by the direct binding of the UBL domain of Parkin to the regulatory 19S proteasome (Sakata et al., 2003).

(D) Substrates involved in cell cycle regulation and cytoskeleton structure

D.1 Septin family of proteins - CDCrel-1 and CDCrel-2a

Cell-division control-related protein-1 (CDCrel-1), also known as Sept5, belongs to the a family of GTPase called septin. It is predominantly expressed in the nervous system and is ubiquitinated and degraded by Parkin (Beites et al., 1999; Zhang et al., 2000). CDCrel-1 accumulates in the brains of AR-JP patients (Choi et al., 2003). CDCrel-2a, another member of the septin family having strong homology to CDCrel-1, coimmunoprecipitates with Parkin from human substantia nigra lysates and is also ubiquitinated by Parkin *in vitro* (Choi et al., 2003). (Beites et al., 1999). Septins are evolutionary conserved proteins with essential functions in cytokinesis, and throughout the cell cycle. These small proteins likely with GTPase activity seem to function in membrane transport and exocytosis (Field and Kellogg, 1999).

D.2 Cyclin E

A number of RING proteins including Skp1, Cullin and F-box (SCF) complexes have been shown to function within multiprotein ubiquitin ligase complexes (Joazeiro and Weissman, 2000; Patton et al., 1998). Parkin is part of a SCF complex containing hSel-1 and Cullin-1 (Staropoli et al., 2003). This complex auto-ubiquitinates Parkin and also ubiquitinates Cyclin E via HSel-1 (Staropoli et al., 2003). Cyclin E is involved in kainite-induced excitotoxicity and primary neuronal apoptosis. Parkin knockdown by RNA interference results in the accumulation of Cyclin E in the primary cerebellar granule cells, while Parkin overexpression inhibits kainite-induced Cyclin E apoptosis (Staropoli et al., 2003).

D.3 α - and β -Tubulin

It was shown that Microtubule-associated protein 1A (MAP1A) binds Parkin in a temperature-sensitive manner with a higher coimmunoprecipitation at 37°C than at 4°C (Feng and Lin, 2001). This suggested that Parkin may be binding to MAP1A via microtubules, as polymerization of microtubules is known to be more efficient at

37°C. Parkin was later identified as a tubulin-binding protein and an E3 ligase for α - and β -Tubulins (Ren et al., 2003). Interestingly, Parkin overexpression stabilized microtubules against colchicine-induced depolymerization (Yang et al., 2005).

***Drosophila parkin* null mutants**

Animal models of PD have facilitated the study of the mechanisms and processes involved in the pathogenesis of the disease. Till date, none of the animal models have been able to recapitulate all the hallmark pathologies and symptoms (Auluck and Bonini, 2002; Hamamichi et al., 2008). These models are generated via knockdown or mutagenesis of genes implicated in PD or by treatment with environmental toxins like 1-methyl, 4-phenyl-1,2,3,6-tetrahydropyridine and paraquat (Dauer and Przedborski, 2003). In our study, we use a *Drosophila* model of PD. To explore the biological role of *parkin*, a series of mutations have been generated in the *Drosophila* ortholog of *parkin*, including deletion, nonsense and missense mutations. Two lines of *parkin* null mutant flies were generated by insertional mutagenesis with activated EP element transposons by two independent groups (Greene et al., 2003; Pesah et al., 2004). In one line (used in our study), *parkin*-deficient males are sterile but the females are reported to be fertile (sterile in our hands) (Greene et al., 2003). Male sterility was further studied and was found to be derived from a spermatid individualization defect at a late stage of spermatogenesis, which may be a result of defective Nebenkern formation and/or function (Greene et al., 2003). The Nebenkern is a specialized mitochondrial derivative in the mature sperm. Ultrastructural analysis of *parkin* mutant males revealed severe disruption of the Nebenkern integrity. In contrast to this study, the other group reported both male and female sterility with homozygous *parkin* deletion (Pesah et al., 2004). They also reported an increased sensitization to oxidative stress and a 30% lower mass of *parkin* mutants in comparison to wild type controls. A reduced cell size and cell number accounted for this mass reduction.

Parkin-deficient flies show a reduced longevity and a partially penetrant downturned wing phenotype (Greene et al., 2003). Locomotor activity assays revealed severe defects in flight and climbing activity. These locomotor defects derive from apoptotic cell death of mitochondria and muscle subsets. Histological analysis of the major flight muscles from *parkin* mutants showed an extensive disruption of muscle

integrity, which was consistent with their inability to fly. Ultrastructural analysis of the flight muscles demonstrated a disintegration of the mitochondria with fragmented cristae which indicate that Parkin might regulate mitochondrial integrity. Similar to *parkin* knockout mice (Goldberg et al., 2003; Itier et al., 2003; Palacino et al., 2004; Perez and Palmiter, 2005; Von Coelln et al., 2004b), no significant dopamine neuron loss was observed in the mutant flies (Greene et al., 2005; Greene et al., 2003; Pesah et al., 2004), notwithstanding the observation of a 20% loss in one study involving only one subset of the dopamine neuron clusters in the fly brain (Whitworth et al., 2005). Transcriptional analysis indicated that genes involved in the innate immune response and oxidative stress components are upregulated in the *parkin* mutant animals (Greene 2005). A genetic screen for mutations in genes that either enhance or suppress the lethality of *Drosophila parkin* mutants revealed that loss-of-function mutations in oxidative stress components, such as glutathione S-transferase S1 enhance the Parkin mutant phenotype (Greene et al., 2005). The lifespan of the mutant flies is further reduced when they are grown on environmental toxins like rotenone or paraquat, which generates toxic free radicals *in vivo* (Pesah et al., 2004).

1.2.2 α -Synuclein (PARK 1)

α -Synuclein (also known as NACP, Synuclein-1 or Synelfin) was first discovered in the *Torpedo* (electric eel) electrical organ as a vesicle-associated component (Clayton and George, 1999) and was identified as the precursor protein of the non-amyloid β component of the senile plaques of Alzheimer's disease (Iwai, 2000). α -Synuclein has 140 amino acids and consists of three domains, including an amino-terminal amphipathic lipid binding α -helix, a hydrophobic β -amyloid binding domain that encodes the non-A β component (NAC) and an acidic carboxy-terminal tail.




Figure 3: Schematic representation of α -Synuclein. The N-terminal region consists of 7 imperfectly conserved KTKEGV repeats. This motif has unknown physiological significance. The NAC domain is extremely hydrophobic and the C-terminal region is negatively charged. The three known missense mutations are localized to the N-terminal repeat region.

Either oxidative stress or mutations of *α -synuclein* may lead to the formation of highly toxic protein aggregates or fibrils, ultimately resulting in neuronal cell death. Missense mutations in the *α -synuclein* gene have been shown to be a cause of rare forms of autosomal dominant familial PD. One of the first identified genetic causes of PD is the missense mutation A53T in the *α -synuclein* gene in a large Italian and several smaller Greek kindreds. (Polymeropoulos et al., 1996; Polymeropoulos et al., 1997). Thereafter, two other missense mutations were identified; A30P and E46K (Kruger et al., 1998; Zarranz et al., 2004). Additionally, a triplication of the wild type *α -synuclein* gene resulting in α -Synuclein overexpression is also associated with familial PD (Singleton et al., 2003), indicating that both the composition and dosage of α -Synuclein appear to play roles in the development of PD. Oxidative stress has been shown to promote the aggregation of α -Synuclein in several studies (Hashimoto, M. 1999, Neuroreport 10: 717-721. Oxidative stress induces amyloid-like aggregate formation of NACP/alpha-synuclein in vitro) (Ostrerova-Golts, N. 2000. J. Neurosci 20: 6048-54) (Paxinou, E. 2001, J.Neurosci 21: 8053-61) (Scherer, TB 2002, J. Neurosci 22: 7006-15). Modifications in the primary structure of the α -Synuclein protein may enhance its oligomerization and subsequent fibril formation (Conway et al., 1998).

Localization and putative functions of α -Synuclein

The location of α -Synuclein is predominantly in the brain and in presynaptic terminals but its normal physiological functions is largely unknown (Clayton and George, 1999; Jakes et al., 1994; Kahle et al., 2000b), though it appears to be involved in the biosynthesis of dopamine, synaptic vesicle recycling (Baptista et al., 2003; Perez et al., 2002) and synaptic plasticity (Clayton 1999). α -Synuclein associates with lipid membranes (Jo et al., 2000) and co-localizes with Synaptophysin and SNAP-25, which promote vesicle docking and fusion (Irizarry et al., 1996; Withers et al., 1997). *In vitro* studies revealed that α -Synuclein can bind these structures through its N-terminal repeat domain, thereby causing dramatic alterations in its secondary structure (Davidson et al., 1998; Jensen et al., 1998). This led to the hypothesis that α -Synuclein has an important function in vesicular transmitter release and synaptic plasticity (Clayton and George, 1998). Different animal models have been created to investigate how α -Synuclein causes PD. α -Synuclein knockout mice display neurodegeneration along with altered compartmentalization of presynaptic dopamine and decreased capacity of dopamine storage pool (Yavich et al., 2004). In addition the reduction in the quantal release of dopamine that usually occurs post repetitive stimulation of the same neurons was inhibited in mice lacking α -Synuclein while the dopamine-dependent response to amphetamine was increased (Abeliovich et al., 2000). These results are indicative of the role of α -Synuclein in the regulation of dopamine homeostasis. However, the involvement of α -Synuclein is not only restricted to dopamine neurons, as its loss has been shown to alter synaptic transmission at glutamatergic synapses as well (Martin et al., 2004). α -Synuclein transgenic flies display a selective dopaminergic neuronal loss and the presence of Lewy body-like inclusions in the neurons (Feany and Bender, 2000). Finally, several studies point towards α -Synuclein's role as a molecular chaperone owing to its 40% homology with the cytoplasmic chaperone 14-3-3 (Kim et al., 2000; Ostrerova et al., 1999).

α -Synuclein aggregation and toxicity

α -Synuclein has a high propensity to aggregate owing to its hydrophobic non-amyloid- β domain. α -Synuclein is transformed from its normal, soluble form to a disease-causing fibrillar state in many neurodegenerative disorders, which are

collectively referred to as α -synucleinopathies (Galvin et al., 2001; Goedert, 1999; Spillantini and Goedert, 2000). While the C-terminal of α -Synuclein is a key regulator of its aggregation *in vivo* (Li et al., 2005), the internal hydrophobic NAC region is capable of fibril formation *in vitro* (Han et al., 1995). The fibrils formed are similar to those observed in Lewy bodies, particularly in the A53T mutation (Conway et al., 1998). The potentially neurotoxic protofibrils formed may damage vesicular membranes by forming pore-like channels (Lashuel et al., 2002; Volles and Lansbury, 2002), similar to some bacterial toxins (Ding et al., 2002; Volles et al., 2001) and permeabilize the membranes of synaptic vesicles and other organelles, resulting in leakage of calcium and dopamine which are known to be cytotoxic at high levels (Caughey and Lansbury, 2003). This provided a mechanism of how protein fibrilization can cause cell death. Interestingly, the dopamine- α -Synuclein adducts stabilize the protofibrils (Conway et al., 2001). A pathological modification involving the phosphorylation of Ser129 in α -Synuclein was shown to promote aggregation besides forming a major component of Lewy bodies (Anderson et al., 2006; Smith et al., 2005). The G-protein-coupled receptor kinase 5 has been recently shown to be responsible for this phosphorylation (Arawaka et al., 2006). α -Synuclein has been reported to directly bind the human dopamine transporter in cotransfected cells (Lee et al., 2001) which may result in elevated dopamine levels and subsequent neuronal death as dopamine may participate in oxidative stress generating reactions, since both dopamine auto-oxidation and its metabolism produce reactive oxygen species (ROS). In addition, α -Synuclein inhibits Tyrosine hydroxylase (Perez et al., 2002). Several studies link mutated α -Synuclein to an increased sensitivity of cells to proteasomal inhibitors by a decreased proteasomal function (Petrucci et al., 2002; Stefanis et al., 2001; Tanaka et al., 2001). Synphilin-1 binds to α -Synuclein *in vivo* and this association promotes the formation of intracellular inclusions (Engelender et al., 1999). In addition, it is suggested that Synphilin-1 regulates the synaptic functions of α -Synuclein, possibly by anchoring α -Synuclein to vesicular membranes (Ribeiro et al., 2002).

α -synuclein was the first gene shown to be associated with a heritable form of PD and was also the first of all PD-related genes to be studied in a *Drosophila* model of PD. Loss of dopamine neurons was observed in a transgenic *Drosophila* model of PD

caused by mutant α -Synuclein (Feany and Bender, 2000) in contrast to observations with several lines of *α -synuclein* transgenic mice which failed to develop dopamine neuron loss (Fernagut and Chesselet, 2004). On the other hand, rats overexpressing mutant synuclein, A30P (Klein et al., 2002; Lo Bianco et al., 2002) or A53T (Kirik et al., 2003), developed a progressive loss of dopaminergic neurons.

1.2.3 PINK1 (PARK6)

The ubiquitously expressed *PINK1* transcript encodes a 581 amino acids protein (Unoki and Nakamura, 2001) with a mitochondrial targeting motif at the N-terminal and a highly conserved protein kinase domain (156-509 amino acids) (Rohe et al., 2004; Valente et al., 2004), which is homologous to the serine/threonine kinases of the calcium/calmodulin family. Known mutations in *PTEN-induced kinase 1* (*PINK1*), associated with the PARK6 locus which result in early-onset autosomal-recessive PD include a W437Stop, nonsense and a G309D, missense mutation within the PINK1 kinase domain (Valente et al., 2004) and an insertion outside the kinase domain causing a frameshift and truncation at the C-terminus of PINK1 (Rohe et al., 2004). The mutants have an increased susceptibility to cellular stress. The wild type PINK 1 may prevent PD by protecting neurons from stress-induced mitochondrial dysfunction through phosphorylation of mitochondrial proteins (Valente et al., 2004). Loss-of-function mutations in the *Drosophila PINK1* ortholog results in morphological abnormalities of the mitochondria and apoptosis in testes and flight muscles with dopaminergic neuronal loss in some cases (Clark et al., 2006; Park et al., 2006; Yang et al., 2006). Parkin overexpression rescues the PINK1 phenotype but the vice versa is not true. This indicates a functional relationship between the two genes implicated in PD, suggesting that *PINK1* functions upstream of *parkin* (Kitada et al., 2007).

1.2.4 Ubiquitin C-terminal hydroxylase L1 (PARK5)

Ubiquitin C-terminal hydroxylase L1 (UCH-L1) is a thiol protease that produces ubiquitin possibly by cleaving ubiquitin polymers or by releasing ubiquitin from cellular adducts with glutathione or amine (Larsen et al., 1998; Larsen et al., 1996). An autosomal dominant missense mutation (I93M) in the gene encoding UCH-L1 was first identified in two siblings having a strong family history of PD (Leroy et al.,

1998). UCH-L1 is an abundant neuronal enzyme that makes up 1-2% of the total brain protein, with an unknown function (Wilkinson et al., 1989). Quite interestingly, UCH-L1 has two enzymatic activities with opposite effects, both of which may play a role in proteasomal protein degradation: a beneficial hydrolase activity and a partly pathogenic dimerization-dependent ligase activity which causes the accumulation of α -synuclein in cultured cells. The I93M mutation decreases the *in vitro* hydrolytic activity of UCH-L1 (Liu et al., 2002). UCH-L1 is also present in Lewy bodies of sporadic PD (Lowe et al., 1990) and I93M mutant was shown to be able to produce inclusions (Ardley et al., 2004). A S18Y polymorphism was also discovered in the UCH-L1 protein and subsequently found to be linked to a decreased susceptibility to PD (Levecque et al., 2001; Maraganore et al., 1999; Satoh and Kuroda, 2001), possibly because it reduces the ligase activity of UCH-L1 while promoting its hydrolase activity (Liu et al., 2002).

1.2.5 DJ-1 (PARK7)

PARK7, which encodes a protein DJ-1 is yet another locus linked to early-onset recessive PD. Various mutations identified in affected individuals include exonic deletions, truncations and point mutations (L166P and M26I) (Abou-Sleiman et al., 2003; Bonifati et al., 2003; Hague et al., 2003; Hedrich et al., 2004). DJ-1 was identified as a protein that transforms mouse NIH3T3 cells in cooperation with activated ras (Nagakubo et al., 1997). Apart from playing a direct role in dopamine neuron physiology (Goldberg et al., 2005), DJ-1 is a component of a cAMP-responsive multiprotein complex that stabilizes (Hod et al., 1999), and it also functions as a positive regulator of the androgen receptor, antagonizing the function of negative regulators, such as DJ-1-binding protein. The human DJ-1 gene, which is comprised of eight exons, is mapped to 1p36.2–36.3, where many chromosome aberrations in cancer have been reported (Taira et al., 2001). DJ-1 was found to be more strongly expressed in the testis than other tissues and also in sperm.

DJ-1 plays the role of a redox-sensitive molecular chaperone *in vitro* and in cells, granting protection against ROS-generated toxicity by getting induced in an oxidative environment and inhibiting the aggregate formation of α -Synuclein (Schapira, 2008;

Shendelman et al., 2004). Also, overexpression of DJ-1 reduces, while DJ-1 knockdown by RNA interference enhances, the susceptibility of cells to oxidative stress (Taira et al., 2004). DJ-1 dimerization is essential for its function and the L166P mutation destabilizes DJ-1 by preventing dimer formation (Macedo et al., 2003; Moore et al., 2003), abolishing DJ-1 chaperone activity (Shendelman et al., 2004). Different studies in DJ-1-null mice have shown that DJ-1 protects against neuronal oxidative stress and that its loss may lead to PD by conferring hypersensitivity to dopaminergic insults (Goldberg et al., 2005; Kim et al., 2005). The *Drosophila* genome encodes two orthologs of the human *DJ-1* gene, *DJ-1a* and *DJ-1b*. while DJ-1b is expressed ubiquitously, DJ-1a is predominant in the testes (Menzies et al., 2005; Meulener et al., 2005; Park et al., 2005).

1.2.6 Leucine-rich repeat kinase 2 (PARK8)

Leucine-rich repeat kinase 2 also known as LRRK2 or Dardarin is a large protein and a member of the leucine-rich repeat kinase family which in humans is encoded by the *LRRK2* (*PARK8*) gene. Dominant mutations of LRRK2 are the most common cause of inherited PD. Several missense mutations and one putative splice site mutation in the *PARK8* gene are linked to the development of PD (Paisan-Ruiz et al., 2004; Zimprich et al., 2004). The pathological markers of the disease vary significantly (Brice, 2005; Wszolek et al., 2004; Zimprich et al., 2004). LRRK 2 is present largely in the cytoplasm but also associates with the mitochondrial outer membrane. It is expressed weakly throughout the brain, with slightly elevated levels in the putamen and the substantia nigra regions. The protein consists of 2527 amino acids and is encoded by 51 exons, encoding multiple conserved domains with an ankyrin repeat region, a leucine-rich repeat (LRR) domain, a kinase domain, a DFG-like motif, a RAS domain, a GTPase domain, an MLK-like domain, and a WD40 domain. (Paisan-Ruiz et al., 2004; Shen, 2004). LRRK2 is known to interact with the RING2 domain of Parkin and is hypothesized to be responsible for the phosphorylation of α -Synuclein and Tau (Shen, 2004).

1.3 Mechanisms of cell death in PD

Although about 95% cases of PD seem to be sporadic, inherited cases have resulted in discovery of PD-linked genes. The genes encoding α -Synuclein, Parkin, DJ-1,

PINK1, UCH-L1 and LRRK2 are known to carry mutations that cause PD. Several homozygous and compound heterozygote mutations may result in loss of protein function, altered protein stability and/or localization or aggregation of accumulated proteins. Key mechanisms contributing to the pathogenesis of PD include mitochondrial impairment, oxidative stress, protein mishandling and altered dopamine homeostasis.




Figure 4: Genetics and molecular mechanisms of PD. Black arrows indicate putative primary causes of PD; dashed arrows indicate effects that are secondary. Blue arrows indicate mechanisms of PD that are secondary to primary causes. mtDNA, mitochondrial DNA. According to Greenamyre and Hastings, 2004.

1.3.1 Oxidative stress and mitochondrial dysfunction

Post mortem studies have repeatedly implicated oxidative stress in PD pathogenesis (Valente et al., 2004). Gene knockout of parkin mouse and flies show increased oxidative stress and mitochondrial dysfunction (Palacino et al., 2004; Pesah et al., 2004). Reactive oxygen species (ROS) can lead to functional alterations in biomolecules like proteins, lipids and DNA. Immunocytochemical analyses revealed an increased level of oxidized RNA and DNA in nigral neurons of PD patients (Alam et al., 1997; Zhang et al., 1999). Leading candidates for generation of oxidative stress include dopamine metabolism, mitochondrial dysfunction, increased free iron levels, impaired free radical defense and environmental chemicals (Foley and Riederer, 2000). ATP production in the mitochondria is regulated by respiratory chain complexes that are involved in the transfer of electrons to oxygen, resulting in the generation of free ROS. While cells tolerate normal levels of ROS, its overproduction combined with insufficient antioxidant defense mechanisms can induce mitochondrial dysfunction, which in turn may enhance the production of more deleterious ROS, further worsening the mitochondrial disturbed state, thus creating a vicious cycle (Tritschler et al., 1994). Nigrostriatal dopaminergic neurons might be particularly exposed to oxidative stress even under physiological conditions, probably due to redox cycling of catechols, resulting in the generation of detrimental ROS. Dopamine (DA) readily oxidizes and reacts with proteins, nucleic acids and lipids forming neurotoxic derivatives, including 6-hydroxydopamine, DA-quinones, superoxide radicals and hydrogen peroxide (Lotharius and Brundin, 2002; Olanow and Tatton, 1999) and its interaction with intracellular iron results in formation of toxic oxygen radicals (Sulzer and Zecca, 2000). Additionally, DA turnover pathway involving the enzymatic deamination of dopamine by monoamine oxidase generates hydrogen peroxide along with the non-toxic dopamine metabolites, 3,4-dihydroxyphenylacetic acid and homovanillic acid (Graham, 1978). To escape from cytosolic DA stress, DA neurons have evolved several defense mechanisms which include feedback inhibition of tyrosine hydroxylase, metabolism by monoamine oxidase, sequestration of DA in synaptic vesicles as well as reduction of DA quinone by glutathione (Nakamura et al., 2001). The quantification of the reaction products of ROS in post-mortem PD brain tissue serves as an indirect index of oxidative stress induced cell damage (Foley and Riederer, 2000).

1.3.2 Endoplasmic reticulum stress

The endoplasmic reticulum (ER) caters to multiple cellular functions. Accumulation of misfolded proteins within the ER disturbs its normal functions, induces a highly specific and evolutionarily conserved unfolded protein response (UPR) and is the major cause of ER stress (Kaufman, 1999). ER stress is aimed initially at compensating for damage but can eventually trigger cell death if ER dysfunction is severe or prolonged. The UPR upregulates ER chaperone and folding catalysts and prevents further protein aggregation by attenuating protein synthesis leading to the activation of the ER-associated protein degradation (ERAD) for eliminating the harmful aggregates (Szegezdi et al., 2003). The mechanisms by which ER stress leads to cell death remain enigmatic, however, several mechanisms including direct activation of proteases, kinases, transcription factors, and Bcl-2-family proteins and their modulators have been implicated. ER stress has been demonstrated to produce mitochondrial dysfunction by interfering with the expression and assembly of the cytochrome c oxidase (Hori et al., 2002). Also, the steady state levels of nuclear-encoded cytochrome c oxidase were reduced and the mitochondria-encoded subunit was rapidly degraded (Hori et al., 2002). UPR causes the ER-resident kinases to induce genes encoding ER chaperones and nuclear transcription factors that ultimately result in either ER stress reduction, achieved by phosphorylation of the eukaryotic initiation factor-2 α (eIF2 α) resulting in the attenuation of protein synthesis (Harding et al., 1999) or cell death if the damage is beyond repair (Mori, 2000).

In a cell culture model of PD, ER stress was inferred from the identification of phosphorylated protein kinase R-like ER kinase (PERK) and Inositol-requiring-1 (IRE1) in rat neuronal PC12 cells upon 6-hydroxydopamine exposure, which was taken up by the dopamine transporters and resulted in selective dopamine neuron death (Ryu et al., 2002). PERK (PKR-like ER kinase) is a serine-threonine protein kinase, the catalytic domain of which shares substantial homology to other kinases of the eukaryotic initiation factor 2 (eIF2) family (Harding et al., 1999; Shi et al., 1998). UPR was evoked by activated PERK and IRE1 (Ryu et al., 2002). Familial mutations in the ER-associated E3 ubiquitin ligase Parkin have also been associated with ER stress-induced cell death and are found in patients with familial PD (Dawson and Dawson, 2003; Takahashi et al., 2003). Importantly, Parkin expression is induced by ER stress, suggesting a role for it in adaptation to ER stress. Knockdown of *parkin* in

cell culture resulted in accumulation of its substrate Pael-R, which caused ER stress and cell death (Imai et al., 2001). These studies indicate that ER stress may be another mechanism underlying PD pathogenesis.

II) Metal-responsive transcription factor-1 (MTF-1)

MTF-1 (also metal-regulatory transcription factor-1), a zinc-finger transcription factor, is evolutionarily conserved from insects to mammals and activates cellular defense mechanisms in response to metals like zinc, copper and cadmium (Lichtlen and Schaffner, 2001; Westin and Schaffner, 1988). Its DNA binding domain consists of six C₂H₂ zinc fingers (Dalton et al., 1997; Radtke et al., 1993) and thus zinc is essential for the binding of MTF-1 to DNA. The DNA-binding motif is termed metal response element (MRE) and has a core consensus sequence of TGCRCNC (Stuart et al., 1984; Stuart et al., 1985). Copper and cadmium regulate MTF-1 indirectly by displacing zinc from zinc-loaded metallothioneins (MTs) or other zinc-binding proteins *in vitro* (Bittel et al., 1998; Zhang et al., 2003). However, *in vivo*, MTF-1 is subject to more complex regulation that includes phosphorylation and nuclear translocation (Saydam et al., 2002; Saydam et al., 2001). In quiescent cells, MTF-1 is predominantly cytoplasmic, but is translocated to the nucleus upon metal load or other stress conditions. In the nucleus, MTF-1 binds to the MREs and activates its target genes including metallothioneins, zinc transporters and placenta growth factor (Cramer et al., 2005; Langmade et al., 2000).

The MTF-1 regulatory protein has been characterized in human (Brugnera et al., 1994), mouse (Radtke et al., 1993), pufferfish (Auf der Maur et al., 1999), *Drosophila* (Zhang et al., 2001), zebrafish (Chen et al., 2002), carp (Ferencz and Hermes, 2008) and capybara (Lindert et al., 2008), but has not been found in *C. elegans* or yeast. Human and mouse MTF-1 show 93% sequence identity, the truncated C-terminus of the mouse notwithstanding. Vertebrate and insect MTF-1 show only 39% sequence similarity of the full-length protein, most of which is accounted for by a 78% similarity in the zinc-finger region.

Human MTF-1 harbors a non-canonical nuclear localization signal (NLS) sequence overlapping with the zinc-finger domain (Lindert et al., 2009) and a nuclear export

signal (NES) sequence that overlaps with the major activation domain (Saydam et al., 2001). MTF-1 also contains three different transactivation domains C-terminal to the DNA binding domain: an acidic, a proline-rich, and a serine/threonine –rich domain.




Figure 5: Schematic view of the functional domains of human-MTF-1 (top) and Drosophila-MTF-1 (below). Domains of human-MTF.1 include a potentiation domain (yellow), C₂H₂-zinc finger domain (blue), acidic activation domain (green), proline-rich domain (orange) and serine/threonine-rich domain (dark blue), auxiliary nuclear localization signal (aNLS) (black box on left), nuclear export signal (NES) (light green box), cysteine-cluster (red box). Drosophila-MTF-1 domains indicate the zinc-finger domain (blue), putative NLS (dark blue), acidic-domain (green), NES1 and 2 (light green), cys-cluster (red).

Targeted disruption of the MTF-1 gene has been achieved in the mouse and in *Drosophila melanogaster*. MTF-1 knockout mice die at the embryonic stage (E13-E14) due to liver degeneration (Gunes et al., 1998). *Drosophila* MTF-1 (dMTF-1) null mutants are viable but highly sensitive to heavy metal stress and also to copper starvation (Egli et al., 2003). dMTF-1 has the unique ability to handle both extremes, namely, copper overload and copper scarcity (Selvaraj et al., 2005). Upon copper stress, it activates the metallothionein genes, while under copper depletion condition it upregulates the copper importer Ctr1B.

Metallothioneins and other MTF-1 target genes

The best characterized target genes of MTF-1 are the metallothioneins. Metallothioneins are small proteins of 61-68 amino acids with a very high cysteine content (up to 30%) (West et al., 2008). Four metallothionein genes have been identified in the mouse. Mt-1 and Mt-2 are ubiquitously expressed and are highly inducible in response to various stress conditions (heavy metal ions, hormones, cytokines, ROS, inflammation). Mt-3 expression is restricted to the central nervous system and Mt-4 to the squamous epithelia, whereby the Mt3 gene contains MRE like sequence motifs but appears not to be regulated by MTF-1. Mice with a combined knockout of Mt-1 and Mt-2 are viable, in contrast to the MTF-1 knockout mice, however, they are sensitive to cadmium and zinc (Kelly et al., 1996; Masters et al., 1994; Michalska and Choo, 1993). In humans, MT-1 has diverged into a family of 7 genes (Mt-1A, -B, -E, -F, -G, -H and -X) while the other isoforms are represented by only one gene each, Mt-2A, Mt-3 and Mt-4 (Sadhu and Gedamu, 1988; Stennard et al., 1994; West et al., 1990). There is increasing evidence for a protective role of metallothioneins against neurodegeneration and CNS injury (West et al., 2008).

In *Drosophila melanogaster* there are at least four metallothioneins, designated MtnA-D, which are preferentially expressed in regions of high metal content, such as the larval intestinal tract and the midgut. They are induced by cadmium, zinc, copper, silver and mercury in an MTF-1 dependent manner (Egli et al., 2006). The quadruple knockout of the four Mtns leads to a severely reduced lifespan in males. Furthermore, flies are highly sensitive to copper, cadmium and to a lesser extent to zinc during development (Egli et al., 2006). Recently a putative fifth Mtn, MtnE, located proximal to MtnD was identified (Susan Celniker, Lawrence Berkeley National Laboratory) and is currently being investigated in our Lab (L. Atanesyan, H. Hua, O. Georgiev and WS, unpublished).

In fungi, metallothionein genes (CUP1 and CRS5 in *Saccharomyces cerevisiae*) are regulated by copper-sensitive transcription factors like Ace1 which have no sequence homology to MTF-1 (Rutherford and Bird, 2004). The metallothionein regulating transcription factor of *Candida glabrata* is Amt1 and that of *Yarrowia lipolytica* (dimorphic yeast) is Crf1. All these factors display a significant sequence similarity in their N-termini (Garcia et al., 2002).

Microarray studies done for the identification of novel target genes of MTF-1 in MTF-1 knockout mice, revealed α -fetoprotein, the liver-enriched transcription factor C/EBP α and tear lipocalin as new target genes, all of them being involved in stress response (Lichtlen and Schaffner, 2001). A zinc transporter, ZnT1 is also under MTF-1 control in the mouse (Langmade et al., 2000). As part of a screen carried out for zinc-responsive transcripts, Kruppel-like factor 4, hepatitis A virus cellular receptor 1 and complement factor B were also proposed as MTF-1 target genes (Kindermann et al., 2005). MTF-1 was shown to be essential for the basal expression of selenoprotein W (Sepw1), a putative antioxidant protein having a glutathione binding site (Wimmer et al., 2005). With regard to stress response, MTF-1 regulates cadmium-induced expression of N-myc downstream regulated gene 1 (Ndr1) as well as the cadmium response of the cysteine- and glycine-rich protein 1 gene (Csrp1) that is involved in cytoskeletal organization (Wimmer et al., 2005). MTF-1 downregulates metal importers, namely, the zinc transporter, Slc39a10 (ZIP10) (Kaler and Prasad, 2007) and in *D. melanogaster* the copper transporter Ctr1B. Additionally, transcription of the gene encoding hypoxia inducible placental growth factor, a pro-angiogenic protein, was shown to depend on MTF-1 (Green et al., 2001). Furthermore, siRNA knockdown experiments in cell culture suggested a cooperation of MTF-1 with Sp1 in the copper dependent positive regulation of the prion protein PrP gene (Bellingham et al., 2009). Misfolding of PrP causes transmissible spongiform encephalitis, a neurodegenerative disorder termed Creutzfeldt-Jakob disease in humans, BSE in cattle and scrapie in sheep. Growing evidence points towards a role of PrP in copper homeostasis (Brown et al., 1997; Pauly and Harris, 1998; Ruiz et al., 2000; Wadsworth et al., 1999). A search for MTF-1 target genes in *Drosophila* revealed results similar to mammals. MTF-1 targets include metallothioneins (MtnA-D), the zinc efflux transporter ZnT35C, the ATP binding cassette (ABC) transporter CG10505 and ferritins, which play a major role in iron homeostasis (Egli et al., 2003; Yepiskoposyan et al., 2006). Furthermore, dMTF-1 is implicated in the copper-responsive expression of DmAtp7, the *Drosophila* ortholog of the Menkes and Wilson's disease proteins Atp7A and Atp7B, in the larval midgut (Burke et al., 2008).

All these verified and inferred target genes highlight the central role of MTF-1 in metal homeostasis, defense against various stresses like heavy metal toxicity, heat shock, ROS and hypoxia as well as in angiogenesis and neuroprotection.

III) Zinc homeostasis

Zinc is an essential heavy metal that functions as a structural component in a number of proteins, including transcription factors and enzymes. Accordingly, zinc is important for several biological processes, such as nucleic acid metabolism, cell growth and proliferation, immune response and brain development. The concentrations of cellular zinc are tightly regulated. Uptake and efflux of zinc requires transporters which are grouped into two families based on their structure and functions. The ZIP protein family (also called as ZRT/IRT-related proteins) import zinc from extracellular space or intracellular vesicles into the cytoplasm (Eide, 2004). The ZnT proteins are involved in intracellular trafficking and excretion of zinc (Liuzzi and Cousins, 2004; Palmiter and Huang, 2004). In mammals, ZIPs are encoded by a solute carrier family 39A (SLC39A) genes and ZnTs (ZnT1-7) are encoded by solute carrier family 30A (SLC30A) genes (Palmiter and Findley, 1995). There are twelve ZIP genes identified so far in humans and three in mouse (Gaither and Eide, 2001). Under zinc-limiting conditions mice lacking ZIP1, ZIP2 or ZIP3 are embryonic lethal (Dufner-Beattie et al., 2006; Peters et al., 2007). Certain mutations in ZIP4 in humans cause a zinc deficiency disorder termed acrodermatitis enteropathica (AE) (Wang et al., 2002). Homozygous ZIP4 mutant mice die at early embryonic stages while heterozygous mutants are highly sensitive to zinc deficiency, similar to AE patients. Furthermore, female mice lacking functional ZnT4, which is expressed in breast epithelial cells, have low zinc levels in the milk which is fed to their pups, resulting in the death of the pups (Huang and Gitschier, 1997). In *Drosophila*, two ZIP genes have been identified, the *fear of intimacy foi* which is the counterpart of mammalian ZIP6/ZIP10 and *catsup* which is the counterpart of mammalian ZIP 7 (Mathews et al., 2005; Stathakis et al., 1999). One of the ZnTs, *ZnT35C* has been described in *Drosophila* (Yepiskoposyan et al., 2006). While *Drosophila* MTs preferentially bind copper and cadmium, mammalian MTs predominantly bind zinc, at least in non-metal stressed cells (Palmiter and Huang, 2004).

Zinc homeostasis in yeast is controlled primarily through the regulation of zinc uptake (Eide, 1997). *S. cerevisiae* has two separate uptake systems to obtain zinc from its environment, a high affinity transporter protein (Zrt1) and a lower affinity transporter (Zrt2), both of which are regulated by zinc availability; zinc limitation induces, and zinc-replete conditions repress their expression (Zhao et al., 1998). This zinc-responsive transcriptional regulation requires the regulatory protein Zap1, the function of which is repressed in high-zinc conditions (Zhao et al., 1998). Zap 1 binds to a conserved sequence, called the zinc response element (ZRE) of consensus 5'-ACCYYNAAGGT-3' which is present in the Zrt1, Zrt2, Zrt3 and ZAP1 promoters (Lyons et al., 2000).

References for Introduction

- Abeliovich, A., Schmitz, Y., Farinas, I., Choi-Lundberg, D., Ho, W. H., Castillo, P. E., Shinsky, N., Verdugo, J. M., Armanini, M., Ryan, A., *et al.* (2000). Mice lacking alpha-synuclein display functional deficits in the nigrostriatal dopamine system. *Neuron* 25, 239-252.
- Abou-Sleiman, P. M., Healy, D. G., Quinn, N., Lees, A. J., and Wood, N. W. (2003). The role of pathogenic DJ-1 mutations in Parkinson's disease. *Ann Neurol* 54, 283-286.
- Alam, Z. I., Jenner, A., Daniel, S. E., Lees, A. J., Cairns, N., Marsden, C. D., Jenner, P., and Halliwell, B. (1997). Oxidative DNA damage in the parkinsonian brain: an apparent selective increase in 8-hydroxyguanine levels in substantia nigra. *J Neurochem* 69, 1196-1203.
- Anderson, J. P., Walker, D. E., Goldstein, J. M., de Laat, R., Banducci, K., Caccavello, R. J., Barbour, R., Huang, J., Kling, K., Lee, M., *et al.* (2006). Phosphorylation of Ser-129 is the dominant pathological modification of alpha-synuclein in familial and sporadic Lewy body disease. *J Biol Chem* 281, 29739-29752.
- Arawaka, S., Wada, M., Goto, S., Karube, H., Sakamoto, M., Ren, C. H., Koyama, S., Nagasawa, H., Kimura, H., Kawanami, T., *et al.* (2006). The role of G-protein-coupled receptor kinase 5 in pathogenesis of sporadic Parkinson's disease. *J Neurosci* 26, 9227-9238.
- Ardley, H. C., Scott, G. B., Rose, S. A., Tan, N. G., and Robinson, P. A. (2004). UCH-L1 aggresome formation in response to proteasome impairment indicates a role in inclusion formation in Parkinson's disease. *J Neurochem* 90, 379-391.
- Auf der Maur, A., Belser, T., Elgar, G., Georgiev, O., and Schaffner, W. (1999). Characterization of the transcription factor MTF-1 from the Japanese pufferfish (*Fugu rubripes*) reveals evolutionary conservation of heavy metal stress response. *Biol Chem* 380, 175-185.
- Auluck, P. K., and Bonini, N. M. (2002). Pharmacological prevention of Parkinson disease in *Drosophila*. *Nat Med* 8, 1185-1186.
- Bae, Y. J., Park, K. S., and Kang, S. J. (2003). Genomic organization and expression of parkin in *Drosophila melanogaster*. *Exp Mol Med* 35, 393-402.
- Baptista, M. J., O'Farrell, C., Daya, S., Ahmad, R., Miller, D. W., Hardy, J., Farrer, M. J., and Cookson, M. R. (2003). Co-ordinate transcriptional regulation of dopamine synthesis genes by alpha-synuclein in human neuroblastoma cell lines. *J Neurochem* 85, 957-968.
- Beites, C. L., Xie, H., Bowser, R., and Trimble, W. S. (1999). The septin CDCrel-1 binds syntaxin and inhibits exocytosis. *Nat Neurosci* 2, 434-439.
- Bellingham, S. A., Coleman, L. A., Masters, C. L., Camakaris, J., and Hill, A. F. (2009). Regulation of prion gene expression by transcription factors SP1 and metal transcription factor-1. *J Biol Chem* 284, 1291-1301.
- Bittel, D., Dalton, T., Samson, S. L., Gedamu, L., and Andrews, G. K. (1998). The DNA binding activity of metal response element-binding transcription factor-1 is activated in vivo and in vitro by zinc, but not by other transition metals. *J Biol Chem* 273, 7127-7133.
- Bonifati, V., Rizzu, P., van Baren, M. J., Schaap, O., Breedveld, G. J., Krieger, E., Dekker, M. C., Squitieri, F., Ibanez, P., Joosse, M., *et al.* (2003). Mutations in the DJ-

1 gene associated with autosomal recessive early-onset parkinsonism. *Science* 299, 256-259.

Brice, A. (2005). How much does dardarin contribute to Parkinson's disease? *Lancet* 365, 363-364.

Brown, D. R., Qin, K., Herms, J. W., Madlung, A., Manson, J., Strome, R., Fraser, P. E., Kruck, T., von Bohlen, A., Schulz-Schaeffer, W., *et al.* (1997). The cellular prion protein binds copper in vivo. *Nature* 390, 684-687.

Brugnera, E., Georgiev, O., Radtke, F., Heuchel, R., Baker, E., Sutherland, G. R., and Schaffner, W. (1994). Cloning, chromosomal mapping and characterization of the human metal-regulatory transcription factor MTF-1. *Nucleic Acids Res* 22, 3167-3173.

Burke, R., Commons, E., and Camakaris, J. (2008). Expression and localisation of the essential copper transporter DmATP7 in *Drosophila* neuronal and intestinal tissues. *Int J Biochem Cell Biol* 40, 1850-1860.

Bushnell, D. M., and Martin, M. L. (1999). Quality of life and Parkinson's disease: translation and validation of the US Parkinson's Disease Questionnaire (PDQ-39). *Qual Life Res* 8, 345-350.

Caughey, B., and Lansbury, P. T. (2003). Protofibrils, pores, fibrils, and neurodegeneration: separating the responsible protein aggregates from the innocent bystanders. *Annu Rev Neurosci* 26, 267-298.

Chen, W. Y., John, J. A., Lin, C. H., and Chang, C. Y. (2002). Molecular cloning and developmental expression of zinc finger transcription factor MTF-1 gene in zebrafish, *Danio rerio*. *Biochem Biophys Res Commun* 291, 798-805.

Choi, P., Snyder, H., Petrucelli, L., Theisler, C., Chong, M., Zhang, Y., Lim, K., Chung, K. K., Kehoe, K., D'Adamio, L., *et al.* (2003). SEPT5_v2 is a parkin-binding protein. *Brain Res Mol Brain Res* 117, 179-189.

Chung, K., Dawson, V. L., and Dawson, T. M. (2003). New insights into Parkinson's disease. *J Neurol* 250, III15-24.

Chung, K. K., Zhang, Y., Lim, K. L., Tanaka, Y., Huang, H., Gao, J., Ross, C. A., Dawson, V. L., and Dawson, T. M. (2001). Parkin ubiquitinates the alpha-synuclein-interacting protein, synphilin-1: implications for Lewy-body formation in Parkinson disease. *Nat Med* 7, 1144-1150.

Clark, I. E., Dodson, M. W., Jiang, C., Cao, J. H., Huh, J. R., Seol, J. H., Yoo, S. J., Hay, B. A., and Guo, M. (2006). *Drosophila* pink1 is required for mitochondrial function and interacts genetically with parkin. *Nature* 441, 1162-1166.

Clayton, D. F., and George, J. M. (1998). The synucleins: a family of proteins involved in synaptic function, plasticity, neurodegeneration and disease. *Trends Neurosci* 21, 249-254.

Clayton, D. F., and George, J. M. (1999). Synucleins in synaptic plasticity and neurodegenerative disorders. *J Neurosci Res* 58, 120-129.

Conway, K. A., Harper, J. D., and Lansbury, P. T. (1998). Accelerated in vitro fibril formation by a mutant alpha-synuclein linked to early-onset Parkinson disease. *Nat Med* 4, 1318-1320.

Conway, K. A., Rochet, J. C., Bieganski, R. M., and Lansbury, P. T., Jr. (2001). Kinetic stabilization of the alpha-synuclein protofibril by a dopamine-alpha-synuclein adduct. *Science* 294, 1346-1349.

Corti, O., Hampe, C., Koutnikova, H., Darios, F., Jacquier, S., Prigent, A., Robinson, J. C., Pradier, L., Ruberg, M., Mirande, M., *et al.* (2003). The p38 subunit of the aminoacyl-tRNA synthetase complex is a Parkin substrate: linking protein biosynthesis and neurodegeneration. *Hum Mol Genet* 12, 1427-1437.

Cramer, M., Nagy, I., Murphy, B. J., Gassmann, M., Hottiger, M. O., Georgiev, O., and Schaffner, W. (2005). NF-kappaB contributes to transcription of placenta growth factor and interacts with metal responsive transcription factor-1 in hypoxic human cells. *Biol Chem* 386, 865-872.

Culetto, E., and Sattelle, D. B. (2000). A role for *Caenorhabditis elegans* in understanding the function and interactions of human disease genes. *Hum Mol Genet* 9, 869-877.

Dalle-Donne, I., Aldini, G., Carini, M., Colombo, R., Rossi, R., and Milzani, A. (2006). Protein carbonylation, cellular dysfunction, and disease progression. *J Cell Mol Med* 10, 389-406.

Dalton, T. P., Bittel, D., and Andrews, G. K. (1997). Reversible activation of mouse metal response element-binding transcription factor 1 DNA binding involves zinc interaction with the zinc finger domain. *Mol Cell Biol* 17, 2781-2789.

Dauer, W., and Przedborski, S. (2003). Parkinson's disease: mechanisms and models. *Neuron* 39, 889-909.

Davidson, W. S., Jonas, A., Clayton, D. F., and George, J. M. (1998). Stabilization of alpha-synuclein secondary structure upon binding to synthetic membranes. *J Biol Chem* 273, 9443-9449.

Dawson, T. M. (2000). New animal models for Parkinson's disease. *Cell* 101, 115-118.

Dawson, T. M., and Dawson, V. L. (2003). Molecular pathways of neurodegeneration in Parkinson's disease. *Science* 302, 819-822.

de Silva, H. R., Khan, N. L., and Wood, N. W. (2000). The genetics of Parkinson's disease. *Curr Opin Genet Dev* 10, 292-298.

Ding, T. T., Lee, S. J., Rochet, J. C., and Lansbury, P. T., Jr. (2002). Annular alpha-synuclein protofibrils are produced when spherical protofibrils are incubated in solution or bound to brain-derived membranes. *Biochemistry* 41, 10209-10217.

Dufner-Beattie, J., Huang, Z. L., Geiser, J., Xu, W., and Andrews, G. K. (2006). Mouse ZIP1 and ZIP3 genes together are essential for adaptation to dietary zinc deficiency during pregnancy. *Genesis* 44, 239-251.

Dunnett, S. B., and Bjorklund, A. (1999). Prospects for new restorative and neuroprotective treatments in Parkinson's disease. *Nature* 399, A32-39.

Egli, D., Selvaraj, A., Yepiskoposyan, H., Zhang, B., Hafen, E., Georgiev, O., and Schaffner, W. (2003). Knockout of 'metal-responsive transcription factor' MTF-1 in *Drosophila* by homologous recombination reveals its central role in heavy metal homeostasis. *Embo J* 22, 100-108.

Egli, D., Yepiskoposyan, H., Selvaraj, A., Balamurugan, K., Rajaram, R., Simons, A., Multhaup, G., Mettler, S., Vardanyan, A., Georgiev, O., and Schaffner, W. (2006). A family knockout of all four *Drosophila* metallothioneins reveals a central role in copper homeostasis and detoxification. *Mol Cell Biol* 26, 2286-2296.

Eide, D. (1997). Molecular biology of iron and zinc uptake in eukaryotes. *Curr Opin Cell Biol* 9, 573-577.

Eide, D. J. (2004). The SLC39 family of metal ion transporters. *Pflugers Arch* 447, 796-800.

Engelender, S., Kaminsky, Z., Guo, X., Sharp, A. H., Amaravi, R. K., Kleiderlein, J. J., Margolis, R. L., Troncoso, J. C., Lanahan, A. A., Worley, P. F., *et al.* (1999). Synphilin-1 associates with alpha-synuclein and promotes the formation of cytosolic inclusions. *Nat Genet* 22, 110-114.

Fallon, L., Moreau, F., Croft, B. G., Labib, N., Gu, W. J., and Fon, E. A. (2002). Parkin and CASK/LIN-2 associate via a PDZ-mediated interaction and are co-localized in lipid rafts and postsynaptic densities in brain. *J Biol Chem* 277, 486-491.

Farrer, M., Chan, P., Chen, R., Tan, L., Lincoln, S., Hernandez, D., Forno, L., Gwinn-Hardy, K., Petrucelli, L., Hussey, J., *et al.* (2001). Lewy bodies and parkinsonism in families with parkin mutations. *Ann Neurol* 50, 293-300.

Feany, M. B., and Bender, W. W. (2000). A *Drosophila* model of Parkinson's disease. *Nature* 404, 394-398.

Feng, J., and Lin, Y. (2001). Parkin interacts with MAP1A and promotes its ubiquitination. *Soc Neurosci Abstr* 27, 10.

Ferencz, A., and Hermes, E. (2008). Identification and characterization of two mtf-1 genes in common carp. *Comp Biochem Physiol C Toxicol Pharmacol* 148, 238-243.

Fernagut, P. O., and Chesselet, M. F. (2004). Alpha-synuclein and transgenic mouse models. *Neurobiol Dis* 17, 123-130.

Field, C. M., and Kellogg, D. (1999). Septins: cytoskeletal polymers or signalling GTPases? *Trends Cell Biol* 9, 387-394.

Finney, N., Walther, F., Mantel, P. Y., Stauffer, D., Rovelli, G., and Dev, K. K. (2003). The cellular protein level of parkin is regulated by its ubiquitin-like domain. *J Biol Chem* 278, 16054-16058.

Foley, P., and Riederer, P. (2000). Influence of neurotoxins and oxidative stress on the onset and progression of Parkinson's disease. *J Neurol* 247 Suppl 2, II82-94.

Gaither, L. A., and Eide, D. J. (2001). Eukaryotic zinc transporters and their regulation. *Biometals* 14, 251-270.

Galvin, J. E., Lee, V. M., and Trojanowski, J. Q. (2001). Synucleinopathies: clinical and pathological implications. *Arch Neurol* 58, 186-190.

Garcia, S., Prado, M., Degano, R., and Dominguez, A. (2002). A copper-responsive transcription factor, CRF1, mediates copper and cadmium resistance in *Yarrowia lipolytica*. *J Biol Chem* 277, 37359-37368.

Gasser, T. (2001). Genetics of Parkinson's disease. *J Neurol* 248, 833-840.

Gelb, D. J., Oliver, E., and Gilman, S. (1999). Diagnostic criteria for Parkinson disease. *Arch Neurol* 56, 33-39.

Goedert, M. (1999). Filamentous nerve cell inclusions in neurodegenerative diseases: tauopathies and alpha-synucleinopathies. *Philos Trans R Soc Lond B Biol Sci* 354, 1101-1118.

Goedert, M., and Spillantini, M. G. (1998). Lewy body diseases and multiple system atrophy as alpha-synucleinopathies. *Mol Psychiatry* 3, 462-465.

Goldberg, M. S., Fleming, S. M., Palacino, J. J., Cepeda, C., Lam, H. A., Bhatnagar, A., Meloni, E. G., Wu, N., Ackerson, L. C., Klapstein, G. J., *et al.* (2003). Parkin-deficient mice exhibit nigrostriatal deficits but not loss of dopaminergic neurons. *J Biol Chem* 278, 43628-43635.

Goldberg, M. S., Pisani, A., Haburcak, M., Vortherms, T. A., Kitada, T., Costa, C., Tong, Y., Martella, G., Tschertter, A., Martins, A., *et al.* (2005). Nigrostriatal dopaminergic deficits and hypokinesia caused by inactivation of the familial Parkinsonism-linked gene DJ-1. *Neuron* 45, 489-496.

Graham, D. G. (1978). Oxidative pathways for catecholamines in the genesis of neuromelanin and cytotoxic quinones. *Mol Pharmacol* 14, 633-643.

Green, C. J., Lichtlen, P., Huynh, N. T., Yanovsky, M., Laderoute, K. R., Schaffner, W., and Murphy, B. J. (2001). Placenta growth factor gene expression is induced by hypoxia in fibroblasts: a central role for metal transcription factor-1. *Cancer Res* 61, 2696-2703.

Greene, J. C., Whitworth, A. J., Andrews, L. A., Parker, T. J., and Pallanck, L. J. (2005). Genetic and genomic studies of *Drosophila parkin* mutants implicate oxidative stress and innate immune responses in pathogenesis. *Hum Mol Genet* 14, 799-811.

Greene, J. C., Whitworth, A. J., Kuo, I., Andrews, L. A., Feany, M. B., and Pallanck, L. J. (2003). Mitochondrial pathology and apoptotic muscle degeneration in *Drosophila parkin* mutants. *Proc Natl Acad Sci U S A* 100, 4078-4083.

Gregersen, N., Bross, P., Vang, S., and Christensen, J. H. (2006). Protein misfolding and human disease. *Annu Rev Genomics Hum Genet* 7, 103-124.

Gunes, C., Heuchel, R., Georgiev, O., Muller, K. H., Lichtlen, P., Bluthmann, H., Marino, S., Aguzzi, A., and Schaffner, W. (1998). Embryonic lethality and liver degeneration in mice lacking the metal-responsive transcriptional activator MTF-1. *Embo J* 17, 2846-2854.

Hague, S., Rogaeva, E., Hernandez, D., Gulick, C., Singleton, A., Hanson, M., Johnson, J., Weiser, R., Gallardo, M., Ravina, B., *et al.* (2003). Early-onset Parkinson's disease caused by a compound heterozygous DJ-1 mutation. *Ann Neurol* 54, 271-274.

Hamamichi, S., Rivas, R. N., Knight, A. L., Cao, S., Caldwell, K. A., and Caldwell, G. A. (2008). Hypothesis-based RNAi screening identifies neuroprotective genes in a Parkinson's disease model. *Proc Natl Acad Sci U S A* 105, 728-733.

Han, H., Weinreb, P. H., and Lansbury, P. T., Jr. (1995). The core Alzheimer's peptide NAC forms amyloid fibrils which seed and are seeded by beta-amyloid: is NAC a common trigger or target in neurodegenerative disease? *Chem Biol* 2, 163-169.

Harding, H. P., Zhang, Y., and Ron, D. (1999). Protein translation and folding are coupled by an endoplasmic-reticulum-resident kinase. *Nature* 397, 271-274.

Hedrich, K., Djarmati, A., Schafer, N., Hering, R., Wellenbrock, C., Weiss, P. H., Hilker, R., Vieregge, P., Ozelius, L. J., Heutink, P., *et al.* (2004). DJ-1 (PARK7) mutations are less frequent than Parkin (PARK2) mutations in early-onset Parkinson disease. *Neurology* 62, 389-394.

Hersch, S. M., Yi, H., Heilman, C. J., Edwards, R. H., and Levey, A. I. (1997). Subcellular localization and molecular topology of the dopamine transporter in the striatum and substantia nigra. *J Comp Neurol* 388, 211-227.

Hod, Y., Pentyala, S. N., Whyard, T. C., and El-Maghrabi, M. R. (1999). Identification and characterization of a novel protein that regulates RNA-protein interaction. *J Cell Biochem* 72, 435-444.

Hori, O., Ichinoda, F., Tamatani, T., Yamaguchi, A., Sato, N., Ozawa, K., Kitao, Y., Miyazaki, M., Harding, H. P., Ron, D., *et al.* (2002). Transmission of cell stress from endoplasmic reticulum to mitochondria: enhanced expression of Lon protease. *J Cell Biol* 157, 1151-1160.

Huang, L., and Gitschier, J. (1997). A novel gene involved in zinc transport is deficient in the lethal milk mouse. *Nat Genet* 17, 292-297.

Huang, Y., Cheung, L., Rowe, D., and Halliday, G. (2004). Genetic contributions to Parkinson's disease. *Brain Res Brain Res Rev* 46, 44-70.

Huynh, D. P., Scoles, D. R., Nguyen, D., and Pulst, S. M. (2003). The autosomal recessive juvenile Parkinson disease gene product, parkin, interacts with and ubiquitinates synaptotagmin XI. *Hum Mol Genet* 12, 2587-2597.

Imai, Y., Soda, M., Inoue, H., Hattori, N., Mizuno, Y., and Takahashi, R. (2001). An unfolded putative transmembrane polypeptide, which can lead to endoplasmic reticulum stress, is a substrate of Parkin. *Cell* 105, 891-902.

Imai, Y., Soda, M., Murakami, T., Shoji, M., Abe, K., and Takahashi, R. (2003). A product of the human gene adjacent to parkin is a component of Lewy bodies and suppresses Pael receptor-induced cell death. *J Biol Chem* 278, 51901-51910.

Irizarry, M. C., Kim, T. W., McNamara, M., Tanzi, R. E., George, J. M., Clayton, D. F., and Hyman, B. T. (1996). Characterization of the precursor protein of the non-A beta component of senile plaques (NACP) in the human central nervous system. *J Neuropathol Exp Neurol* 55, 889-895.

Ishizawa, T., Mattila, P., Davies, P., Wang, D., and Dickson, D. W. (2003). Colocalization of tau and alpha-synuclein epitopes in Lewy bodies. *J Neuropathol Exp Neurol* 62, 389-397.

Itier, J. M., Ibanez, P., Mena, M. A., Abbas, N., Cohen-Salmon, C., Bohme, G. A., Laville, M., Pratt, J., Corti, O., Pradier, L., *et al.* (2003). Parkin gene inactivation alters behaviour and dopamine neurotransmission in the mouse. *Hum Mol Genet* 12, 2277-2291.

Iwai, A. (2000). Properties of NACP/alpha-synuclein and its role in Alzheimer's disease. *Biochim Biophys Acta* 1502, 95-109.

Iwatsubo, T. (2003). Aggregation of alpha-synuclein in the pathogenesis of Parkinson's disease. *J Neurol* 250 Suppl 3, III11-14.

Jakes, R., Spillantini, M. G., and Goedert, M. (1994). Identification of two distinct synucleins from human brain. *FEBS Lett* 345, 27-32.

Jensen, P. H., Nielsen, M. S., Jakes, R., Dotti, C. G., and Goedert, M. (1998). Binding of alpha-synuclein to brain vesicles is abolished by familial Parkinson's disease mutation. *J Biol Chem* 273, 26292-26294.

Jiang, H., Jiang, Q., and Feng, J. (2004). Parkin increases dopamine uptake by enhancing the cell surface expression of dopamine transporter. *J Biol Chem* 279, 54380-54386.

Jo, E., McLaurin, J., Yip, C. M., St George-Hyslop, P., and Fraser, P. E. (2000). alpha-Synuclein membrane interactions and lipid specificity. *J Biol Chem* 275, 34328-34334.

Joazeiro, C. A., and Weissman, A. M. (2000). RING finger proteins: mediators of ubiquitin ligase activity. *Cell* 102, 549-552.

Kahle, P. J., Leimer, U., and Haass, C. (2000a). Does failure of parkin-mediated ubiquitination cause juvenile parkinsonism? *Trends Biochem Sci* 25, 524-527.

Kahle, P. J., Neumann, M., Ozmen, L., Muller, V., Jacobsen, H., Schindzielorz, A., Okochi, M., Leimer, U., van Der Putten, H., Probst, A., *et al.* (2000b). Subcellular localization of wild-type and Parkinson's disease-associated mutant alpha -synuclein in human and transgenic mouse brain. *J Neurosci* 20, 6365-6373.

Kaler, P., and Prasad, R. (2007). Molecular cloning and functional characterization of novel zinc transporter rZip10 (Slc39a10) involved in zinc uptake across rat renal brush-border membrane. *Am J Physiol Renal Physiol* 292, F217-229.

Kaufman, R. J. (1999). Stress signaling from the lumen of the endoplasmic reticulum: coordination of gene transcriptional and translational controls. *Genes Dev* 13, 1211-1233.

Kelly, E. J., Quaife, C. J., Froelick, G. J., and Palmiter, R. D. (1996). Metallothionein I and II protect against zinc deficiency and zinc toxicity in mice. *J Nutr* 126, 1782-1790.

Kim, R. H., Smith, P. D., Aleyasin, H., Hayley, S., Mount, M. P., Pownall, S., Wakeham, A., You-Ten, A. J., Kalia, S. K., Horne, P., *et al.* (2005). Hypersensitivity of DJ-1-deficient mice to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) and oxidative stress. *Proc Natl Acad Sci U S A* 102, 5215-5220.

Kim, T. D., Paik, S. R., Yang, C. H., and Kim, J. (2000). Structural changes in alpha-synuclein affect its chaperone-like activity in vitro. *Protein Sci* 9, 2489-2496.

Kindermann, B., Doring, F., Budczies, J., and Daniel, H. (2005). Zinc-sensitive genes as potential new target genes of the metal transcription factor-1 (MTF-1). *Biochem Cell Biol* 83, 221-229.

Kirik, D., Annett, L. E., Burger, C., Muzyczka, N., Mandel, R. J., and Bjorklund, A. (2003). Nigrostriatal alpha-synucleinopathy induced by viral vector-mediated overexpression of human alpha-synuclein: a new primate model of Parkinson's disease. *Proc Natl Acad Sci U S A* 100, 2884-2889.

Kitada, T., Asakawa, S., Hattori, N., Matsumine, H., Yamamura, Y., Minoshima, S., Yokochi, M., Mizuno, Y., and Shimizu, N. (1998). Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* 392, 605-608.

Kitada, T., Pisani, A., Porter, D. R., Yamaguchi, H., Tscherter, A., Martella, G., Bonsi, P., Zhang, C., Pothos, E. N., and Shen, J. (2007). Impaired dopamine release and synaptic plasticity in the striatum of PINK1-deficient mice. *Proc Natl Acad Sci U S A* 104, 11441-11446.

Klein, C., Liu, L., Doheny, D., Kock, N., Muller, B., de Carvalho Aguiar, P., Leung, J., de Leon, D., Bressman, S. B., Silverman, J., *et al.* (2002). Epsilon-sarcoglycan mutations found in combination with other dystonia gene mutations. *Ann Neurol* 52, 675-679.

Klein, C., Pramstaller, P. P., Kis, B., Page, C. C., Kann, M., Leung, J., Woodward, H., Castellan, C. C., Scherer, M., Vieregge, P., *et al.* (2000). Parkin deletions in a family with adult-onset, tremor-dominant parkinsonism: expanding the phenotype. *Ann Neurol* 48, 65-71.

Kosel, S., Hofhaus, G., Maassen, A., Vieregge, P., and Graeber, M. B. (1999). Role of mitochondria in Parkinson disease. *Biol Chem* 380, 865-870.

Kruger, R., Kuhn, W., Muller, T., Woitalla, D., Graeber, M., Kosel, S., Przuntek, H., Epplen, J. T., Schols, L., and Riess, O. (1998). Ala30Pro mutation in the gene encoding alpha-synuclein in Parkinson's disease. *Nat Genet* 18, 106-108.

Kubo, S. I., Kitami, T., Noda, S., Shimura, H., Uchiyama, Y., Asakawa, S., Minoshima, S., Shimizu, N., Mizuno, Y., and Hattori, N. (2001). Parkin is associated with cellular vesicles. *J Neurochem* 78, 42-54.

Langmade, S. J., Ravindra, R., Daniels, P. J., and Andrews, G. K. (2000). The transcription factor MTF-1 mediates metal regulation of the mouse ZnT1 gene. *J Biol Chem* 275, 34803-34809.

Lansbury, P. T., Jr., and Brice, A. (2002). Genetics of Parkinson's disease and biochemical studies of implicated gene products. *Curr Opin Cell Biol* 14, 653-660.

Larsen, C. N., Krantz, B. A., and Wilkinson, K. D. (1998). Substrate specificity of deubiquitinating enzymes: ubiquitin C-terminal hydrolases. *Biochemistry* 37, 3358-3368.

Larsen, C. N., Price, J. S., and Wilkinson, K. D. (1996). Substrate binding and catalysis by ubiquitin C-terminal hydrolases: identification of two active site residues. *Biochemistry* 35, 6735-6744.

Lashuel, H. A., Petre, B. M., Wall, J., Simon, M., Nowak, R. J., Walz, T., and Lansbury, P. T., Jr. (2002). Alpha-synuclein, especially the Parkinson's disease-associated mutants, forms pore-like annular and tubular protofibrils. *J Mol Biol* 322, 1089-1102.

Lee, F. J., Liu, F., Pristupa, Z. B., and Niznik, H. B. (2001). Direct binding and functional coupling of alpha-synuclein to the dopamine transporters accelerate dopamine-induced apoptosis. *Faseb J* 15, 916-926.

Leroy, E., Boyer, R., Auburger, G., Leube, B., Ulm, G., Mezey, E., Harta, G., Brownstein, M. J., Jonnalagada, S., Chernova, T., *et al.* (1998). The ubiquitin pathway in Parkinson's disease. *Nature* 395, 451-452.

Levecque, C., Destee, A., Mouroux, V., Becquet, E., Defebvre, L., Amouyel, P., and Chartier-Harlin, M. C. (2001). No genetic association of the ubiquitin carboxy-terminal hydrolase-L1 gene S18Y polymorphism with familial Parkinson's disease. *J Neural Transm* 108, 979-984.

Li, W., West, N., Colla, E., Pletnikova, O., Troncoso, J. C., Marsh, L., Dawson, T. M., Jakala, P., Hartmann, T., Price, D. L., and Lee, M. K. (2005). Aggregation promoting C-terminal truncation of alpha-synuclein is a normal cellular process and is enhanced by the familial Parkinson's disease-linked mutations. *Proc Natl Acad Sci U S A* 102, 2162-2167.

Lichtlen, P., and Schaffner, W. (2001). Putting its fingers on stressful situations: the heavy metal-regulatory transcription factor MTF-1. *Bioessays* 23, 1010-1017.

Lim, K. L., Chew, K. C., Tan, J. M., Wang, C., Chung, K. K., Zhang, Y., Tanaka, Y., Smith, W., Engelender, S., Ross, C. A., *et al.* (2005). Parkin mediates nonclassical, proteasomal-independent ubiquitination of synphilin-1: implications for Lewy body formation. *J Neurosci* 25, 2002-2009.

Lindert, U., Cramer, M., Meuli, M., Georgiev, O., and Schaffner, W. (2009). Metal-responsive transcription factor 1 (MTF-1) activity is regulated by a nonconventional nuclear localization signal and a metal-responsive transactivation domain. *Mol Cell Biol* 29, 6283-6293.

Lindert, U., Leuzinger, L., Steiner, K., Georgiev, O., and Schaffner, W. (2008). Characterization of metal-responsive transcription factor (MTF-1) from the giant rodent capybara reveals features in common with human as well as with small rodents (mouse, rat). Short communication. *Chem Biodivers* 5, 1485-1494.

Liu, Y., Fallon, L., Lashuel, H. A., Liu, Z., and Lansbury, P. T., Jr. (2002). The UCH-L1 gene encodes two opposing enzymatic activities that affect alpha-synuclein degradation and Parkinson's disease susceptibility. *Cell* 111, 209-218.

Liuzzi, J. P., and Cousins, R. J. (2004). Mammalian zinc transporters. *Annu Rev Nutr* 24, 151-172.

Lo Bianco, C., Ridet, J. L., Schneider, B. L., Deglon, N., and Aebischer, P. (2002). alpha-Synucleinopathy and selective dopaminergic neuron loss in a rat lentiviral-based model of Parkinson's disease. *Proc Natl Acad Sci U S A* 99, 10813-10818.

Lo Bianco, C., Schneider, B. L., Bauer, M., Sajadi, A., Brice, A., Iwatsubo, T., and Aebischer, P. (2004). Lentiviral vector delivery of parkin prevents dopaminergic degeneration in an alpha-synuclein rat model of Parkinson's disease. *Proc Natl Acad Sci U S A* 101, 17510-17515.

Lohmann, E., Periquet, M., Bonifati, V., Wood, N. W., De Michele, G., Bonnet, A. M., Fraix, V., Broussolle, E., Horstink, M. W., Vidailhet, M., *et al.* (2003). How much phenotypic variation can be attributed to parkin genotype? *Ann Neurol* 54, 176-185.

Lotharius, J., and Brundin, P. (2002). Pathogenesis of Parkinson's disease: dopamine, vesicles and alpha-synuclein. *Nat Rev Neurosci* 3, 932-942.

Lowe, J., McDermott, H., Landon, M., Mayer, R. J., and Wilkinson, K. D. (1990). Ubiquitin carboxyl-terminal hydrolase (PGP 9.5) is selectively present in ubiquitinated inclusion bodies characteristic of human neurodegenerative diseases. *J Pathol* 161, 153-160.

Lyons, T. J., Gasch, A. P., Gaither, L. A., Botstein, D., Brown, P. O., and Eide, D. J. (2000). Genome-wide characterization of the Zap1p zinc-responsive regulon in yeast. *Proc Natl Acad Sci U S A* 97, 7957-7962.

Macedo, M. G., Anar, B., Bronner, I. F., Cannella, M., Squitieri, F., Bonifati, V., Hoogeveen, A., Heutink, P., and Rizzu, P. (2003). The DJ-1L166P mutant protein associated with early onset Parkinson's disease is unstable and forms higher-order protein complexes. *Hum Mol Genet* 12, 2807-2816.

Manetto, V., Perry, G., Tabaton, M., Mulvihill, P., Fried, V. A., Smith, H. T., Gambetti, P., and Autilio-Gambetti, L. (1988). Ubiquitin is associated with abnormal cytoplasmic filaments characteristic of neurodegenerative diseases. *Proc Natl Acad Sci U S A* 85, 4501-4505.

Maraganore, D. M., Farrer, M. J., Hardy, J. A., Lincoln, S. J., McDonnell, S. K., and Rocca, W. A. (1999). Case-control study of the ubiquitin carboxy-terminal hydrolase L1 gene in Parkinson's disease. *Neurology* 53, 1858-1860.

Martin, E. D., Gonzalez-Garcia, C., Milan, M., Farinas, I., and Cena, V. (2004). Stressor-related impairment of synaptic transmission in hippocampal slices from alpha-synuclein knockout mice. *Eur J Neurosci* 20, 3085-3091.

Masters, B. A., Kelly, E. J., Quaife, C. J., Brinster, R. L., and Palmiter, R. D. (1994). Targeted disruption of metallothionein I and II genes increases sensitivity to cadmium. *Proc Natl Acad Sci U S A* 91, 584-588.

Mathews, W. R., Wang, F., Eide, D. J., and Van Doren, M. (2005). *Drosophila* fear of intimacy encodes a Zrt/IRT-like protein (ZIP) family zinc transporter functionally related to mammalian ZIP proteins. *J Biol Chem* 280, 787-795.

Matsumine, H. (1999). [A Parkin gene (PARK 2) and Parkinson's disease]. *Rinsho Shinkeigaku* 39, 9-12.

McNaught, K. S., Belizaire, R., Isacson, O., Jenner, P., and Olanow, C. W. (2003). Altered proteasomal function in sporadic Parkinson's disease. *Exp Neurol* 179, 38-46.

Menzies, F. M., Yenissetti, S. C., and Min, K. T. (2005). Roles of *Drosophila* DJ-1 in survival of dopaminergic neurons and oxidative stress. *Curr Biol* 15, 1578-1582.

Meulener, M., Whitworth, A. J., Armstrong-Gold, C. E., Rizzu, P., Heutink, P., Wes, P. D., Pallanck, L. J., and Bonini, N. M. (2005). *Drosophila* DJ-1 mutants are selectively sensitive to environmental toxins associated with Parkinson's disease. *Curr Biol* 15, 1572-1577.

Michalska, A. E., and Choo, K. H. (1993). Targeting and germ-line transmission of a null mutation at the metallothionein I and II loci in mouse. *Proc Natl Acad Sci U S A* 90, 8088-8092.

Mizuno, Y., Hattori, N., Mori, H., Suzuki, T., and Tanaka, K. (2001). Parkin and Parkinson's disease. *Curr Opin Neurol* 14, 477-482.

Moore, D. J., Zhang, L., Dawson, T. M., and Dawson, V. L. (2003). A missense mutation (L166P) in DJ-1, linked to familial Parkinson's disease, confers reduced protein stability and impairs homo-oligomerization. *J Neurochem* 87, 1558-1567.

Mori, H., Kondo, T., Yokochi, M., Matsumine, H., Nakagawa-Hattori, Y., Miyake, T., Suda, K., and Mizuno, Y. (1998). Pathologic and biochemical studies of juvenile parkinsonism linked to chromosome 6q. *Neurology* 51, 890-892.

Mori, K. (2000). Tripartite management of unfolded proteins in the endoplasmic reticulum. *Cell* 101, 451-454.

Munch, G., Gerlach, M., Sian, J., Wong, A., and Riederer, P. (1998). Advanced glycation end products in neurodegeneration: more than early markers of oxidative stress? *Ann Neurol* 44, S85-88.

Murakami, T., Shoji, M., Imai, Y., Inoue, H., Kawarabayashi, T., Matsubara, E., Harigaya, Y., Sasaki, A., Takahashi, R., and Abe, K. (2004). Pael-R is accumulated in Lewy bodies of Parkinson's disease. *Ann Neurol* 55, 439-442.

Nagakubo, D., Taira, T., Kitaura, H., Ikeda, M., Tamai, K., Iguchi-Ariga, S. M., and Ariga, H. (1997). DJ-1, a novel oncogene which transforms mouse NIH3T3 cells in cooperation with ras. *Biochem Biophys Res Commun* 231, 509-513.

Nakamura, K., Bindokas, V. P., Kowlessur, D., Elas, M., Milstien, S., Marks, J. D., Halpern, H. J., and Kang, U. J. (2001). Tetrahydrobiopterin scavenges superoxide in dopaminergic neurons. *J Biol Chem* 276, 34402-34407.

Olanow, C. W., and Tatton, W. G. (1999). Etiology and pathogenesis of Parkinson's disease. *Annu Rev Neurosci* 22, 123-144.

Ostrerova, N., Petrucelli, L., Farrer, M., Mehta, N., Choi, P., Hardy, J., and Wolozin, B. (1999). alpha-Synuclein shares physical and functional homology with 14-3-3 proteins. *J Neurosci* 19, 5782-5791.

Paisan-Ruiz, C., Jain, S., Evans, E. W., Gilks, W. P., Simon, J., van der Brug, M., Lopez de Munain, A., Aparicio, S., Gil, A. M., Khan, N., *et al.* (2004). Cloning of the gene containing mutations that cause PARK8-linked Parkinson's disease. *Neuron* 44, 595-600.

Palacino, J. J., Sagi, D., Goldberg, M. S., Krauss, S., Motz, C., Wacker, M., Klose, J., and Shen, J. (2004). Mitochondrial dysfunction and oxidative damage in parkin-deficient mice. *J Biol Chem* 279, 18614-18622.

Palmiter, R. D., and Findley, S. D. (1995). Cloning and functional characterization of a mammalian zinc transporter that confers resistance to zinc. *Embo J* 14, 639-649.

Palmiter, R. D., and Huang, L. (2004). Efflux and compartmentalization of zinc by members of the SLC30 family of solute carriers. *Pflugers Arch* 447, 744-751.

Park, J., Kim, S. Y., Cha, G. H., Lee, S. B., Kim, S., and Chung, J. (2005). Drosophila DJ-1 mutants show oxidative stress-sensitive locomotive dysfunction. *Gene* 361, 133-139.

Park, J., Lee, S. B., Lee, S., Kim, Y., Song, S., Kim, S., Bae, E., Kim, J., Shong, M., Kim, J. M., and Chung, J. (2006). Mitochondrial dysfunction in Drosophila PINK1 mutants is complemented by parkin. *Nature* 441, 1157-1161.

Parkinson, J. (2002). An essay on the shaking palsy. 1817. *J Neuropsychiatry Clin Neurosci* 14, 223-236; discussion 222.

Patton, E. E., Willems, A. R., and Tyers, M. (1998). Combinatorial control in ubiquitin-dependent proteolysis: don't Skp the F-box hypothesis. *Trends Genet* 14, 236-243.

Pauly, P. C., and Harris, D. A. (1998). Copper stimulates endocytosis of the prion protein. *J Biol Chem* 273, 33107-33110.

Pawlyk, A. C., Giasson, B. I., Sampathu, D. M., Perez, F. A., Lim, K. L., Dawson, V. L., Dawson, T. M., Palmiter, R. D., Trojanowski, J. Q., and Lee, V. M. (2003). Novel monoclonal antibodies demonstrate biochemical variation of brain parkin with age. *J Biol Chem* 278, 48120-48128.

Perez, F. A., and Palmiter, R. D. (2005). Parkin-deficient mice are not a robust model of parkinsonism. *Proc Natl Acad Sci U S A* 102, 2174-2179.

Perez, R. G., Waymire, J. C., Lin, E., Liu, J. J., Guo, F., and Zigmond, M. J. (2002). A role for alpha-synuclein in the regulation of dopamine biosynthesis. *J Neurosci* 22, 3090-3099.

Pesah, Y., Pham, T., Burgess, H., Middlebrooks, B., Verstreken, P., Zhou, Y., Harding, M., Bellen, H., and Mardon, G. (2004). Drosophila parkin mutants have

decreased mass and cell size and increased sensitivity to oxygen radical stress. *Development* 131, 2183-2194.

Peters, J. L., Dufner-Beattie, J., Xu, W., Geiser, J., Lahner, B., Salt, D. E., and Andrews, G. K. (2007). Targeting of the mouse *Slc39a2* (*Zip2*) gene reveals highly cell-specific patterns of expression, and unique functions in zinc, iron, and calcium homeostasis. *Genesis* 45, 339-352.

Petrucelli, L., O'Farrell, C., Lockhart, P. J., Baptista, M., Kehoe, K., Vink, L., Choi, P., Wolozin, B., Farrer, M., Hardy, J., and Cookson, M. R. (2002). Parkin protects against the toxicity associated with mutant alpha-synuclein: proteasome dysfunction selectively affects catecholaminergic neurons. *Neuron* 36, 1007-1019.

Pickart, C. M. (2001). Mechanisms underlying ubiquitination. *Annu Rev Biochem* 70, 503-533.

Pollanen, M. S., Dickson, D. W., and Bergeron, C. (1993). Pathology and biology of the Lewy body. *J Neuropathol Exp Neurol* 52, 183-191.

Polymeropoulos, M. H., Higgins, J. J., Golbe, L. I., Johnson, W. G., Ide, S. E., Di Iorio, G., Sanges, G., Stenroos, E. S., Pho, L. T., Schaffer, A. A., *et al.* (1996). Mapping of a gene for Parkinson's disease to chromosome 4q21-q23. *Science* 274, 1197-1199.

Polymeropoulos, M. H., Lavedan, C., Leroy, E., Ide, S. E., Dehejia, A., Dutra, A., Pike, B., Root, H., Rubenstein, J., Boyer, R., *et al.* (1997). Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science* 276, 2045-2047.

Radtke, F., Heuchel, R., Georgiev, O., Hergersberg, M., Gariglio, M., Dembic, Z., and Schaffner, W. (1993). Cloned transcription factor MTF-1 activates the mouse metallothionein I promoter. *Embo J* 12, 1355-1362.

Ren, Y., Zhao, J., and Feng, J. (2003). Parkin binds to alpha/beta tubulin and increases their ubiquitination and degradation. *J Neurosci* 23, 3316-3324.

Ribeiro, C. S., Carneiro, K., Ross, C. A., Menezes, J. R., and Engelender, S. (2002). Synphilin-1 is developmentally localized to synaptic terminals, and its association with synaptic vesicles is modulated by alpha-synuclein. *J Biol Chem* 277, 23927-23933.

Rohe, C. F., Montagna, P., Breedveld, G., Cortelli, P., Oostra, B. A., and Bonifati, V. (2004). Homozygous PINK1 C-terminus mutation causing early-onset parkinsonism. *Ann Neurol* 56, 427-431.

Ruiz, F. H., Silva, E., and Inestrosa, N. C. (2000). The N-terminal tandem repeat region of human prion protein reduces copper: role of tryptophan residues. *Biochem Biophys Res Commun* 269, 491-495.

Rutherford, J. C., and Bird, A. J. (2004). Metal-responsive transcription factors that regulate iron, zinc, and copper homeostasis in eukaryotic cells. *Eukaryot Cell* 3, 1-13.

Ryu, E. J., Harding, H. P., Angelastro, J. M., Vitolo, O. V., Ron, D., and Greene, L. A. (2002). Endoplasmic reticulum stress and the unfolded protein response in cellular models of Parkinson's disease. *J Neurosci* 22, 10690-10698.

Sadhu, C., and Gedamu, L. (1988). Regulation of human metallothionein (MT) genes. Differential expression of MTI-F, MTI-G, and MTII-A genes in the hepatoblastoma cell line (HepG2). *J Biol Chem* 263, 2679-2684.

Sakata, E., Yamaguchi, Y., Kurimoto, E., Kikuchi, J., Yokoyama, S., Yamada, S., Kawahara, H., Yokosawa, H., Hattori, N., Mizuno, Y., *et al.* (2003). Parkin binds the Rpn10 subunit of 26S proteasomes through its ubiquitin-like domain. *EMBO Rep* 4, 301-306.

Satoh, J., and Kuroda, Y. (2001). A polymorphic variation of serine to tyrosine at codon 18 in the ubiquitin C-terminal hydrolase-L1 gene is associated with a reduced risk of sporadic Parkinson's disease in a Japanese population. *J Neurol Sci* 189, 113-117.

Saydam, N., Adams, T. K., Steiner, F., Schaffner, W., and Freedman, J. H. (2002). Regulation of metallothionein transcription by the metal-responsive transcription factor MTF-1: identification of signal transduction cascades that control metal-inducible transcription. *J Biol Chem* 277, 20438-20445.

Saydam, N., Georgiev, O., Nakano, M. Y., Greber, U. F., and Schaffner, W. (2001). Nucleo-cytoplasmic trafficking of metal-regulatory transcription factor 1 is regulated by diverse stress signals. *J Biol Chem* 276, 25487-25495.

Schapira, A. H. (2008). Mitochondria in the aetiology and pathogenesis of Parkinson's disease. *Lancet Neurol* 7, 97-109.

Schlossmacher, M. G., Frosch, M. P., Gai, W. P., Medina, M., Sharma, N., Forno, L., Ochiishi, T., Shimura, H., Sharon, R., Hattori, N., *et al.* (2002). Parkin localizes to the Lewy bodies of Parkinson disease and dementia with Lewy bodies. *Am J Pathol* 160, 1655-1667.

Schmidt, M. L., Murray, J., Lee, V. M., Hill, W. D., Wertkin, A., and Trojanowski, J. Q. (1991). Epitope map of neurofilament protein domains in cortical and peripheral nervous system Lewy bodies. *Am J Pathol* 139, 53-65.

Selvaraj, A., Balamurugan, K., Yepiskoposyan, H., Zhou, H., Egli, D., Georgiev, O., Thiele, D. J., and Schaffner, W. (2005). Metal-responsive transcription factor (MTF-1) handles both extremes, copper load and copper starvation, by activating different genes. *Genes Dev* 19, 891-896.

Shashidharan, P., Good, P. F., Hsu, A., Perl, D. P., Brin, M. F., and Olanow, C. W. (2000). TorsinA accumulation in Lewy bodies in sporadic Parkinson's disease. *Brain Res* 877, 379-381.

Shen, J. (2004). Protein kinases linked to the pathogenesis of Parkinson's disease. *Neuron* 44, 575-577.

Shendelman, S., Jonason, A., Martinat, C., Leete, T., and Abeliovich, A. (2004). DJ-1 is a redox-dependent molecular chaperone that inhibits alpha-synuclein aggregate formation. *PLoS Biol* 2, e362.

Shi, Y., Vattam, K. M., Sood, R., An, J., Liang, J., Stramm, L., and Wek, R. C. (1998). Identification and characterization of pancreatic eukaryotic initiation factor 2 alpha-subunit kinase, PEK, involved in translational control. *Mol Cell Biol* 18, 7499-7509.

Shimura, H., Hattori, N., Kubo, S., Mizuno, Y., Asakawa, S., Minoshima, S., Shimizu, N., Iwai, K., Chiba, T., Tanaka, K., and Suzuki, T. (2000). Familial Parkinson disease gene product, parkin, is a ubiquitin-protein ligase. *Nat Genet* 25, 302-305.

Shimura, H., Hattori, N., Kubo, S., Yoshikawa, M., Kitada, T., Matsumine, H., Asakawa, S., Minoshima, S., Yamamura, Y., Shimizu, N., and Mizuno, Y. (1999). Immunohistochemical and subcellular localization of Parkin protein: absence of protein in autosomal recessive juvenile parkinsonism patients. *Ann Neurol* 45, 668-672.

Shimura, H., Schlossmacher, M. G., Hattori, N., Frosch, M. P., Trockenbacher, A., Schneider, R., Mizuno, Y., Kosik, K. S., and Selkoe, D. J. (2001). Ubiquitination of a new form of alpha-synuclein by parkin from human brain: implications for Parkinson's disease. *Science* 293, 263-269.

Singleton, A. B., Farrer, M., Johnson, J., Singleton, A., Hague, S., Kachergus, J., Hulihan, M., Peuralinna, T., Dutra, A., Nussbaum, R., *et al.* (2003). alpha-Synuclein locus triplication causes Parkinson's disease. *Science* 302, 841.

Smith, W. W., Margolis, R. L., Li, X., Troncoso, J. C., Lee, M. K., Dawson, V. L., Dawson, T. M., Iwatsubo, T., and Ross, C. A. (2005). Alpha-synuclein phosphorylation enhances eosinophilic cytoplasmic inclusion formation in SH-SY5Y cells. *J Neurosci* 25, 5544-5552.

Sorkina, T., Doolen, S., Galperin, E., Zahniser, N. R., and Sorkin, A. (2003). Oligomerization of dopamine transporters visualized in living cells by fluorescence resonance energy transfer microscopy. *J Biol Chem* 278, 28274-28283.

Spacey, S. D., and Wood, N. W. (1999). The genetics of Parkinson's disease. *Curr Opin Neurol* 12, 427-432.

Spillantini, M. G., and Goedert, M. (2000). The alpha-synucleinopathies: Parkinson's disease, dementia with Lewy bodies, and multiple system atrophy. *Ann N Y Acad Sci* 920, 16-27.

Spillantini, M. G., Schmidt, M. L., Lee, V. M., Trojanowski, J. Q., Jakes, R., and Goedert, M. (1997). Alpha-synuclein in Lewy bodies. *Nature* 388, 839-840.

Staropoli, J. F., McDermott, C., Martinat, C., Schulman, B., Demireva, E., and Abeliovich, A. (2003). Parkin is a component of an SCF-like ubiquitin ligase complex and protects postmitotic neurons from kainate excitotoxicity. *Neuron* 37, 735-749.

Stathakis, D. G., Burton, D. Y., McIvor, W. E., Krishnakumar, S., Wright, T. R., and O'Donnell, J. M. (1999). The catecholamines up (Catsup) protein of *Drosophila melanogaster* functions as a negative regulator of tyrosine hydroxylase activity. *Genetics* 153, 361-382.

Stefanis, L., Larsen, K. E., Rideout, H. J., Sulzer, D., and Greene, L. A. (2001). Expression of A53T mutant but not wild-type alpha-synuclein in PC12 cells induces alterations of the ubiquitin-dependent degradation system, loss of dopamine release, and autophagic cell death. *J Neurosci* 21, 9549-9560.

Stennard, F. A., Holloway, A. F., Hamilton, J., and West, A. K. (1994). Characterisation of six additional human metallothionein genes. *Biochim Biophys Acta* 1218, 357-365.

Stichel, C. C., Augustin, M., Kuhn, K., Zhu, X. R., Engels, P., Ullmer, C., and Lubbert, H. (2000). Parkin expression in the adult mouse brain. *Eur J Neurosci* 12, 4181-4194.

Stuart, G. W., Searle, P. F., Chen, H. Y., Brinster, R. L., and Palmiter, R. D. (1984). A 12-base-pair DNA motif that is repeated several times in metallothionein gene promoters confers metal regulation to a heterologous gene. *Proc Natl Acad Sci U S A* 81, 7318-7322.

Stuart, G. W., Searle, P. F., and Palmiter, R. D. (1985). Identification of multiple metal regulatory elements in mouse metallothionein-I promoter by assaying synthetic sequences. *Nature* 317, 828-831.

Sulzer, D., and Zecca, L. (2000). Intraneuronal dopamine-quinone synthesis: a review. *Neurotox Res* 1, 181-195.

Szegezdi, E., Fitzgerald, U., and Samali, A. (2003). Caspase-12 and ER-stress-mediated apoptosis: the story so far. *Ann N Y Acad Sci* 1010, 186-194.

Taira, T., Saito, Y., Niki, T., Iguchi-Ariga, S. M., Takahashi, K., and Ariga, H. (2004). DJ-1 has a role in antioxidative stress to prevent cell death. *EMBO Rep* 5, 213-218.

Taira, T., Takahashi, K., Kitagawa, R., Iguchi-Arigo, S. M., and Ariga, H. (2001). Molecular cloning of human and mouse DJ-1 genes and identification of Sp1-dependent activation of the human DJ-1 promoter. *Gene* 263, 285-292.

Takahashi, H., Ohama, E., Suzuki, S., Horikawa, Y., Ishikawa, A., Morita, T., Tsuji, S., and Ikuta, F. (1994). Familial juvenile parkinsonism: clinical and pathologic study in a family. *Neurology* 44, 437-441.

Takahashi, R., Imai, Y., Hattori, N., and Mizuno, Y. (2003). Parkin and endoplasmic reticulum stress. *Ann N Y Acad Sci* 991, 101-106.

Tanaka, Y., Engelender, S., Igarashi, S., Rao, R. K., Wanner, T., Tanzi, R. E., Sawa, A., V. L. D., Dawson, T. M., and Ross, C. A. (2001). Inducible expression of mutant alpha-synuclein decreases proteasome activity and increases sensitivity to mitochondria-dependent apoptosis. *Hum Mol Genet* 10, 919-926.

Tompkins, M. M., and Hill, W. D. (1997). Contribution of somal Lewy bodies to neuronal death. *Brain Res* 775, 24-29.

Torres, G. E., Gainetdinov, R. R., and Caron, M. G. (2003). Plasma membrane monoamine transporters: structure, regulation and function. *Nat Rev Neurosci* 4, 13-25.

Tritschler, H. J., Packer, L., and Medori, R. (1994). Oxidative stress and mitochondrial dysfunction in neurodegeneration. *Biochem Mol Biol Int* 34, 169-181.

Twelves, D., Perkins, K. S., and Counsell, C. (2003). Systematic review of incidence studies of Parkinson's disease. *Mov Disord* 18, 19-31.

Unoki, M., and Nakamura, Y. (2001). Growth-suppressive effects of BPOZ and EGR2, two genes involved in the PTEN signaling pathway. *Oncogene* 20, 4457-4465.

Valente, E. M., Abou-Sleiman, P. M., Caputo, V., Muqit, M. M., Harvey, K., Gispert, S., Ali, Z., Del Turco, D., Bentivoglio, A. R., Healy, D. G., *et al.* (2004). Hereditary early-onset Parkinson's disease caused by mutations in PINK1. *Science* 304, 1158-1160.

Volles, M. J., and Lansbury, P. T., Jr. (2002). Vesicle permeabilization by protofibrillar alpha-synuclein is sensitive to Parkinson's disease-linked mutations and occurs by a pore-like mechanism. *Biochemistry* 41, 4595-4602.

Volles, M. J., Lee, S. J., Rochet, J. C., Shtilerman, M. D., Ding, T. T., Kessler, J. C., and Lansbury, P. T., Jr. (2001). Vesicle permeabilization by protofibrillar alpha-synuclein: implications for the pathogenesis and treatment of Parkinson's disease. *Biochemistry* 40, 7812-7819.

von Coelln, R., Dawson, V. L., and Dawson, T. M. (2004a). Parkin-associated Parkinson's disease. *Cell Tissue Res* 318, 175-184.

Von Coelln, R., Thomas, B., Savitt, J. M., Lim, K. L., Sasaki, M., Hess, E. J., Dawson, V. L., and Dawson, T. M. (2004b). Loss of locus coeruleus neurons and reduced startle in parkin null mice. *Proc Natl Acad Sci U S A* 101, 10744-10749.

Wadsworth, J. D., Jackson, G. S., Hill, A. F., and Collinge, J. (1999). Molecular biology of prion propagation. *Curr Opin Genet Dev* 9, 338-345.

Wakabayashi, K., Engelender, S., Yoshimoto, M., Tsuji, S., Ross, C. A., and Takahashi, H. (2000). Synphilin-1 is present in Lewy bodies in Parkinson's disease. *Ann Neurol* 47, 521-523.

Wang, K., Zhou, B., Kuo, Y. M., Zemansky, J., and Gitschier, J. (2002). A novel member of a zinc transporter family is defective in acrodermatitis enteropathica. *Am J Hum Genet* 71, 66-73.

West, A. B., Lockhart, P. J., O'Farrell, C., and Farrer, M. J. (2003). Identification of a novel gene linked to parkin via a bi-directional promoter. *J Mol Biol* 326, 11-19.

West, A. K., Hidalgo, J., Eddins, D., Levin, E. D., and Aschner, M. (2008). Metallothionein in the central nervous system: Roles in protection, regeneration and cognition. *Neurotoxicology* 29, 489-503.

West, A. K., Stallings, R., Hildebrand, C. E., Chiu, R., Karin, M., and Richards, R. I. (1990). Human metallothionein genes: structure of the functional locus at 16q13. *Genomics* 8, 513-518.

Westin, G., and Schaffner, W. (1988). A zinc-responsive factor interacts with a metal-regulated enhancer element (MRE) of the mouse metallothionein-I gene. *Embo J* 7, 3763-3770.

Whitworth, A. J., Theodore, D. A., Greene, J. C., Benes, H., Wes, P. D., and Pallanck, L. J. (2005). Increased glutathione S-transferase activity rescues dopaminergic neuron loss in a *Drosophila* model of Parkinson's disease. *Proc Natl Acad Sci U S A* 102, 8024-8029.

Wilkinson, K. D., Lee, K. M., Deshpande, S., Duerksen-Hughes, P., Boss, J. M., and Pohl, J. (1989). The neuron-specific protein PGP 9.5 is a ubiquitin carboxyl-terminal hydrolase. *Science* 246, 670-673.

Wimmer, U., Wang, Y., Georgiev, O., and Schaffner, W. (2005). Two major branches of anti-cadmium defense in the mouse: MTF-1/metallothioneins and glutathione. *Nucleic Acids Res* 33, 5715-5727.

Withers, G. S., George, J. M., Banker, G. A., and Clayton, D. F. (1997). Delayed localization of synelfin (synuclein, NACP) to presynaptic terminals in cultured rat hippocampal neurons. *Brain Res Dev Brain Res* 99, 87-94.

Wszolek, Z. K., Pfeiffer, R. F., Tsuboi, Y., Uitti, R. J., McComb, R. D., Stoessl, A. J., Strongosky, A. J., Zimprich, A., Muller-Myhsok, B., Farrer, M. J., *et al.* (2004). Autosomal dominant parkinsonism associated with variable synuclein and tau pathology. *Neurology* 62, 1619-1622.

Yang, F., Jiang, Q., Zhao, J., Ren, Y., Sutton, M. D., and Feng, J. (2005). Parkin stabilizes microtubules through strong binding mediated by three independent domains. *J Biol Chem* 280, 17154-17162.

Yang, Y., Gehrke, S., Imai, Y., Huang, Z., Ouyang, Y., Wang, J. W., Yang, L., Beal, M. F., Vogel, H., and Lu, B. (2006). Mitochondrial pathology and muscle and dopaminergic neuron degeneration caused by inactivation of *Drosophila* Pink1 is rescued by Parkin. *Proc Natl Acad Sci U S A* 103, 10793-10798.

Yang, Y., Nishimura, I., Imai, Y., Takahashi, R., and Lu, B. (2003). Parkin suppresses dopaminergic neuron-selective neurotoxicity induced by Pael-R in *Drosophila*. *Neuron* 37, 911-924.

Yavich, L., Tanila, H., Vepsäläinen, S., and Jakala, P. (2004). Role of alpha-synuclein in presynaptic dopamine recruitment. *J Neurosci* 24, 11165-11170.

Yepiskoposyan, H., Egli, D., Fergestad, T., Selvaraj, A., Treiber, C., Multhaup, G., Georgiev, O., and Schaffner, W. (2006). Transcriptome response to heavy metal stress in *Drosophila* reveals a new zinc transporter that confers resistance to zinc. *Nucleic Acids Res* 34, 4866-4877.

Zarranz, J. J., Alegre, J., Gomez-Esteban, J. C., Lezcano, E., Ros, R., Ampuero, I., Vidal, L., Hoenicka, J., Rodriguez, O., Atares, B., *et al.* (2004). The new mutation, E46K, of alpha-synuclein causes Parkinson and Lewy body dementia. *Ann Neurol* 55, 164-173.

Zhang, B., Egli, D., Georgiev, O., and Schaffner, W. (2001). The *Drosophila* homolog of mammalian zinc finger factor MTF-1 activates transcription in response to heavy metals. *Mol Cell Biol* 21, 4505-4514.

Zhang, B., Georgiev, O., Hagmann, M., Gunes, C., Cramer, M., Faller, P., Vasak, M., and Schaffner, W. (2003). Activity of metal-responsive transcription factor 1 by toxic heavy metals and H₂O₂ in vitro is modulated by metallothionein. *Mol Cell Biol* 23, 8471-8485.

Zhang, J., Perry, G., Smith, M. A., Robertson, D., Olson, S. J., Graham, D. G., and Montine, T. J. (1999). Parkinson's disease is associated with oxidative damage to cytoplasmic DNA and RNA in substantia nigra neurons. *Am J Pathol* 154, 1423-1429.

Zhang, Y., Dawson, V. L., and Dawson, T. M. (2000). Oxidative stress and genetics in the pathogenesis of Parkinson's disease. *Neurobiol Dis* 7, 240-250.

Zhao, H., Butler, E., Rodgers, J., Spizzo, T., Duesterhoeft, S., and Eide, D. (1998). Regulation of zinc homeostasis in yeast by binding of the ZAP1 transcriptional activator to zinc-responsive promoter elements. *J Biol Chem* 273, 28713-28720.

Zimprich, A., Biskup, S., Leitner, P., Lichtner, P., Farrer, M., Lincoln, S., Kachergus, J., Hulihan, M., Uitti, R. J., Calne, D. B., *et al.* (2004). Mutations in LRRK2 cause autosomal-dominant parkinsonism with pleomorphic pathology. *Neuron* 44, 601-607.

Neurobiol Dis. 2010 Oct;40(1):82-92. Epub 2010 May 17.

Extended lifespan of *Drosophila parkin* mutants through sequestration of redox-active metals and enhancement of anti-oxidative pathways.

Saini N, Oelhafen S, Hua H, Georgiev O, Schaffner W, Büeler H.

Institute for Molecular Life Sciences, University of Zürich,
Winterthurerstrasse 190, CH-8051 Zürich, Switzerland.

Abstract

The mechanisms underlying neuron death in Parkinson's disease are unknown, but both genetic defects and environmental factors are implicated in its pathogenesis. Mutations in the *parkin* gene lead to autosomal recessive juvenile Parkinsonism (AR-JP). Here we report that compared to control flies, *Drosophila* lacking *parkin* show significantly reduced lifespan but no difference in dopamine neuron numbers when raised on food supplemented with environmental pesticides or mitochondrial toxins. Moreover, chelation of redox-active metals, antioxidants and overexpression of superoxide dismutase 1 all significantly reversed the reduced longevity of *parkin*-deficient flies. Finally, *parkin* deficiency exacerbated the rough eye phenotype of *Drosophila* caused by overexpression of the copper importer B (Ctr1B). Taken together, our results demonstrate an important function of *parkin* in the protection against redox-active metals and pesticides implicated in the etiology of Parkinson's disease. They also corroborate that oxidative stress, perhaps as a consequence of mitochondrial dysfunction, is a major determinant of morbidity in *parkin* mutant flies.

Introduction

A combination of mitochondrial dysfunction, oxidative stress and abnormal protein aggregation underlies the pathogenesis of PD ([Bueler, 2009], [Henchcliffe and Beal, 2008], [Cookson, 2005], [Dawson and Dawson, 2003], [Moore et al., 2005] and [Shen and Cookson, 2004]). About 10% of PD cases are heritable and caused by mutations in at least five genes ([Gasser, 2009], [Hardy et al., 2009] and [Hatano et al., 2009]). Parkin (PARK2) mutations lead to autosomal recessive juvenile Parkinsonism (AR-JP) ([Abbas et al., 1999] and [Kitada et al., 1998]). Parkin functions as an E3 ubiquitin ligase in the proteasome-mediated turnover of several proteins in vitro ([Corti et al., 2003], [Imai et al., 2001], [Shimura et al., 2000] and [Zhang et al., 2000]), some of which accumulate in the brains of AR-JP patients and/or mice carrying a targeted deletion of the Parkin gene ([Fukae et al., 2009], [Murakami et al., 2004] and [Periquet et al., 2005]). Ablation of Parkin in mice resulted in mitochondrial respiratory defects, decreased levels of antioxidant proteins (Palacino et al., 2004) and abnormalities in dopamine neurotransmission, but not loss of dopamine neurons ([Goldberg et al., 2003] and [Itier et al., 2003]). Recently, Parkin has been linked to DNA repair ([Kao, 2009a], [Kao, 2009b] and [Rothfuss et al., 2009]). Parkin was shown to enhance the replication, transcription and repair of mitochondrial DNA, thereby protecting mitochondrial DNA against oxidative damage (Rothfuss et al., 2009). *Drosophila* lacking Parkin display increased sensitivity to oxidative stress and severe structural mitochondrial abnormalities in muscle and germline tissues associated with apoptotic muscle degeneration and male sterility ([Greene et al., 2005], [Greene et al., 2003] and [Pesah et al., 2004]). There is controversy as to whether or not frank dopamine neuron death occurs in flies with *parkin* loss-of-function mutations. Limited dopamine neuron death restricted to the PPL1/DL1 cluster has been detected in 20-day-old *parkin* mutant flies (park25 allele) by tyrosine hydroxylase (TH) immunohistochemistry combined with whole-mount confocal microscopy (Whitworth et al., 2005). However, the original characterization of the same flies showed that dopamine neuron numbers determined in paraffin-embedded sections were normal, even in older *Drosophila* (30 days) (Greene et al.,

2003). Another line of *parkin* mutant flies (Pesah et al., 2004) also lacked dopamine neuron death, but the PPL1 cluster showing neuron loss (Whitworth et al., 2005) was not analyzed in this study. Although (Greene et al., 2003) and (Whitworth et al., 2005) used different methods to quantify TH neurons, it cannot be concluded that confocal microscopy is more likely to detect TH neuron loss than using paraffin-embedded sections. In fact, in α -synuclein transgenic flies, TH neuron loss was detected using paraffin-embedded sections (Feany and Bender, 2000), but not with whole-mount confocal microscopy (Pesah et al., 2005), although both groups used the same flies again. Finally, several groups failed to detect dopamine neuron loss in Parkin-deficient mice ([Goldberg et al., 2003], [Itier et al., 2003] and [Palacino et al., 2004]). Taken together, the data suggest that Parkin ablation may lead to neuron death in one of several dopamine neuron clusters in the *Drosophila* brain, while Parkin appears to be dispensible for dopamine neuron survival in mice.

Several epidemiological studies showed that long-term exposure to pesticides, such as rotenone and paraquat, correlates with an increased risk of developing idiopathic PD ([Abbott et al., 2003], [Baldi et al., 2003], [Gorell et al., 1998] and [Petrovitch et al., 2002]). Intravenous administration of an illicit drug, which contained 1-methyl 4-phenyl 1,2,3,6-tetrahydropyridine (MPTP), caused an acute and permanent state of Parkinsonism in humans (Langston et al., 1983). Moreover, excessive accumulation of iron in the substantia nigra has been implicated in PD pathogenesis ([Dexter et al., 1992], [Friedman et al., 2007] and [Zecca et al., 2004]). Free iron is toxic to cells as it acts as a catalyst in the formation of free radicals from hydrogen peroxide (H₂O₂) via the Fenton reaction (Orino et al., 2001). Because H₂O₂ is a normal product of the dopamine metabolism, dopaminergic neurons may be particularly vulnerable to accumulation of free iron. Support for increased free iron levels in PD comes from studies that showed reduced amounts of l-ferritin, which is important for iron sequestration (Harrison and Arosio, 1996), in PD compared to control brains ([Connor et al., 1995] and [Koziorowski et al., 2007]). In addition to iron, copper toxicity may play a role in various neurological disorders (Desai and Kaler, 2008). In Wilson's disease, increased copper

accumulation leads to liver dysfunction and neurological symptoms (Ala et al., 2007). Interestingly, patients with Wilson's disease often present with parkinsonism (Oder et al., 1993 W. Oder et al., Wilson's disease: evidence of subgroups derived from clinical findings and brain lesions, *Neurology* 43 (1993), pp. 120–124. View Record in Scopus | Cited By in Scopus (59)Oder et al., 1993) and deficits in dopamine neurotransmission have been reported (Barthel et al., 2003). Furthermore, neurological symptoms were shown to correlate with accumulation of copper in brain areas affected in PD, such as the basal ganglia and substantia nigra (Sudmeyer et al., 2006). This suggests that the nigrostriatal system may be particularly sensitive to copper accumulation. In agreement with this notion, long-term occupational exposure to copper was associated with an increased risk for PD (Gorell et al., 1997), and treatment of cultured dopaminergic neurons with copper led to mitochondrial defects, decreased dopamine content and cell death (Shi et al., 2008).

Because no single, unified mechanism leads to PD, it has been postulated that exogenous and/or endogenous stress factors may synergize with genetic defects to cause the precipitation of PD. This “gene–environmental factor interaction hypothesis” states that mutations in familial PD genes may increase the vulnerability to PD-associated environmental factors and toxins. Here, we hypothesized that *parkin* represents a genetic susceptibility factor in the vulnerability of flies to redox-active metals, pesticides and mitochondrial toxins implicated in PD. To test this hypothesis, flies carrying a null mutation (*park25/25*) of the *Drosophila* homologue of *parkin* (Greene et al., 2003) were raised on food supplemented with redox-active metals, metal chelators, rotenone, paraquat and MPP⁺, and their survival was compared to *park25/+* control flies. The interaction of copper with *parkin* deficiency was further studied in flies overexpressing the copper importer Ctr1B. Finally we increased the anti-oxidant defense in flies by various means to corroborate the causative role of oxidative stress in the reduced longevity observed in *parkin*-deficient flies. Our results demonstrate that loss of Parkin can greatly increase the flies' vulnerability to environmental factors, mitochondrial toxins and certain redox-active metals implicated in the etiology of PD, and reinforce that

oxidative stress is a major pathway in causing the morbidity of *parkin*-deficient flies.

Materials and methods

Fly stocks and maintenance

All flies were maintained at 25 °C on a 12:12 h light–dark cycle. Unless specified otherwise, heterozygous *parkin* mutant flies (*w*;+;*park25/TM6B*, *w*+, hereafter referred to as *park25/+*) were used as controls for the homozygous *parkin* mutant flies (*w*;+;*park25/park25*, hereafter referred to as *park25/25*). *park25/25* is null mutation of the *Drosophila parkin* gene (Greene et al., 2003). The TM6B balancer in the control flies was confirmed to have no effect on any of the experiments performed (by having it crossed out). The *w*; *UAS-SOD1/Cyo*;+ line expressing SOD1 under the mesoderm-specific twist-Gal4 driver (Greig and Akam, 1993) was received from the Bloomington stock center. These flies were crossed to the *w*;+;*park25/TM6B*, *w*+ and *twi-Gal4*;+;+ flies to obtain *twi-Gal4*; *UAS-SOD1*; *park25/+* animals, which were bred to obtain *twi-Gal4*; *UAS-SOD1*; *park25/25* flies. The Ctr1B coding region was amplified by PCR (with primers 5'-TGGACTAGATCTCAACATGGATCACGGCTCGGAT-3' and 5'-GTCAGT-AGATCTCTAAGGACAGCACTCGCT-3') from a Ctr1B cDNA clone and inserted into the SmaI site of pBluescript (Stratagene). The resulting construct was digested with EcoRI and NotI and the fragment was cloned between the EcoRI and NotI sites of the P element vector pUAST. The GMR-Ctr1B transgenic flies were generated with a Ctr1B ORF driven by the eye-specific GMR promoter/enhancer. The *w*; *GMR-Ctr1B/Cyo*; *park25/+* transgenic flies were obtained by crossing the *w*; *GMR-Ctr1B/Cyo* flies with *w*;+;*park25/TM6B*, *w*+ flies.

Preparation of fly food for oxidative stress, metal toxicity and metal chelator experiments

One liter standard fly food (normal food, NF) contained 55 g cornmeal, 10 g wheat flour, 100 g yeast, 75 g glucose, 8 g agar and 15 ml anti-fungal agent

nipagin (15% in ethanol). Normal food was supplemented with 200 μ M rotenone, 10 mM paraquat and 25 mM 1-methyl-4-phenylpyridinium iodide (MPP+) (final concentrations, all from Sigma). For the metal toxicity/chelation and antioxidant experiments, normal food was supplemented with 500 μ M ammonium ferric citrate, 500 μ M copper (II) sulfate, 0.3 mM bathocuproine disulfonate (copper chelator), 0.1 mM bathophenanthroline sulfonated sodium salt (iron chelator), 20 or 50 μ M silver nitrate, 15 mM N-acetyl-cysteine and 100, 250 or 500 mM ascorbic acid.

Lifespan determination

Lifespan (survival) experiments were carried out in triplicate. In each experiment, twenty 1- to 2-day-old flies per genotype were analyzed. Surviving flies were counted daily and transferred to fresh food vials every second day, with the exception of the SOD1 overexpression experiment where flies were counted only on the four indicated days. As a control, flies of the same genotype were grown on food lacking the toxins, metals, metal-chelating agents or antioxidants. The variations in the average lifespan of control flies in different experiments can be attributed to subtle experimental variations.

Exposure to hypoxia and heat shock

park25/+ and *park25/25* flies (six sets of 20 flies each) were exposed for 7 min to 0.5% oxygen at 25 °C in a closed hypoxia chamber, and the time to regain consciousness (ability to move) was recorded. To achieve a stable atmosphere of 0.5% oxygen, ambient air was mixed with nitrogen gas (Digamix M302 a-F, H Wösthoff e.H.G) and pumped into the chamber (VENT2 pump, EMKA Technologies) for 4 min at a flow rate of 25 l per minute. In the heat shock experiment, flies were incubated at 38 °C for 30 or 60 min and surviving animals were counted 24 h later.

Quantification of food uptake (gustatory assay)

For the adult gustatory assay, we performed a protocol similar to the one described by Hilliker and colleagues (Bahadorani et al., 2008). Briefly, newly eclosed flies were reared on normal food for 2–3 days before being starved for 18 h on Whatman paper soaked with distilled water. Subsequently, starved flies ($n = 20$) were transferred into a vial with fresh normal food (control) or chelator-supplemented food, both containing 0.2 % sulforhodamine B sodium salt (acid-red), and allowed to feed for 2 h. After 2 h of feeding, flies were decapitated and their bodies (without heads) were homogenized individually in 100 μ l ice-cold buffer (0.32 M sucrose, 20 mM HEPES, 1 mM $MgCl_2$, 0.5 mM PMSF, pH 7.4). 50 μ l of the homogenate supernatants were transferred to separate wells of a microtiter (ELISA) plate and absorbance was measured at 565 nm (absorbance maximum for acid-red) using a multi-label counter (Wallac 1429, Perkin Elmer).

Quantification of brain ROS levels

Twenty heads dissected from adult flies were homogenized in 3 volumes of ice-cold buffer (0.32 mM sucrose, 20 mM HEPES, 1 mM $MgCl_2$, 0.5 mM PMSF, pH 7.4) and centrifuged for 20 min at 4 °C at 20,000g. Total protein concentration in the supernatant was measured with the Bradford method (Biorad kit) and adjusted to 0.4 μ g/ μ l. 20 μ l sample (8 μ g protein) was mixed with 2 μ l 40 μ M DCF-DA in a fluorimeter cuvette (Hellma, 1.5 \times 1.5 mm). Excitation and emission maxima of DCF are at 485 nm and 520 nm, respectively. The shutter size of the fluorimeter was fixed at 2.5 nm and sample fluorescence intensity was monitored for 45 min at 26 °C between 500 and 600 nm. Fluorescence intensity was recorded every 5 min. The maximum peak values were then read as the final value, by integrating the area under the peak of the graphs.

Analysis of eye phenotype

Flies were frozen at –20 °C and pictures of their eyes were taken with a scanning electron microscope (model JEOL JSM-630LV) in the imaging

facility of the University of Zurich. Ten eyes per genotype and treatment condition were analyzed.

Detection of TH-positive neurons in brain sections by DAB immunohistochemistry

Flies were lined up on a Heisenberg collar and fixed in Carnoy's solution (60% dried ethanol, 30% chloroform, 10% acetic acid) overnight at 4 °C (Heisenberg and Böhl, 1979). They were then dehydrated for 40 min in absolute ethanol, twice for 40 min in dried ethanol and twice for 60 min in n-butanol. Paraffin-embedded flies were decapitated, cut into 5-micron-thick sections and transferred to Super Frost Plus slides. Slides were placed in xylol for 2–5 min for de-waxing and then transferred to absolute ethanol for 2–5 min, before incubation in 95% ethanol for 2–3 min. Then they were placed into 70% ethanol for 2–3 min and the ethanol was washed out. Sections were fixed in 0.5% formaldehyde/phosphate-buffered saline (PBS) for 30 min, rinsed three times in PBS and blocked by incubation for 30 min in 1% bovine serum albumin/0.05% Triton X-100. After washing four times for 5 min in PBS/0.05% Triton X-100 they were incubated in PBS/1% bovine serum albumin (4 °C overnight) with 1:200 diluted monoclonal anti-tyrosine hydroxylase antibody specific for dopamine neurons (Chemicon, MAB 318). After five washes with PBS/0.05% Triton X-100, 1:1000 diluted secondary biotin-conjugated goat anti-mouse IgG (Jackson ImmunoResearch) was added and the sections were incubated for 2 h at room temperature. Finally, they were washed five times with PBS and TH neurons were stained with 3,3'-diaminobenzidine (DAB) using the Vectastain Elite ABC Kit (Vector Laboratories).

Quantification of TH neurons in whole mount *Drosophila* brains by confocal microscopy

Fly brains were isolated and fixed for 2 h on ice in 100 mM Pipes/2 mM magnesium sulphate/1 mM EDTA/4% formaldehyde. They were rinsed twice with PBST (PBS/0.1% Tween-20), post-fixed with ice-cold methanol for 3

min, rinsed at least four times with PBST and blocked in PBST/10% fetal bovine serum (FBS) for 1 h at room temperature. They were then incubated in PBST/2% FBS with 1:200 diluted mouse anti-tyrosine hydroxylase antibody (Chemicon, MAB318) for at least 36 h at 4 °C, washed three times for 1 h in PBST at 4 °C, and incubated at 4 °C overnight with 1:500 diluted goat anti-mouse IgG conjugated to Alexa 488 (Invitrogen) in PBST/2% FBS. After three 1-h washes in PBST, the brains were mounted in 80% glycerol/0.1 M Tris-HCl pH 7.5. TH-positive neurons of the DM and DL1 clusters were counted in about thirty 2 micron-thick confocal brain sections per fly.

Statistical analysis

JMP software (SAS Institute) was used for statistical evaluations. Lifespan (survival) assays were analyzed with the Kaplan–Meier log-rank statistical test. Brain ROS levels were compared by one-way ANOVA. For all other experiments, data were compared using the Student's t-test. Results are expressed as mean \pm standard deviation.

Results

Increased vulnerability of *parkin*-deficient *Drosophila* to environmental pesticides and mitochondrial toxins associated with Parkinson's disease

For our studies we used the *park25/25* flies in which 75% of the Parkin coding sequence, including the start codon, had been removed by imprecise excision of a P-element insertion at the 5' end of the fly *parkin* gene (Greene et al., 2003). The deletion is thus a *parkin* null allele. As controls, we used *park25/+* (heterozygous mutant *parkin*) flies. To investigate whether *parkin* deficiency increases the susceptibility to environmental substances that are known to cause oxidative stress and have been associated with PD ([Abbott et al., 2003], [Dauer and Przedborski, 2003], [Langston et al., 1983] and [Petrovitch et al., 2002]), we compared the survival of *park25/25* and *park25/+* flies exposed to rotenone, MPP⁺ and paraquat. Flies were raised on food supplemented with 200 μ M rotenone, 25 mM MPP⁺ or 10 mM paraquat and their survival was monitored daily (Fig. 1). The toxin concentrations we used were in the range

reported for the same pesticides in *Drosophila* in previous publications, namely 2–40 mM for paraquat ([Jones et al., 2009], [Lee et al., 2009], [Pesah et al., 2004], [Schriner et al., 2009] and [Wang et al., 2008]) and 25–500 μ M for rotenone ([Coulom and Birman, 2004], [Hosamani and Muralidhara, 2009], [Wang et al., 2007] and [Wang et al., 2008]). For MPP+, we found no reference for *Drosophila* and choose a concentration that is at least 25 times higher than required to induce significant cell death in mammalian neuroblastoma cells (Bando et al., 2005). *park25/25* flies were more sensitive to and showed significantly shortened lifespan after treatment with all three toxins, when compared to either *park25/+* ($p < 0.001$) or *yw* ($p < 0.001$) controls (Fig. 1A–C). In contrast, *park25/+* flies were not more vulnerable to the toxins than *yw* control flies. All toxins further reduced the lifespan of *park25/25 Drosophila* ($p < 0.01$), showing that the experiment was able to detect an effect of the toxins on the *parkin*-deficient background. Unexpectedly, *park25/+* flies displayed increased lifespan compared to *yw* control flies on MPP+, an observation that we have not investigated further and for which we have no explanation at present. We believe it is unlikely that *park25/+* flies developed mitochondrial resistance to MPP+, because both MPP+ and rotenone are potent inhibitors of the mitochondrial complex I, but *park25/+* flies are not resistant to rotenone (Fig. 1A).

Lack of dopamine neuron death in Parkin-deficient flies exposed to environmental toxins

Although dopamine neurons of mice, and most of the dopamine neuron clusters in flies, survive in the absence of Parkin ([Goldberg et al., 2003], [Greene et al., 2003], [Itier et al., 2003], [Pesah et al., 2004] and [Whitworth et al., 2005]), mitochondrial abnormalities and the upregulation of components of the oxidative stress response in *parkin*-deficient *Drosophila* suggested that Parkin generally functions to protect mitochondria and cells against oxidative stress ([Greene et al., 2005], [Greene et al., 2003] and Palacino et al., 2004 J.J. Palacino et al., Mitochondrial dysfunction and oxidative damage in *parkin*-deficient mice (Palacino et al., 2004)). Given the reduced lifespan of *park25/25* flies exposed to pesticides and MPP+ (Fig. 1), we tested whether

dopamine neurons of *parkin* null flies are more sensitive to oxidative stress and mitochondrial complex I inhibition evoked by the same toxins. Dopamine-synthesizing neurons in whole *Drosophila* brains were identified with an antibody against tyrosine hydroxylase (TH). To maximize the chance of detecting a degenerative process, brains of survivors were analyzed when at least 50% of the *parkin*-deficient flies had died. Owing to the small size of the fly brain, TH-positive neurons could be directly counted in serial optical sections of whole-mount fly brains using confocal laser microscopy (Fig. 2A and B). *Drosophila* dopamine neurons are organized in several distinct clusters. We quantified the number of dopamine neurons of the protocerebral posterior lateral (PPL1, also named DL1) and protocerebral posterior medial (PPM, also named DM) clusters (Whitworth et al., 2005). However, no differences in the numbers of TH-positive neurons between wild type and *parkin* null flies exposed to the various toxins were observed (Fig. 2C and D).

Sequestration of redox-active trace metals prolongs the lifespan of *parkin* mutant flies

As discussed above, trace metals can be toxic when insufficiently sequestered and have been implicated in several neurodegenerative disorders ([Friedman et al., 2007], [Rhodes and Ritz, 2008], [Rossi et al., 2004], [Sayre et al., 1999], [Zatta et al., 2009] and [Zecca et al., 2004]). To investigate whether *parkin* deficiency may sensitize flies to endogenous iron and copper toxicity, *park25/25* *Drosophila* were raised on food supplemented with chelators of copper and iron. We used the copper-specific chelator bathocuproine disulfonate (BCS) and bathophenanthroline sulfonated sodium salt (BPS), which preferentially sequesters iron and to a lesser degree copper. Various concentrations of the chelators were tested. We found that both 0.1 mM BPS and 0.3 mM BCS significantly prolonged the median lifespan of *park25/25* flies, which was 5 days on normal food (Fig. 3A). In addition, the combination of BPS and BCS extended the median lifespan even more to 19 days (Fig. 3A). In contrast, metal chelation did not increase the lifespan of *park25/+* flies observed up to 50 days of age (data not shown). Because increased longevity has been associated with calorie restriction in several species including

Drosophila (Guarente and Picard, 2005), we wanted to exclude the possibility that the flies consumed reduced amounts of food containing metal chelators. For this purpose we carried out the adult gustatory assay and quantified for food uptake. This assay results in red abdomen coloration due to the chemical acid-red mixed into the food ([Bahadorani et al., 2008] and [Orgad et al., 1998]). As shown in Fig. 3B and C, there was no difference in the amount of food ingested by *park25/25* and *park25/+* flies grown on metal chelator-supplemented food, compared to normal food. This demonstrates that *parkin* mutant flies were not repelled from eating food containing metal chelators. Therefore, our results show that the increased lifespan of *parkin* mutant flies is due to metal chelation and suggest that *parkin* mutants are more sensitive to endogenous levels of the redox-active metals iron and copper, which have been implicated in PD ([Desai and Kaler, 2008], [Friedman et al., 2007], [Hirsch, 2006], [Wright et al., 2009] and [Zecca et al., 2004]). The mechanisms by which iron and copper chelators increased the lifespan of *parkin*-deficient flies are unknown, but possibly include reduced oxidative stress, as both iron and copper have the ability to accept and donate electrons. This can lead to formation of ROS and oxidative modification of proteins, lipids and other cell constituents ([Barnham and Bush, 2008] and [Halliwell and Gutteridge, 1990]). Consistent with this idea, heads of *parkin*-deficient flies raised on chelator-supplemented food showed a reduced level of ROS measured by 2',7'-dichlorofluorescein diacetate (DCF-DA), a cell-permeable non-fluorescent dye that turns highly fluorescent upon oxidation within cells (Fig. 3D). DCF fluorescence of isolated *park25/25* brains, as observed in the confocal microscope, was also significantly lower in presence of metal chelators (Fig. 3D, inset). To further probe an interaction of iron and copper with *parkin* deficiency, we exposed *park25/25* and *park25/+* flies to increased levels of iron and copper. Iron slightly but significantly reduced the lifespan of *park25/25* flies ($p < 0.01$) without exerting an effect on *park25/+* flies (Fig. 3E). In contrast, increasing the copper concentration in the food failed to shorten the lifespan of either genotype of flies (Fig. 3F). It is possible that uptake of iron and copper is subject to different regulation, and that excess copper in the food not necessarily results in increased copper uptake through the copper transporter (see Discussion).

Lack of Parkin aggravates the rough eye phenotype caused by overexpression of the Ctr1B copper importer

High affinity copper transporters are present in all eukaryotes including *Drosophila*, which encodes three members (Ctr1A, B and C) ([Dancis et al., 1994], Lee et al., 2000 J. Lee et al., Isolation of a murine copper transporter gene, tissue specific expression and functional complementation of a yeast copper transport mutant ([Lee et al., 2000], [Selvaraj et al., 2005], [Turski and Thiele, 2009] and [Zhou et al., 2003])). Ectopic overexpression of the *Drosophila* Ctr1B gene results in copper hypersensitivity leading to death when flies are grown on food with normal copper content (Selvaraj et al., 2005). To study whether *parkin* deficiency may sensitize to copper-mediated cell death in the eye, we overexpressed Ctr1B under the control of the eye-specific GMR promoter in *park25/+* and *park25/25* *Drosophila*. Control GMR-Ctr1B transgenic flies showed normal eyes when grown on regular food (Fig. 4A), but developed a rough eye phenotype in presence of 20 μ M copper (Fig. 4B). *park25/25* *Drosophila* also had normal eyes (Fig. 4C). In contrast, GMR-driven overexpression of Ctr1B in *park25/25* flies resulted in a rough eye phenotype, even on regular food (Fig. 4D). This phenotype was alleviated by copper scarcity and exacerbated by increased copper levels (data not shown). Importantly, the rough eye phenotype was not observed when Ctr1B was overexpressed in the eyes of *park25/+* flies (Fig. 4E). These results show that loss of Parkin and Ctr1B overexpression synergize in the induction of the rough eye phenotype, and suggest that *parkin* deficiency increases the vulnerability to increased copper import through Ctr1B. To investigate whether the phenotype was indeed specific for copper, *park25/25* flies overexpressing Ctr1B were raised on food supplemented with 20 μ M or 50 μ M silver nitrate (AgNO₃). Silver effectively competes with and inhibits copper import through the human transporter hCtr1 (Lee et al., 2002) but, in contrast to copper, does not promote oxidative stress by engaging in cellular redox reactions. Silver failed to cause a rough eye phenotype in *park25/25*

flies overexpressing Ctr1B (Fig. 4F–G), showing that this phenotype was specific to copper and likely depends on copper-mediated oxidative stress.

Beneficial effects of antioxidants and SOD1 overexpression on *parkin*-deficient flies

Because the studies described above suggested that reduced ROS production may, at least in part, be responsible for the increased lifespan of *parkin*-deficient flies raised on metal chelator-supplemented food (Fig. 3A and D), we further investigated the role of oxidative stress in the reduced longevity of *park25/25* flies and the effect of combined anti-oxidant and metal chelator treatment on the survival of *park25/25* flies. Cytosolic Cu/Zn superoxide dismutase (SOD1), catalase and glutathione reductase provide the first line of defense against ROS. In addition, the glutathione precursor N-acetylcysteine (NAC), which can be taken up by cells, reduces oxidative stress through its ability to scavenge free radicals. Ascorbic acid (vitamin C) is the most important water-soluble antioxidant and protects cellular DNA and low-density lipoproteins from oxidative damage by quenching free radicals (Duarte and Lunec, 2005). Both 15 mM NAC (Fig. 5A) and 100 mM ascorbic acid (Fig. 5B) significantly increased the median lifespan of *park25/25* flies from 5 days to 14 days (NAC) and 10 days (ascorbic acid). However, higher concentrations of ascorbic acid were ineffective and even shortened lifespan. Although this was unexpected, studies in humans have shown that optimal dosing of vitamin C is critical to achieve beneficial effects (Padayatty et al., 2003). Flies grown on food containing both NAC and metal chelators had a longer maximum lifespan (>38 days) than either flies grown on NAC-supplemented food alone (22 days) or metal chelators alone (30 days). However, the median lifespan of flies treated with NAC and metal chelators (15 days) was similar to flies treated with NAC alone (14 days), suggesting that the combination of both treatments provides no further benefit on overall survival (although it extends maximum lifespan). Collectively, these results suggest that the lifespan-extending effect of metal chelation is at least partially mediated through the reduction of oxidative stress. Next, we compared the survival of *parkin*-deficient flies with or without overexpression of SOD1 in

mesoderm-derived tissues. In this experiment, survival was monitored at the four specific ages indicated in Fig. 5C. SOD1-expressing *park25/25* flies up to 4 days of age did not show a difference in survival compared to non-transgenic *park25/25* flies (Fig. 5C). However, at 11 days of age only 10% of the *park25/25* flies were alive compared to 80% of the *park25/25* flies overexpressing SOD1. None of the *park25/25* flies survived to 17 days at which time point 80% of the SOD-1 expressing *park25/25* animals were still alive (Fig. 5C). In conclusion, both non-enzymatic and enzymatic reduction of oxidative stress and ROS led to increased survival of *park25/25* flies. The effect of SOD1 was particularly strong, perhaps because the *twi*-GAL4 driver results in enhanced SOD1 expression in flight muscles ([Estrada et al., 2007] and [Greig and Akam, 1993]), which underwent apoptotic degeneration in *parkin*-deficient *Drosophila* (Greene et al., 2003).

Loss of *parkin* leads to increased susceptibility to hypoxia and heat shock

It has been proposed that Parkin functions as a multipurpose neuroprotective agent in the defense against oxidative stress and the accumulation of specific Parkin substrates ([Feany and Pallanck, 2003] and [Moore, 2006]). In addition, *parkin* mutant flies were reported to be more sensitive to cold stress (Pesah et al., 2004). A recent study identified an E3 ligase complex comprised of Parkin and two other recessive PD-associated proteins, PINK1 and DJ-1 (PPD complex), whose function is to degrade unfolded and misfolded proteins (Xiong et al., 2009). In order to analyze whether Parkin may also protect against non-oxidative stressors and conditions that induce global protein denaturation and aggregation, we exposed *park25/25* and *park25/+* flies to transient hypoxia and heat shock. The cellular response to hypoxia is conserved in *Drosophila* (Lavista-Llanos et al., 2002) and the recovery from oxygen deprivation is significantly delayed by mutations in *hypnos* genes (Haddad et al., 1997), one of which encodes an RNA-directed adenosine deaminase (ADAR) (Ma et al., 2001). ADAR is known to edit the transcripts of at least three fly ion channels, suggesting that regulation of specific ion channels might be important for recovery from hypoxia (Ma et al., 2001). Interestingly, the recovery of *park25/25* flies from a 7-min exposure to 0.5%

oxygen in a hypoxia chamber was also significantly delayed when compared to *park25/+* control flies (Fig. 6A). In addition, a heat shock for 30 or 60 min at 38 °C resulted in dramatically increased mortality in the absence of Parkin, with only 15% and 7% of the *park25/25* flies surviving, while at least 95% of *park25/+* control flies survived the same treatments (Fig. 6B). These results suggest that the protective function of Parkin extends to non-oxidative stressors and conditions that impair global protein homeostasis.

Discussion

Here, we addressed whether the lack of Parkin sensitizes flies to environmental pesticides, mitochondrial toxins and metals that have been implicated as risk factors in the etiology of Parkinson's disease. Earlier studies in *Drosophila* had shown that *parkin* ablation increased the vulnerability of flies to paraquat (Pesah et al., 2004), but other toxins and metals have not been studied previously. Our experiments confirm the reduced lifespan after exposure to paraquat and show that Parkin-deficient flies are also more sensitive to the mitochondrial complex I inhibitors rotenone and MPP⁺. In rats, chronic and systemic exposure to rotenone reproduces features of PD, including nigrostriatal degeneration and α -synuclein-positive cytosolic inclusions ([Betarbet et al., 2000] and [Sherer et al., 2003]). *Drosophila* expressing mutant forms of LRRK2 exhibited late-onset dopamine neuron loss that was aggravated by rotenone (Ng et al., 2009). Interestingly, co-expression of human Parkin conferred significant protection against rotenone-induced dopamine neuron loss in LRRK2 G2019S transgenic flies, suggesting that Parkin reversed the increased rotenone sensitivity caused by mutant LRRK2 (Ng et al., 2009). MPP⁺ is the toxic metabolite of MPTP, which caused an acute and permanent state of Parkinsonism in humans ([Langston et al., 1983] and [Langston and Ballard, 1983]). This discovery subsequently led to the development of MPTP-based PD models in monkeys and mice ([Burns et al., 1983] and [Heikkilä et al., 1984]). While both rotenone and MPP⁺ significantly shortened the lifespan of *park25/25 Drosophila*, we failed to detect an adverse effect of these chemicals on the survival of dopamine-synthesizing (tyrosine hydroxylase-positive) neurons in these flies. The same

was true for paraquat that is known to produce nigrostriatal dysfunction and degenerative changes reminiscent of PD in mice ([Brooks et al., 1999], [Manning-Bog et al., 2002] and [McCormack et al., 2002]). Although no loss of TH-positive neurons was observed in *park25/25* flies, we cannot exclude the possibility that dopamine neuron death may occur after longer toxin exposure in *parkin* mutant flies. Limited dopamine neuron death restricted to the PPL1 cluster has been reported in 20-day-old *parkin* mutant flies (*park25* allele) (Whitworth et al., 2005). However, the original characterization of the same flies showed that dopamine neuron numbers were normal, even in 30-day-old *Drosophila* (Greene et al., 2003). In addition, although we used the same flies, the maximum lifespan of *park25/25* flies was only 10–13 days in different experiments in our hands (see Figs. 1C, 3A, E, 5A–B), preventing us from analyzing dopamine neuron numbers at later ages. Alternatively, feeding the toxins may not result in their accumulation in the brain at concentrations sufficiently high to affect neurons. In this regard, it is noteworthy that flies with a knockout of DJ-1, another gene associated with Parkinsonism, also displayed a shortened lifespan but no dopamine neuron death when exposed to paraquat and rotenone ([Menzies et al., 2005] and [Meulener et al., 2005]). Collectively, these results show that the dopaminergic system of flies is largely insensitive to paraquat, rotenone and MPP⁺ (at least under the conditions used here), but that lifespan determination is a feasible strategy to assess the effects of a specific gene ablation on susceptibility to PD-associated toxins and other factors. Therefore, in all subsequent studies we focused on the survival of *parkin*-deficient flies compared to *park25/+* *Drosophila* to determine whether a given treatment affected the flies in a *parkin*-dependent manner.

Iron and copper have been implicated in the pathogenesis of various neurodegenerative disorders ([Chinta and Andersen, 2008], [Desai and Kaler, 2008], [Rhodes and Ritz, 2008], [Waggoner et al., 1999], [Zatta et al., 2009] and [Zecca et al., 2004]). The total iron content is increased in the substantia nigra in PD, and a shift in the Fe³⁺/Fe²⁺ ratio with a higher level of reactive labile Fe²⁺ may contribute to increased oxidative stress in PD (Berg and Hochstrasser, 2006). In our experiments, we found that *park25/25* but not

park25/+ flies showed decreased lifespan when grown on iron-supplemented food. Moreover, addition of the iron-chelating agent BPS to food containing normal levels of iron increased the lifespan of *parkin*-deficient flies. Likewise, mixing the copper-chelating agent BCS into the food significantly increased the lifespan of *park25/25* flies, and the combination of both BCS and BPS extended lifespan even more. In contrast, *park25/+* flies showed similar survival curves up to 50 days of age when grown on either normal or chelator (BPS + BCS) -supplemented food (data not shown), indicating that sequestration of free iron and copper benefited flies only in the absence of Parkin. Given that sequestration of copper extended the lifespan of *park25/25 Drosophila*, we were surprised to find that increased concentrations of copper in the food did not impair survival of *parkin*-deficient flies. It is conceivable that the uptake of copper is tightly regulated and not proportional to the amount of copper added to the food. In fact, copper uptake by the human Ctr1 transporter (hCtr1) was saturated at about 25 μ M copper (Lee et al., 2002), a concentration that is 20 times lower than we used in the copper feeding experiment (Fig. 3E). In contrast, overexpression of the copper transporter Ctr1B in the eye of *park25/25* flies resulted in a rough eye phenotype even on food with normal copper concentration, while this phenotype was not observed in *park25/+* controls. Because it has been shown that overexpression of hCtr1 increases copper uptake even at low copper concentrations (Lee et al., 2002), these results suggest that the uptake, rather than availability of copper in the food, may be limiting for toxicity in *park25/25* flies. Compared to *park25/+* control flies, *park25/25* animals showed significantly increased levels of ROS in the head/brain, as measured by DCF fluorescence. Rearing *park25/25* flies on food containing iron- and copper-chelating agents decreased the ROS levels, suggesting that reduction of oxidative stress through sequestration of redox-active iron and copper may be responsible for the lifespan extension observed under these conditions. In agreement with this idea, transcriptional profiling of *Drosophila parkin* mutants revealed that genes of the oxidative stress response are up-regulated (Greene et al., 2005). Furthermore, the phenotype of *parkin*-deficient flies was exacerbated by loss-of-function alleles in genes that limit oxidative damage, such as glutathione S-transferase S1 (GstS1), thioredoxin (Trx) and Rlu-A1,

which is involved in the synthesis of the anti-oxidant vitamin B2 (Greene et al., 2005). Conversely, increased glutathione S-transferase activity rescued neuron loss in a different line of *parkin* mutant flies (Whitworth et al., 2005). Finally, our own studies described here corroborate that oxidative stress is responsible for the reduced longevity of *park25/25* flies. Both N-acetylcysteine and ascorbic acid (vitamin C) supplementation significantly increased the lifespan of *park25/25* flies. In addition, overexpression of SOD1 using the mesoderm-specific driver *twi-GAL4* dramatically prolonged the survival of *parkin* mutant flies, possibly by enhancing SOD1 expression in muscle tissue, which was affected by apoptotic degeneration in *park25/25* flies (Greene et al., 2003). Combining N-acetyl cysteine with metal chelators extended the maximum but not the median lifespan of *park25/25* flies, suggesting that metal sequestration extends lifespan, at least in part, by reducing oxidative stress. This is consistent with the ability of the iron- and copper-chelating agents to reduce ROS levels in the brain of *parkin*-deficient flies (Fig. 3D).

Recent evidence has placed Parkin in a common pathway with PINK1 in the maintenance of mitochondrial function and integrity ([Clark et al., 2006], [Park et al., 2006] and [Yang et al., 2006]). Mitochondrial dysfunction and, in *Drosophila*, degeneration have been observed in the absence of either protein (Bueler, 2009). However, a unique feature of Parkin is its ability to associate with dysfunctional mitochondria and promote their degradation through mitophagy (Narendra et al., 2008), suggesting that impaired removal of defective mitochondria in the absence of Parkin is pathogenic (Narendra et al., 2009). As a consequence, endogenous and exogenous factors that impose additional stress on mitochondria may result in further accumulation of dysfunctional organelles in Parkin-deficient cells, which could increase toxicity. Both rotenone and MPP⁺ are potent inhibitors of mitochondrial complex I. In addition, iron is an essential component of many mitochondrial proteins (iron-sulfur cluster proteins) that are readily oxidized, and copper-mediated oxidative stress has been linked to mitochondrial dysfunction in neurodegenerative disorders associated with copper imbalance (Rossi et al., 2004). In line with this, treatment of dopaminergic neurons with copper led to

mitochondrial dysfunction, decreased dopamine content and cell death (Shi et al., 2008). Thus, the various toxins and metals tested in this study may further accelerate and enhance mitochondrial dysfunction and degeneration, overwhelming the mitophagy system that is already compromised in *parkin*-deficient flies. To study whether chelator treatment mitigated mitochondrial degeneration in *park25/25* flies, we examined mitochondrial morphology in sections of the indirect flight muscles by electron microscopy. However, we failed to detect a rescue of swollen mitochondria with disintegrated cristae after chelator treatment (data not shown), showing that the life-extending effect of metal chelation is likely not due to improved mitochondrial structure, but rather due to reduction of oxidative stress that may occur downstream of mitochondrial dysfunction in *park25/25 Drosophila*. In this context, it is worth mentioning that food supplementation with zinc, a non-redox-active metal that under most conditions functions as an antioxidant, exerts a beneficial effect on *parkin* mutant flies (Saini and Schaffner, 2010). Future studies with flies expressing mitochondrial-targeted GFP (Hollenbeck and Saxton, 2005) may be interesting to investigate whether metal chelators can improve other aspects of mitochondrial biology, such as transport and dynamics, in *parkin* mutants.

Finally, our studies show that *park25/25* flies are more sensitive to hypoxia and heat shock. Interestingly, the three recessive PD proteins Parkin, PINK1 and DJ-1 associate to form a functional E3 ligase complex (PPD complex) that promotes the degradation of Parkin substrates as well as unfolded and misfolded proteins after heat shock (Xiong et al., 2009). This strongly suggests that the drastically reduced survival of *park25/25* flies after heat shock is due to impaired removal of misfolded and aggregated proteins that may accumulate under these conditions.

Our studies may have therapeutic implications. The extension of lifespan in *park25/25* flies by a combination of different metal-chelating agents hints towards a possible strategy of reducing oxidative stress in familial PD, where carriers of PARK2 (Parkin) mutations could be identified by genetic screening and treatment may be administered early to inhibit or delay dopamine neuron death. PARK2 mutations are by far the most common in recessive

Parkinsonism ([Abbas et al., 1999] and [Gasser, 2005]), accounting for 49–63% of early-onset familial cases ([Lohmann et al., 2003] and [Lucking et al., 2000]). In addition, PARK2 mutations are found in 19% of isolated PD cases with onset before 55 years ([Lohmann et al., 2003] and [Periquet et al., 2003]). Thus, metal sequestration might benefit a significant number of patients. A previous study showed that the orally administered iron chelator clioquinol was well tolerated in mice, where it resulted in a partial protection against MPTP-induced dopamine neuron degeneration in a model of sporadic PD (Kaur et al., 2003). Our experiments did not directly address whether chelators, in addition to reducing ROS, rescued muscle cell apoptosis (Greene et al., 2003) or non-dopaminergic neuronal cell loss that might occur in the brain of *parkin*-deficient flies. Such studies will be important in the future to refine the mechanisms of metal chelator-mediated lifespan extension in *parkin* mutant flies. In addition, studies in “familial PD mice” that show frank dopamine neuron loss (Lu et al., 2009) will be necessary to test whether metal chelation indeed protects against dopamine neuron death caused by Parkin mutations.

Acknowledgments

We thank Drs. Jessica Greene and Leo Pallanck (University of Washington) for providing the park25/25 flies, Christian Frei (ETH Zurich) for help with the hypoxia experiments, Ivan Ostojic (FMI Basel) for statistical analyses, and Till Strassen for maintenance of fly stocks. This work was supported by grants of the Kanton Zürich and the Swiss National Science Foundation (to W.S.) as well as the Bonizzi-Theler Foundation and Stammbach Foundation (to H.B.).

References

Abbas et al., 1999 N. Abbas et al., A wide variety of mutations in the parkin gene are responsible for autosomal recessive parkinsonism in Europe. French Parkinson's Disease Genetics Study Group and the European Consortium on Genetic Susceptibility in Parkinson's Disease, Hum. Mol. Genet. 8 (1999), pp. 567–574.

- Abbott et al., 2003 R.D. Abbott et al., Environmental, life-style, and physical precursors of clinical Parkinson's disease: recent findings from the Honolulu-Asia Aging Study, *J. Neurol.* 250 (Suppl 3) (2003), pp. III30–III39.
- Ala et al., 2007 A. Ala et al., Wilson's disease, *Lancet* 369 (2007), pp. 397–408.
- Bahadorani et al., 2008 S. Bahadorani et al., The effects of vitamin supplementation on *Drosophila* life span under normoxia and under oxidative stress, *J. Gerontol. A Biol. Sci. Med. Sci.* 63 (2008), pp. 35–42.
- Baldi et al., 2003 I. Baldi et al., Association between Parkinson's disease and exposure to pesticides in southwestern France, *Neuroepidemiology* 22 (2003), pp. 305–310.
- Bando et al., 2005 Y. Bando et al., RA410/Sly1 suppresses MPP⁺ and 6-hydroxydopamine-induced cell death in SH-SY5Y cells, *Neurobiol. Dis.* 18 (2005), pp. 143–151.
- Barnham and Bush, 2008 K.J. Barnham and A.I. Bush, Metals in Alzheimer's and Parkinson's diseases, *Curr. Opin. Chem. Biol.* 12 (2008), pp. 222–228.
- Barthel et al., 2003 H. Barthel et al., Concordant pre- and postsynaptic deficits of dopaminergic neurotransmission in neurologic Wilson disease, *AJNR Am. J. Neuroradiol.* 24 (2003), pp. 234–238.
- Berg and Hochstrasser, 2006 D. Berg and H. Hochstrasser, Iron metabolism in Parkinsonian syndromes, *Mov. Disord.* 21 (2006), pp. 1299–1310.
- Betarbet et al., 2000 R. Betarbet et al., Chronic systemic pesticide exposure reproduces features of Parkinson's disease, *Nat. Neurosci.* 3 (2000), pp. 1301–1306.
- Brooks et al., 1999 A.I. Brooks et al., Paraquat elicited neurobehavioral syndrome caused by dopaminergic neuron loss, *Brain Res.* 823 (1999), pp. 1–10.
- Bueler, 2009 H. Bueler, Impaired mitochondrial dynamics and function in the pathogenesis of Parkinson's disease, *Exp. Neurol.* 218 (2009), pp. 235–246.
- Burns et al., 1983 R.S. Burns et al., A primate model of parkinsonism: selective destruction of dopaminergic neurons in the pars compacta of the substantia nigra by N-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine, *Proc. Natl. Acad. Sci. USA* 80 (1983), pp. 4546–4550.

Chinta and Andersen, 2008 S.J. Chinta and J.K. Andersen, Redox imbalance in Parkinson's disease, *Biochim. Biophys. Acta* 1780 (2008), pp. 1362–1367.

Clark et al., 2006 I.E. Clark et al., *Drosophila pink1* is required for mitochondrial function and interacts genetically with parkin, *Nature* 441 (2006), pp. 1162–1166.

Connor et al., 1995 J.R. Connor et al., A quantitative analysis of isoferitins in select regions of aged, parkinsonian, and Alzheimer's diseased brains, *J. Neurochem.* 65 (1995), pp. 717–724.

Cookson, 2005 M.R. Cookson, The biochemistry of Parkinson's disease, *Annu. Rev. Biochem.* 74 (2005), pp. 29–52.

Corti et al., 2003 O. Corti et al., The p38 subunit of the aminoacyl-tRNA synthetase complex is a Parkin substrate: linking protein biosynthesis and neurodegeneration, *Hum. Mol. Genet.* 12 (2003), pp. 1427–1437.

Coulom and Birman, 2004 H. Coulom and S. Birman, Chronic exposure to rotenone models sporadic Parkinson's disease in *Drosophila melanogaster*, *J. Neurosci.* 24 (2004), pp. 10993–10998.

Dancis et al., 1994 A. Dancis et al., Molecular characterization of a copper transport protein in *S. cerevisiae*: an unexpected role for copper in iron transport, *Cell* 76 (1994), pp. 393–402.

Dauer and Przedborski, 2003 W. Dauer and S. Przedborski, Parkinson's disease: mechanisms and models, *Neuron* 39 (2003), pp. 889–909

Dawson and Dawson, 2003 T.M. Dawson and V.L. Dawson, Molecular pathways of neurodegeneration in Parkinson's disease, *Science* 302 (2003), pp. 819–822.

Desai and Kaler, 2008 V. Desai and S.G. Kaler, Role of copper in human neurological disorders, *Am. J. Clin. Nutr.* 88 (2008), pp. 855S–858S.

Dexter et al., 1992 D.T. Dexter et al., Alterations in levels of iron, ferritin, and other trace metals in neurodegenerative diseases affecting the basal ganglia. The Royal Kings and Queens Parkinson's Disease Research Group, *Ann. Neurol.* 32 (1992), pp. S94–S100 Suppl.

Duarte and Lunec, 2005 T.L. Duarte and J. Lunec, Review: Part of the Series: From Dietary Antioxidants to Regulators in Cellular Signalling and Gene Expression: Review: when is an antioxidant not an antioxidant? A review of

novel actions and reactions of vitamin C, *Free Radic. Res.* 39 (2005), pp. 671–686.

Estrada et al., 2007 B. Estrada et al., The MARVEL domain protein, Singles Bar, is required for progression past the pre-fusion complex stage of myoblast fusion, *Dev. Biol.* 307 (2007), pp. 328–339.

Feany and Bender, 2000 M.B. Feany and W.W. Bender, A *Drosophila* model of Parkinson's disease, *Nature* 404 (2000), pp. 394–398.

Feany and Pallanck, 2003 M.B. Feany and L.J. Pallanck, Parkin: a multipurpose neuroprotective agent?, *Neuron* 38 (2003), pp. 13–16.

Friedman et al., 2007 A. Friedman et al., Iron as a trigger of neurodegeneration in Parkinson's disease, *Handb. Clin. Neurol.* 83 (2007), pp. 493–505.

Fukae et al., 2009 J. Fukae et al., Programmed cell death-2 isoform1 is ubiquitinated by parkin and increased in the substantia nigra of patients with autosomal recessive Parkinson's disease, *FEBS Lett.* 583 (2009), pp. 521–525.

Gasser, 2005 T. Gasser, Genetics of Parkinson's disease, *Curr. Opin. Neurol.* 18 (2005), pp. 363–369.

Gasser, 2009 T. Gasser, Mendelian forms of Parkinson's disease, *Biochim. Biophys. Acta* 1792 (2009), pp. 587–596.

Goldberg et al., 2003 M.S. Goldberg et al., Parkin-deficient mice exhibit nigrostriatal deficits but not loss of dopaminergic neurons, *J. Biol. Chem.* 278 (2003), pp. 43628–43635.

Gorell et al., 1997 J.M. Gorell et al., Occupational exposures to metals as risk factors for Parkinson's disease, *Neurology* 48 (1997), pp. 650–658.

Gorell et al., 1998 J.M. Gorell et al., The risk of Parkinson's disease with exposure to pesticides, farming, well water, and rural living, *Neurology* 50 (1998), pp. 1346–1350.

Greene et al., 2005 J.C. Greene et al., Genetic and genomic studies of *Drosophila* parkin mutants implicate oxidative stress and innate immune responses in pathogenesis, *Hum. Mol. Genet.* 14 (2005), pp. 799–811

Greene et al., 2003 J.C. Greene et al., Mitochondrial pathology and apoptotic muscle degeneration in *Drosophila* parkin mutants, *Proc. Natl. Acad. Sci. USA* 100 (2003), pp. 4078–4083.

- Greig and Akam, 1993 S. Greig and M. Akam, Homeotic genes autonomously specify one aspect of pattern in the *Drosophila* mesoderm, *Nature* 362 (1993), pp. 630–632.
- Guarente and Picard, 2005 L. Guarente and F. Picard, Calorie restriction—the SIR2 connection, *Cell* 120 (2005), pp. 473–482.
- Haddad et al., 1997 G.G. Haddad et al., Genetic basis of tolerance to O₂ deprivation in *Drosophila melanogaster*, *Proc. Natl. Acad. Sci. USA* 94 (1997), pp. 10809–10812.
- Halliwell and Gutteridge, 1990 B. Halliwell and J.M. Gutteridge, Role of free radicals and catalytic metal ions in human disease: an overview, *Methods Enzymol.* 186 (1990), pp. 1–85.
- Hardy et al., 2009 J. Hardy et al., The genetics of Parkinson's syndromes: a critical review, *Curr. Opin. Genet. Dev.* 19 (2009), pp. 254–265.
- Harrison and Arosio, 1996 P.M. Harrison and P. Arosio, The ferritins: molecular properties, iron storage function and cellular regulation, *Biochim. Biophys. Acta* 1275 (1996), pp. 161–203.
- Hatano et al., 2009 T. Hatano et al., Pathogenesis of familial Parkinson's disease: new insights based on monogenic forms of Parkinson's disease, *J. Neurochem.* 111 (2009), pp. 1075–1093.
- Heikkila et al., 1984 R.E. Heikkila et al., Dopaminergic neurotoxicity of 1-methyl-4-phenyl-1, 2, 5, 6-tetrahydropyridine in mice, *Science* 224 (1984), pp. 1451–1453.
- Heisenberg and Böhl, 1979 M. Heisenberg and K. Böhl, Isolation of anatomical brain mutants of *Drosophila* by histological means, *Z. Naturforsch.* 34 (1979), pp. 143–147.
- Henchcliffe and Beal, 2008 C. Henchcliffe and M.F. Beal, Mitochondrial biology and oxidative stress in Parkinson disease pathogenesis, *Nat. Clin. Pract. Neurol.* 4 (2008), pp. 600–609.
- Hirsch, 2006 E.C. Hirsch, Altered regulation of iron transport and storage in Parkinson's disease, *J. Neural Transm. Suppl.* (2006), pp. 201–204.
- Hollenbeck and Saxton, 2005 P.J. Hollenbeck and W.M. Saxton, The axonal transport of mitochondria, *J. Cell Sci.* 118 (2005), pp. 5411–5419.
- Hosamani and Muralidhara, 2009 R. Hosamani and Muralidhara, Neuroprotective efficacy of *Bacopa monnieri* against rotenone induced

oxidative stress and neurotoxicity in *Drosophila melanogaster*, *Neurotoxicology* 30 (2009), pp. 977–985.

Imai et al., 2001 Y. Imai et al., An unfolded putative transmembrane polypeptide, which can lead to endoplasmic reticulum stress, is a substrate of Parkin, *Cell* 105 (2001), pp. 891–902.

Itier et al., 2003 J.M. Itier et al., Parkin gene inactivation alters behaviour and dopamine neurotransmission in the mouse, *Hum. Mol. Genet.* 12 (2003), pp. 2277–2291.

Jones et al., 2009 M.A. Jones et al., A forward genetic screen in *Drosophila* implicates insulin signaling in age-related locomotor impairment, *Exp. Gerontol.* 44 (2009), pp. 532–540.

Kao, 2009a S.Y. Kao, DNA damage induces nuclear translocation of parkin, *J. Biomed. Sci.* 16 (2009), p. 67.

Kao, 2009b S.Y. Kao, Regulation of DNA repair by parkin, *Biochem. Biophys. Res. Commun.* 382 (2009), pp. 321–325.

Kaur et al., 2003 D. Kaur et al., Genetic or pharmacological iron chelation prevents MPTP-induced neurotoxicity in vivo: a novel therapy for Parkinson's disease, *Neuron* 37 (2003), pp. 899–909.

Kitada et al., 1998 T. Kitada et al., Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism, *Nature* 392 (1998), pp. 605–608.

Koziorowski et al., 2007 D. Koziorowski et al., ELISA reveals a difference in the structure of substantia nigra ferritin in Parkinson's disease and incidental Lewy body compared to control, *Parkinsonism Relat. Disord.* 13 (2007), pp. 214–218.

Langston et al., 1983 J.W. Langston et al., Chronic Parkinsonism in humans due to a product of meperidine-analog synthesis, *Science* 219 (1983), pp. 979–980.

Langston and Ballard, 1983 J.W. Langston and P.A. Ballard Jr., Parkinson's disease in a chemist working with 1-methyl-4-phenyl-1, 2, 5, 6-tetrahydropyridine, *N Engl J. Med.* 309 (1983), p. 310.

Lavista-Llanos et al., 2002 S. Lavista-Llanos et al., Control of the hypoxic response in *Drosophila melanogaster* by the basic helix-loop-helix PAS protein similar, *Mol. Cell. Biol.* 22 (2002), pp. 6842–6853.

- Lee et al., 2002 J. Lee et al., Biochemical characterization of the human copper transporter Ctr1, *J. Biol. Chem.* 277 (2002), pp. 4380–4387.
- Lee et al., 2000 J. Lee et al., Isolation of a murine copper transporter gene, tissue specific expression and functional complementation of a yeast copper transport mutant, *Gene* 254 (2000), pp. 87–96.
- Lee et al., 2009 K.S. Lee et al., JNK/FOXO-mediated neuronal expression of fly homologue of peroxiredoxin II reduces oxidative stress and extends life span, *J. Biol. Chem.* 284 (2009), pp. 29454–29461.
- Lohmann et al., 2003 E. Lohmann et al., How much phenotypic variation can be attributed to parkin genotype?, *Ann. Neurol.* 54 (2003), pp. 176–185.
- Lu et al., 2009 X.H. Lu et al., Bacterial artificial chromosome transgenic mice expressing a truncated mutant parkin exhibit age-dependent hypokinetic motor deficits, dopaminergic neuron degeneration, and accumulation of proteinase K-resistant alpha-synuclein, *J. Neurosci.* 29 (2009), pp. 1962–1976.
- Lucking et al., 2000 C.B. Lucking et al., Association between early-onset Parkinson's disease and mutations in the parkin gene, *N Engl J. Med.* 342 (2000), pp. 1560–1567.
- Ma et al., 2001 E. Ma et al., Mutation in pre-mRNA adenosine deaminase markedly attenuates neuronal tolerance to O₂ deprivation in *Drosophila melanogaster*, *J. Clin. Invest.* 107 (2001), pp. 685–693.
- Manning-Bog et al., 2002 A.B. Manning-Bog et al., The herbicide paraquat causes up-regulation and aggregation of alpha-synuclein in mice: paraquat and alpha-synuclein, *J. Biol. Chem.* 277 (2002), pp. 1641–1644.
- McCormack et al., 2002 A.L. McCormack et al., Environmental risk factors and Parkinson's disease: selective degeneration of nigral dopaminergic neurons caused by the herbicide paraquat, *Neurobiol. Dis.* 10 (2002), pp. 119–127.
- Menzies et al., 2005 F.M. Menzies et al., Roles of *Drosophila* DJ-1 in survival of dopaminergic neurons and oxidative stress, *Curr. Biol.* 15 (2005), pp. 1578–1582.
- Meulener et al., 2005 M. Meulener et al., *Drosophila* DJ-1 mutants are selectively sensitive to environmental toxins associated with Parkinson's disease, *Curr. Biol.* 15 (2005), pp. 1572–1577.

- Moore, 2006 D.J. Moore, Parkin: a multifaceted ubiquitin ligase, *Biochem. Soc. Trans.* 34 (2006), pp. 749–753.
- Moore et al., 2005 D.J. Moore et al., Molecular pathophysiology of Parkinson's disease, *Annu. Rev. Neurosci.* 28 (2005), pp. 57–87.
- Murakami et al., 2004 T. Murakami et al., Pael-R is accumulated in Lewy bodies of Parkinson's disease, *Ann. Neurol.* 55 (2004), pp. 439–442.
- Narendra et al., 2008 D. Narendra et al., Parkin is recruited selectively to impaired mitochondria and promotes their autophagy, *J. Cell Biol.* 183 (2008), pp. 795–803.
- Narendra et al., 2009 D. Narendra et al., Parkin-induced mitophagy in the pathogenesis of Parkinson disease, *Autophagy* 5 (2009), pp. 706–708.
- Ng et al., 2009 C.-H. Ng et al., Parkin protects against LRRK2 G2019S mutant-induced dopaminergic neurodegeneration in *Drosophila*, *J. Neurosci.* 29 (2009), pp. 11257–11262.
- Oder et al., 1993 W. Oder et al., Wilson's disease: evidence of subgroups derived from clinical findings and brain lesions, *Neurology* 43 (1993), pp. 120–124.
- Orgad et al., 1998 S. Orgad et al., Metal ions suppress the abnormal taste behavior of the *Drosophila* mutant *malvolio*, *J. Exp. Biol.* 201 (1998), pp. 115–120.
- Orino et al., 2001 K. Orino et al., Ferritin and the response to oxidative stress, *Biochem. J.* 357 (2001), pp. 241–247.
- Padayatty et al., 2003 S.J. Padayatty et al., Vitamin C as an antioxidant: evaluation of its role in disease prevention, *J. Am. Coll. Nutr.* 22 (2003), pp. 18–35.
- Palacino et al., 2004 J.J. Palacino et al., Mitochondrial dysfunction and oxidative damage in parkin-deficient mice, *J. Biol. Chem.* 279 (2004), pp. 18614–18622.
- Park et al., 2006 J. Park et al., Mitochondrial dysfunction in *Drosophila* PINK1 mutants is complemented by parkin, *Nature* 441 (2006), pp. 1157–1161.
- Periquet et al., 2005 M. Periquet et al., Proteomic analysis of parkin knockout mice: alterations in energy metabolism, protein handling and synaptic function, *J. Neurochem.* 95 (2005), pp. 1259–1276.

Periquet et al., 2003 M. Periquet et al., Parkin mutations are frequent in patients with isolated early-onset parkinsonism, *Brain* 126 (2003), pp. 1271–1278.

Pesah et al., 2004 Y. Pesah et al., *Drosophila* parkin mutants have decreased mass and cell size and increased sensitivity to oxygen radical stress, *Development* 131 (2004), pp. 2183–2194

Pesah et al., 2005 Y. Pesah et al., Whole mount analysis reveals normal numbers of dopaminergic neurons following misexpression of alpha-synuclein in *Drosophila*, *Genesis* 41 (2005), pp. 154–159.

Petrovitch et al., 2002 H. Petrovitch et al., Plantation work and risk of Parkinson disease in a population-based longitudinal study, *Arch. Neurol.* 59 (2002), pp. 1787–1792.

Rhodes and Ritz, 2008 S.L. Rhodes and B. Ritz, Genetics of iron regulation and the possible role of iron in Parkinson's disease, *Neurobiol. Dis.* 32 (2008), pp. 183–195.

Rossi et al., 2004 L. Rossi et al., Mitochondrial dysfunction in neurodegenerative diseases associated with copper imbalance, *Neurochem. Res.* 29 (2004), pp. 493–504.

Rothfuss et al., 2009 O. Rothfuss et al., Parkin protects mitochondrial genome integrity and supports mitochondrial DNA repair, *Hum. Mol. Genet.* 18 (2009), pp. 3832–3850.

Saini and Schaffner, 2010 N. Saini and W. Schaffner, Zinc supplement greatly improves the condition of parkin mutant *Drosophila*, *Biol. Chem.* 391 (2010), pp. 513–518.

Sayre et al., 1999 L.M. Sayre et al., Redox metals and neurodegenerative disease, *Curr. Opin. Chem. Biol.* 3 (1999), pp. 220–225.

Schriner et al., 2009 S.E. Schriner et al., Decreased mitochondrial superoxide levels and enhanced protection against paraquat in *Drosophila melanogaster* supplemented with *Rhodiola rosea*, *Free Radic. Res.* 43 (2009), pp. 836–843.

Selvaraj et al., 2005 A. Selvaraj et al., Metal-responsive transcription factor (MTF-1) handles both extremes, copper load and copper starvation, by activating different genes, *Genes Dev.* 19 (2005), pp. 891–896

- Shen and Cookson, 2004 J. Shen and M.R. Cookson, Mitochondria and dopamine: new insights into recessive parkinsonism, *Neuron* 43 (2004), pp. 301–304.
- Sherer et al., 2003 T.B. Sherer et al., Subcutaneous rotenone exposure causes highly selective dopaminergic degeneration and alpha-synuclein aggregation, *Exp. Neurol.* 179 (2003), pp. 9–16.
- Shi et al., 2008 L.M. Shi et al., Mitochondria dysfunction was involved in copper-induced toxicity in MES23.5 cells, *Neurosci. Bull.* 24 (2008), pp. 79–83.
- Shimura et al., 2000 H. Shimura et al., Familial Parkinson disease gene product, parkin, is a ubiquitin-protein ligase, *Nat. Genet.* 25 (2000), pp. 302–305.
- Sudmeyer et al., 2006 M. Sudmeyer et al., Wilson's disease tremor is associated with magnetic resonance imaging lesions in basal ganglia structures, *Mov. Disord.* 21 (2006), pp. 2134–2139.
- Turski and Thiele, 2009 M.L. Turski and D.J. Thiele, New roles for copper metabolism in cell proliferation, signaling, and disease, *J. Biol. Chem.* 284 (2009), pp. 717–721.
- Waggoner et al., 1999 D.J. Waggoner et al., The role of copper in neurodegenerative disease, *Neurobiol. Dis.* 6 (1999), pp. 221–230.
- Wang et al., 2007 C. Wang et al., Drosophila overexpressing parkin R275W mutant exhibits dopaminergic neuron degeneration and mitochondrial abnormalities, *J. Neurosci.* 27 (2007), pp. 8563–8570.
- Wang et al., 2008 D. Wang et al., Dispensable role of Drosophila ortholog of LRRK2 kinase activity in survival of dopaminergic neurons, *Mol. Neurodegener.* 3 (2008), p. 3.
- Whitworth et al., 2005 A.J. Whitworth et al., Increased glutathione S-transferase activity rescues dopaminergic neuron loss in a Drosophila model of Parkinson's disease, *Proc. Natl. Acad. Sci. USA* 102 (2005), pp. 8024–8029.
- Wright et al., 2009 J.A. Wright et al., Unique copper-induced oligomers mediate alpha-synuclein toxicity, *FASEB J.* 23 (2009), pp. 2384–2393.

- Xiong et al., 2009 H. Xiong et al., Parkin, PINK1, and DJ-1 form a ubiquitin E3 ligase complex promoting unfolded protein degradation, *J. Clin. Invest.* 119 (2009), pp. 650–660.
- Yang et al., 2006 Y. Yang et al., Mitochondrial pathology and muscle and dopaminergic neuron degeneration caused by inactivation of *Drosophila* Pink1 is rescued by Parkin, *Proc. Natl. Acad. Sci. USA* 103 (2006), pp. 10793–10798.
- Zatta et al., 2009 P. Zatta et al., Alzheimer's disease, metal ions and metal homeostatic therapy, *Trends Pharmacol. Sci.* 30 (2009), pp. 346–355.
- Zecca et al., 2004 L. Zecca et al., Iron, brain ageing and neurodegenerative disorders, *Nat. Rev. Neurosci.* 5 (2004), pp. 863–873.
- Zhang et al., 2000 Y. Zhang et al., Parkin functions as an E2-dependent ubiquitin- protein ligase and promotes the degradation of the synaptic vesicle-associated protein, CDCrel-1, *Proc. Natl. Acad. Sci. USA* 97 (2000), pp. 13354–13359.
- Zhou et al., 2003 H. Zhou et al., A copper-regulated transporter required for copper acquisition, pigmentation, and specific stages of development in *Drosophila melanogaster*, *J. Biol. Chem.* 278 (2003), pp. 48210–48218.




Fig. 1. Lack of Parkin increases sensitivity of flies to environmental toxins and MPP⁺. Twenty 1- to 2-day-old flies of each genotype were grown on food supplemented with (A) 200 μ M rotenone, (B) 25 mM MPP⁺ or (C) 10 mM paraquat and were scored daily for survival. Error bars indicate standard deviation. Data were analyzed using the Kaplan-Meier log-rank statistical test. *park*^{25/25} vs. *park*^{25/+} for all toxins tested, $p < 0.001$; *park*^{25/25} normal food vs. *park*^{25/25} food plus toxin for all toxins tested, $p < 0.01$.




Fig. 2. Normal counts of dopamine neurons in *parkin* mutant flies exposed to environmental toxins. (A) TH-positive neurons of DM/PPM and DL1/PPL1 clusters of the *Drosophila* brain revealed by TH-DAB immunohistochemistry in 5- μ m-thick paraffin sections of a wildtype fly brain. (B) TH-positive neurons labeled by fluorescent immunohistochemistry in a whole brain of an adult fly (whole-mount). DL1/PPL1, protocerebral posterior lateral TH neurons; DM/PPM, protocerebral posterior medial TH neurons. (C–D) Lack of dopamine neuron loss in flies exposed to MPP⁺, paraquat or rotenone. Dopamine neurons were stained with the anti-TH antibody and quantified by confocal microscopy in whole brains of flies of the various genotypes. Two- to 3-day-old flies were grown on food supplemented with 25 mM MPP⁺, 10 mM paraquat or 200 μ M rotenone. Neurons of at least seven brains per genotype and toxin were counted for the DL1/PPL1 cluster (C) and DM/PPM cluster (D). Error bars indicate standard deviation.




Fig. 3. Lifespan of parkin mutant flies grown on food supplemented with metal-chelating agents and redox-active metals. All experiments were done in triplicate with $n = 20$ flies per genotype and experiment. (A) Flies grown on food containing 0.1 mM of the iron chelator bathophenanthroline sulfonated sodium salt (BPS), 0.3 mM of the copper chelator bathocuproine disulfonate (BCS), or a combination of both metal-chelating agents. $park^{25/+}$ normal food vs. $park^{25/+}$ food plus BCS, $p < 0.01$; $park^{25/25}$ normal food vs. $park^{25/25}$ food plus BPS, $p < 0.01$; $park^{25/25}$ normal food vs. $park^{25/25}$ food plus BCS and BPS, $p < 0.001$. Survival of $park^{25/+}$ control flies was unaffected by the metal chelators up to 50 days of age (data not shown). (B) Qualitative and (C) quantitative gustatory assay, showing that flies consumed comparable amounts of normal and metal chelator-supplemented food (for details, see Materials and Methods). Twenty flies per genotype were analyzed. Pictures in panel B were taken at similar but not identical magnification to qualitatively show abdominal redness. (D) ROS-dependent DCF fluorescence in the heads and brains of $park^{25/25}$ and $park^{25/+}$ flies, showing that parkin ablation increases oxidative stress, which is reduced after metal sequestration by BPS and BCS. Pools of twenty brains per treatment and genotype were analyzed. *** $p < 0.001$ and ** $p < 0.01$ compared to normal food for the same genotype (black stars), or compared to the same treatment between genotypes (red stars). (E) Survival of $park^{25/25}$ and $park^{25/+}$ flies on food containing 500 μ M ammonium ferric citrate, $p < 0.01$. (F) Survival of $park^{25/25}$ and $park^{25/+}$ flies on food supplemented with 500 μ M Cu(II)-sulfate, p value not significant.




Fig. 4. Overexpression of *Ctrl1B* causes a rough eye phenotype in *park^{25/25}* but not *park^{25/+}* flies. Control *w; GMR-Ctrl1B/Cyo* transgenic flies grown on (A) regular food and (B) food supplemented with 20 μ M Cu(II)-sulfate . (C) *park^{25/25}* flies. (D) Rough eye phenotype in *w;GMR-Ctrl1B/Cyo; park^{25/25}* flies. (E) Normal eye phenotype in *w; GMR-Ctrl1B/Cyo; park^{25/+}* flies. Supplementation of food with (F) 20 μ M and (G) 50 μ M AgNO_3 did not cause a rough eye phenotype in *w;GMR-Ctrl1B/Cyo; park^{25/25}* flies, showing that the rough eye phenotype in *park^{25/25}* flies overexpressing *Ctrl1B* is copper-dependent (see main text).




Fig. 5. Effects of antioxidants and SOD1 overexpression on the lifespan of parkin mutant flies. (A) *park*^{25/25} flies grown on food containing 15 mM N-acetylcysteine (NAC), the metal-chelating agents BPS (0.1 mM) and BCS (0.3 mM), or both NAC and metal chelators display increased lifespan compared to *park*^{25/25} flies grown on regular food. Kaplan-Meier log-rank test, $p < 0.01$ for all treatments compared to normal food. (B) Ascorbic acid increased lifespan of *park*^{25/25} flies at 100 mM (Kaplan-Meier log-rank test, $p < 0.01$) but not 250 mM and 500 mM. (C) Overexpression of SOD1 using the *twi-Gal4* driver significantly increased the number of surviving *park*^{25/25} flies at 11 days and 17 days (paired Student's *t*-test, $p < 0.05$), at which time point the experiment was terminated because all *park*^{25/25} flies had died. Experiments were carried out in triplicate with 20 flies per experiment. Data are expressed as mean \pm standard deviation.




Fig. 6. Parkin-deficient flies are more vulnerable to hypoxia and heat shock. (A) *park^{25/+}* and *park^{25/25}* flies (six sets of 20 animals per genotype) were exposed for 7 min to 0.5% oxygen at room temperature (25 °C) in a closed hypoxia chamber, and the time to regain consciousness (ability to move) was recorded. Student's *t*-test, $p < 0.001$. (B) *park^{25/+}* and *park^{25/25}* flies ($n = 20$ per genotype) were exposed to 38 °C for either 30 min or 60 min and survival was recorded 24 h later. Experiments were done three times. Student's *t*-test, $p < 0.01$.

Molecular and Cellular Biology, 2011 May;31(10):2151-61.

***Parkin* mutant in the fly is largely rescued by metal-responsive transcription factor (MTF-1)**

Nidhi Saini¹, Oleg Georgiev¹, Walter Schaffner^{1*}

¹Institute of Molecular Life Sciences, University of Zürich

Abstract

The gene for Parkin, an E3 ubiquitin ligase, is mutated in some familial forms of Parkinson's disease, a severe neurodegenerative disorder. A homozygous mutant of the *Drosophila* ortholog of human *parkin* is viable but results in severe motoric impairment including an inability to fly, female and male sterility, and a decreased lifespan. Here we show that a double mutant of the genes for Parkin and the metal-responsive transcription factor MTF-1 is not viable. MTF-1, which is conserved from insects to mammals, is a key regulator of heavy metal homeostasis and detoxification and plays additional roles in other stress conditions, notably oxidative stress. In contrast to the synthetic lethality of the double mutant, elevated expression of MTF-1 dramatically ameliorates the *parkin* mutant phenotype, as evidenced by prolonged lifespan, motoric improvement including short flight episodes, and female fertility. At the cellular level, muscle and mitochondrial structures are substantially improved. A beneficial effect is also seen with a transgene encoding human MTF-1. We propose that Parkin and MTF-1 provide complementary functions in metal homeostasis, oxidative stress and other cellular stress responses.

***Parkin* mutant in the fly is largely rescued by metal-responsive
transcription factor (MTF-1)**

Nidhi Saini¹, Oleg Georgiev¹, Walter Schaffner^{1*}

¹Institute of Molecular Life Sciences
University of Zürich
Winterthurerstrasse 190
CH-8051, Switzerland

*Corresponding author: Walter Schaffner
Institute of Molecular Life Sciences
University of Zurich
Winterthurerstrasse 190
CH-8057 Zurich
Switzerland
phone: +41 44 635 31 50
fax: +41 44 635 68 11
Email: walter.schaffner@imls.uzh.ch

Running title: *Parkin* mutant largely rescued by MTF-1

word count for the introduction, results, and discussion sections: 3343

word count for the Materials and methods section: 2084

Abstract

The gene for Parkin, an E3 ubiquitin ligase, is mutated in some familial forms of Parkinson's disease, a severe neurodegenerative disorder. A homozygous mutant of the *Drosophila* ortholog of human *parkin* is viable but results in severe motoric impairment including an inability to fly, female and male sterility, and a decreased lifespan. Here we show that a double mutant of the genes for Parkin and the metal-responsive transcription factor MTF-1 is not viable. MTF-1, which is conserved from insects to mammals, is a key regulator of heavy metal homeostasis and detoxification and plays additional roles in other stress conditions, notably oxidative stress. In contrast to the synthetic lethality of the double mutant, elevated expression of MTF-1 dramatically ameliorates the *parkin* mutant phenotype, as evidenced by prolonged lifespan, motoric improvement including short flight episodes, and female fertility. At the cellular level, muscle and mitochondrial structures are substantially improved. A beneficial effect is also seen with a transgene encoding human MTF-1. We propose that Parkin and MTF-1 provide complementary functions in metal homeostasis, oxidative stress and other cellular stress responses.

Keywords: *Drosophila*/ MTF-1/ metal homeostasis/ *parkin*/ Parkinson's disease

Introduction

Parkinson's disease (hereafter referred to as PD) is the second most prevalent progressive neurodegenerative disorder and the most common age-related movement disorder (10, 13, 43, 59). Many molecular aspects of PD pathogenesis still need to be clarified. Extensive studies point to oxidative stress as a major contributor to the disease (28). Besides the gene for Parkin, an E3 ubiquitin ligase, four other genes: *PINK1*, *DJI*, *UCHL1* and α -synuclein have been implicated in rare, early-onset, familial forms of PD while *LRRK2* is predominantly responsible for late onset PD (20, 57, 70). Much effort has gone into the development of animal models of PD, including models in the fly *Drosophila melanogaster*. In our studies presented here, we use a strain in which the ortholog of the human *parkin* gene has been disrupted by insertion of a P-element transposon into the coding region (24, 48).

In mammals, the proteins PINK1 and Parkin cooperate to ensure proper quality control of mitochondria and Parkin is particularly important for autophagy of faulty mitochondria (reviewed in (8, 73)). In agreement with this notion, Parkin deficient flies suffer from mitochondrial malfunction (24, 45, 48), which distorts muscle structure and causes severe locomotor defects and an inability to fly (24, 48). Furthermore, both male and female *parkin* null mutant *Drosophila* are sterile (52), exhibit an increased sensitivity to multiple stresses, including oxidative stress, and have a reduced lifespan (23, 48).

Maintenance of metal homeostasis is an essential requirement for the proper functioning of all organisms. An adequate supply of essential trace metals, like copper and zinc, is important, whereas an excess can be highly toxic. Alterations in copper homeostasis due to mutations in copper transporters cause Wilson's and Menkes disease (11, 31, 65). An imbalance in trace metal levels has also been implicated in neurodegenerative disorders such as Parkinson's and Alzheimer's disease, as well as in senescence processes (42, 53, 54). To investigate the possible interplay of Parkin function with metal homeostasis, we modulated the concentration of the metal responsive transcription factor-1 (MTF-1) in *parkin* mutant *Drosophila*. MTF-1 is conserved in evolution and its homologs have been characterized in humans (7, 36, 44), mice (25, 51, 69), fish (3, 9) and *Drosophila* (17, 60, 75). MTF-1, also referred to

as metal response element binding transcription factor 1, is a zinc-finger protein that regulates transcription of its target genes by binding to DNA sequence motifs known as metal response elements (MREs), which are typically located proximal to the transcription start (12, 27, 32, 41, 64, 68). The majority of MTF-1 preferentially localizes to the cytoplasm in quiescent, non-stressed cells but translocates to the nucleus upon heavy metal load and a number of other stressful conditions (34, 58, 62).

Apart from counteracting the effects of heavy metal load, MTF-1 also induces transcription of metallothionein genes in response to oxidative stress and infection (2, 21, 22). Metallothioneins (MTs), are small, cysteine-rich, metal-binding proteins with a major role in metal homeostasis and detoxification (30, 46). MTs occur in all eukaryotes, as well as in some prokaryotes. Heavy metals like zinc, copper and cadmium are complexed by the cysteine sulfhydryl groups, which can also exert antioxidant function (2, 29). In this context it is noteworthy that in a mouse PD model, dopaminergic (DA) neurons of a MT-knockout mutant are more vulnerable to L-DOPA toxicity than neurons from mice with wild type MT (15, 39). This suggests that MTs play a protective role against DA-quinone induced neurotoxicity. There are more than ten functional metallothionein genes in humans, four in the mouse (50, 71) and at least four in *Drosophila*, termed *MtnA-MtnD* (17, 40). The *Drosophila* MTs are involved to different degrees in the defense against heavy metal stress. *MtnA* is the most important under copper load, while *MtnB* preferentially binds cadmium and protects against cadmium intoxication. *MtnC* and *MtnD*, despite sharing 67% amino acid identity with *MtnB*, have only a minor role in protection against heavy metals, at least when *MtnA* and *MtnB* are present (16, 18). The *MtnA* and *MtnB* genes (also referred to as *Mtn* and *Mto*, respectively) are differentially regulated during development (61). In addition to metallothioneins, MTF-1 also regulates, in *Drosophila*, the expression of ferritins, the copper importer Ctr1B, the zinc exporter ZnT35C, glutathione S-transferase and an ABC transporter (60, 63, 74).

MTF-1 proteins of human and *Drosophila* are highly similar in their DNA-binding zinc finger region but quite divergent outside of it. Nevertheless, they can largely complement each other in the protection against metal stress (6, 75). A major difference between mammals and *Drosophila* is that metallothionein genes in

mammals are mainly induced by zinc and cadmium whereas in *Drosophila* they are best induced by copper and cadmium (18, 75, 76). Moreover, disruption of the *MTF-1* gene in the mouse results in embryonic lethality (25), which is not the case for fly mutants, which are viable and fertile. However, the fly mutants do display sensitivity to cadmium, zinc and copper load as manifested by a reduced lifespan on heavy-metal supplemented food, and also cannot tolerate copper starvation (17, 60). The *Drosophila* allele *MTF-1*^{140-1R} carrying a 4.1 kb deletion of the coding region has the strongest phenotype and is considered a null mutation (17). We have used this allele for experiments requiring a *MTF-1* loss of function.

A link between cellular heavy metal handling and the *parkin* mutant phenotype was suggested by our previous finding of a partial rescue of *parkin* mutant flies upon zinc supplementation or chelation of redox-active metals (55, 56). In our present study we set out to investigate the interaction between metal homeostasis and Parkin function, or more specifically, MTF-1 and Parkin. We found that *parkin* mutants combined with a knockout of *MTF-1* are not viable, a genetic constellation termed synthetic lethality. *Parkin* mutant *Drosophila* suffer from oxidative stress as a result of heightened ROS production. A strong ubiquitous MTF-1 expression dramatically ameliorates the *parkin* mutant phenotype: our results show that MTF-1 decreases oxidative stress, normalizes concentration of essential trace metals, increases the frequency of development to adulthood, restores female fertility, improves muscle/mitochondrial morphology, locomotion and considerably extends the lifespan of *parkin* mutant flies.

Materials and Methods

Fly food and maintenance

One liter of standard fly food was composed of 55 g cornmeal, 10 g wheat flour, 100 g yeast, 75 g glucose, 8 g agar and 15 ml anti-fungal agent Nipagin (15% in ethanol). For the experiments of survival, development, eclosure frequency, ROS measurements, real-time PCR and TEM muscle analysis, several conditions were tested, namely, normal food (NF), NF supplemented with zinc chloride (4 mM) or N-acetylcysteine (15 mM) or bathocuproine disulfonate (BCS, copper chelator) (0.3

mM), and bathophenanthroline sulfonated sodium salt (BPS, iron (copper) chelator) (0.1 mM). All flies were maintained at 25°C on a 12:12 hours light-dark cycle.

Construction of transgenic flies and fly stocks

UAS-MTF-1 flies were generated using the full-length MTF-1 cDNA cloned into the pUAST vector. *Tub-MTF-1* constructs were made by cloning the MTF-1 cDNA under the control of the constitutive α -tubulin promoter. Both the constructs were injected into the *w¹¹¹⁸* fly strain along with p(Δ 2-3) helper plasmid and transformants were selected based on the eye color (red/orange). The *MTF-1^{I40-1R}* null allele, generated by homologous recombination, was characterized previously (17). The *UAS-MTF-1*, *MT* (*tub-MtnA*), *MTF-1^{I40-1R}* strains were generated by recombination in our laboratory. The *w⁺; +; tub-MtnA/TM6B* (*tub-MtnA*), *w⁺; tub-MTF-1/Cyo⁺* (*tub-MTF-1*) and *w⁺; Actin-Gal4; UAS-MTF-1/TM6B* (*Actin-Gal4; UAS-MTF-1*) flies were combined with *w⁺; +; park^{25/25}* (*park^{25/25}*) by recombination. Unless specified otherwise, *w⁺; +; park²⁵/TM6B, w⁺* flies (heterozygous *parkin* mutants hereafter referred to as *park^{25/+}*) were used as controls for the *w⁺; +; park^{25/25}* flies (homozygous *parkin* mutants hereafter referred to as *park^{25/25}*). *park^{25/25}* is a null mutation of the *Drosophila parkin* gene (24). The TM6B balancer in the control flies was confirmed to have no effect on any of the experiments performed (by removing it).

Lifespan determinations, eclosure frequency and fertility assays

For lifespan experiments, 1-2 day old flies (20 per vial) maintained at 25°C on a 12:12 hours light-dark cycle were examined for each genotype at least in triplicate. Surviving flies were transferred to fresh food vials every 2 days and counted daily. In the experiment with *w⁺; Actin-Gal4; UAS-MTF-1, park²⁵/TM6B* flies, the survivors in 23 parallel independent sets were counted at regular time intervals. In each lifespan assay testing different conditions, the controls of *park^{25/+}* and *park^{25/25}* flies raised on NF were the same. The variations in the average lifespan of control flies in different experiments can be attributed to subtle experimental variations. The metal chelator concentrations selected for ROS determination had no significant effect on feeding behavior of the flies (55). Fertility was assayed by placing single *parkin* mutant males with 3-4 virgin *yw* females and by placing single virgin *parkin* mutant females with two *yw* males. Vials were checked 3-7 days later for the presence of larvae and eclosing adults. For the analysis of eclosure frequency, the number of days allowed

for egg laying and the parent population was the same in all vials of normal food (NF) or zinc (Zn)-supplemented food and progeny flies were counted at the same time.

Behavioral assay (climbing performance and locomotion ability)

The Climbing assay was performed as described (47). Flies of each genotype (*park*^{25/+}, *park*^{25/25}, *tub-MTF-1, park*^{25/+}, *tub-MTF-1, park*^{25/25}) were anesthetized with CO₂ and individually counted and placed in food vials 24 hours before the assays were performed to enable a full recovery from the effects of CO₂. Ten flies were placed in an empty 110-by 27-mm vial; a horizontal line was drawn 100 mm above the bottom of the vial and another identical vial was used as a cover to provide more mobile space. After the flies had acclimated for 10 min at room temperature, each genotype was assayed in triplicate for five trials per set per genotype. The procedure involved gently tapping the flies (on a soft surface) down to the bottom of the vial. The flies were given 30 seconds to climb the vial and the number of flies which crossed the 100 mm mark each time were recorded. These values were then averaged and a group mean and standard error was calculated. The mean values of various fly groups were statistically compared using an unpaired Student t-test. The study was repeatedly performed with the same group of flies on every alternate day up to 10 days in an isolation room at 25°C, 60-70% humidity under standard lighting conditions. Preliminary studies indicated no significant difference in the outcome of climbing assays performed in normal light or red light conditions.

Fluorescent protein (EYFP) reporter

The *Drosophila* metallothionein *MtnA* promoter (-446 to +74) was cloned from genomic DNA using the primer pair 5'-CGG GAT CCA GGT ATG GGC TAT TTA GGC C-3' and 5'-GGG ATG GCC CCA AAG GAT CTG-3' in a pCasper4-derived vector carrying EYFP-coding sequence and the SV40 polyA site. Details were reported previously (6, 18). Transgenics of *MtnA-EYFP* combined with *parkin* heterozygous or homozygous knockouts were made. Both fly types were then frozen at the same age and photographs of EYFP expression were taken with a Leica MZ FLIII fluorescence stereomicroscope and a Nikon Coolpix 950 digital camera (Leica, Heidelberg, Germany) at an exposure of 730 ms.

RNA isolation and real-time analysis

Total RNA was purified from adult *Drosophila* tissue using the Nucleospin RNA II protocol (Macherey-Nagel) and eluted in 60 µl of RNase-free water. cDNA was prepared using the Transcriptor High Fidelity cDNA Synthesis kit from Roche. The cDNA obtained was further purified using the AM 1906 Ambion DNA free kit and used for analysis by real-time PCR on the Tecan Genesis 200/8 robot using the Eurogentec Mesa Green qPCR Mastermix Plus for SYBR assays. The qPCR run was performed on an Applied Biosystem machine (ABI Prism SDS 7900 HT) in a 384 well format with a reaction volume of 10 µl. $\Delta\Delta\text{Ct}$ values were calculated by subtracting the ΔCt calibrator from the ΔCt sample, ΔCt values were calculated by subtracting ΔCt endogenous control from ΔCt target gene/calibrator. The normalization strategy used has been described in Vandesompele *et al* (67). All the fold-change values are normalized to respective *park*^{25/+} values on normal food (NF). The housekeeping genes used were *actin5c*, *TBP* and *GAPDH*. Two sets of primer sequences were used for each of the transcripts quantified: for *parkin*, the first primer set was 5'-AAG ATC ATA TTT GCC GGT AAG GAA-3' and 5'-CGC TTT GCT GAC CCA AGT C-3' which amplify a 73 bp fragment only from the *parkin* heterozygous control flies and the second set was 5'-CAA AGC CCT GTC CAA AAT GC-3' and 5'-GCG CGT GTG CAG ACC AT-3'; for *MTF-1*, the first primer set was 5'-TGT CCG GCT GCG ATA AGG-3' and 5'-GCC ATT GTG CAG ACG AAG GT-3' which amplify a 68 bp fragment from wild-type MTF-1 containing flies and the second set was GCA TTC AAC ACG CGC TAC A-3' and 5'-ACA GTT GAA CGT CTC GCC ATT-3'; for *MtnB*, the first primer set was 5'-TTG CAA GGG TTG TGG AAC AAA-3' and 5'-TGC AGG CGC AGT TGT CC-3' which amplify a 65 bp fragment and the second set was 5'-AAG TCG AGA AAT AGA TAC ATA CAA GAT GGT-3' and 5'-CGC ACT TTT GGG CCG AG-3'; for *foi*, the first primer set was 5'-GTG GCT GCG GGT CTG TTC-3' and 5'-TTT GTG CGA GGC CGA GAT-3' which amplify a 69 bp fragment and the second set was 5'-TGG CGA TGC CCT ACT TCA C-3' and 5'-TGA TCA TCC CCC GCT CAT-3'.

Detection of ROS levels

Fresh dichlorofluorescein diacetate (2,7-DCFH-DA) from Invitrogen – Molecular Probes Cat # C369 was mixed with DMSO to make a 1 mM stock solution (=2,7-

DCFH-DA/DMSO). A 40 μ M working solution was prepared in HEPES buffer (30 mM). 50 heads of frozen adult flies of a specific genotype (same age) were removed and collected in an Eppendorf tube. This was done in triplicate for each condition. Each sample was then homogenized using cold protein homogenization buffer (1:3 w/v of 0.32 mM sucrose, 20 mM HEPES, 1 mM $MgCl_2$, 0.5 mM PMSF protease inhibitor at pH 7.4) and centrifuged for 20 minutes at 4°C and 20,000 g. Protein content of the supernatant was determined using the Bio Rad diagnostics kit and a final concentration of 0.4 mg/ml was used as the standard for the ROS assay of each genotype tested. The fluorescence intensity (emission acquisition) was monitored for 45 minutes after the sample (20 μ l) was kept in a cuvette (1.5 x 1.5 nm) in the fluorimeter immediately after the addition of the DCFH-DA dye (2 μ l) to prevent loss of signal due to fading of fluorescence. Excitation of dye was at 485 nm and emission at 520 nm. The curve area of fluorescence intensity which was recorded every 5 minutes (for 45 minutes) in the range of 500-600 nm was integrated and the total area was used for comparison, with the final result obtained in counts per second. The fluorimeter was standardized using 0.05% H_2O_2 , a positive ROS generating species.

Dissection of ovaries

Ovaries from female parents were dissected in Grace's insect medium (1X), GIBCO, Invitrogen at room temperature (RT). The dissected ovaries were immediately fixed in 4% paraformaldehyde, 0.2% Triton-X-100 dissolved in Grace's medium for 20 minutes without shaking. The fixative was washed three times with phosphate buffer saline (PBS)+0.5% Triton (PBST) for 10 minutes each. The samples were then labeled for one hour with fluorescently labeled phalloidin (Phalloidin-alexa 568, molecular probes), diluted 1:200 in PBST at RT. This was followed by three washes with PBST for 10 minutes each and a second labeling with Toto (nuclear stain) diluted 1:1000 in PBST at RT. The samples were then washed three times with PBST for 10 minutes each, followed by two washes with PBS for 10 minutes each and then embedded in vectashield (mounting medium) overnight at -20°C. Ovarioles were dissected from the ovaries and mounted on glass slides. Pictures were taken using the confocal at 20X magnification.

Quantification of metal content

Female flies (*park*^{25/+}, *tub-MTF-1*, *park*^{25/+} and *yw* and *MTF-1* knockout controls) were allowed to lay eggs on normal food or metal-supplemented food (100 μ M cadmium sulphate/ 500 μ M copper sulphate/ 500 μ M ammonium ferric citrate/ 4 mM zinc chloride) for four days and removed afterwards. The resulting progeny were collected at regular intervals and frozen. This procedure was repeated until the required number of 50 flies was obtained in triplicate for each genotype. Each sample set of frozen flies was then subjected to homogenization using cold protein homogenization buffer (0.32 mM sucrose, 20 mM HEPES, 1 mM MgCl₂, 0.5 mM PMSF protease inhibitor at pH 7.4) and the samples were normalized for protein content. A final concentration of 1 mg/ml was prepared by diluting the samples in 0.2M HNO₃ to obtain a total assay volume of 1 ml. A highly sensitivity flame atomic absorption spectrophotometer (FAAS; GTA-120/PSD-120, Varian Australia Pty Ltd, Mulgrave, VIC, Australia) was used to detect the metal content in each genotype assessed. Cd and Zn concentrations were recorded by the same flame. Likewise Cu and Fe concentrations were measured together.

Muscle section and TEM

Dissected thoraces of two-day old anesthetized adult flies (*park*^{25/+}, *park*^{25/25} and *Act-Gal4; UAS-MTF-1, park*^{25/25}) were kept in ice-cold fixative (2.5% glutaraldehyde/0.1 M sodium cacodylate buffer adjusted to 328 mOsm/l with sucrose, pH 7.4) for 4 hours at 4°C. Postfixation was performed with 1% OsO₄ and 0.1 M sodium cacodylate, pH 7.4 for 2 hours at 4°C and the sample was washed overnight with 0.1 M sodium cacodylate, pH 7.4 before going through a series of progressive dehydration steps in a graded ethanol series of 70/80/96% alcohol for 10 minutes each, followed by three 10 min washes in 100% alcohol and a final 20 min wash in propylene oxide. The sample was treated with a 1:1 propylene oxide: Epon (Epon 812) mix for 2 hours and embedded in Epon overnight at 70°C. Blocks of thoraces were trimmed and semi-thin sections of dorsal longitudinal muscles were stained with Toluidine Blue dye, which labels nucleic acids, hence staining both nuclei and cytoplasm (5). Ultrathin sections of 70 nm thickness were made with an ultramicrotome for the selected sections. For TEM, sections were contrast-stained on the grid first with 2% uranyl acetate, then with 2.5% lead citrate (Reynold's), each of

them for twenty minutes at room temperature. The sections were inspected with a Philips CM100 TEM with GATAN Orius Camera.

Statistical Analysis

JMP software (SAS Institute) was used for statistical evaluations. Lifespan (survival) assays were analyzed with the Kaplan–Meier log-rank statistical test. Brain ROS levels and qPCR results were compared by one-way ANOVA. Results are expressed as mean \pm standard deviation.

Results

Synthetic lethality of combined *parkin* and *MTF-1* mutants

The metal-responsive transcription factor (MTF-1) is a key regulator of metal homeostasis in *Drosophila* (17, 60, 75). Based on our findings of a connection between the *parkin* knockout and trace metal status (55), we tested the effect of an *MTF-1* knockout in a *parkin* mutant background. The result was clear cut, in that no surviving double mutant was ever observed in 50 independent crosses. This “synthetic lethality,” observed at the pupal stage, was rescued by a cDNA transgene of *MTF-1* driven by the constitutive tubulin enhancer/promoter. In a genetic cross between parents heterozygous for the *parkin* and *MTF-1* recombined deletions (Figure 1), 69 out of a total of 217 progeny (32%) were *parkin* and *MTF-1* homozygous knockouts also expressing an *MTF-1* transgene (statistical expectation, 40%). This rescue by an MTF-1 cDNA transgene confirms the absence of any secondary hits as the cause of the observed lethality. Furthermore, it also excludes the possibility that an intronic open reading frame located within the *MTF-1* gene (D. Steiger, K. Steiner and WS, unpublished) is responsible for the effect and not MTF-1 itself. To find out what condition could overcome the pupal lethality, we maintained the heterozygous *parkin*, *MTF-1* parent flies on N-acetylcysteine (NAC), which is a precursor to glutathione, an established antioxidant. Since either condition, lack of a functional *parkin* gene or of an *MTF-1* gene, increases reactive oxygen species (ROS) (see below), we reasoned that keeping a lower ROS level by other means might also overcome the synthetic lethality of the double mutants. After testing different concentrations of NAC an optimal supplement of 15 mM was chosen for the main experiment. Indeed, raising

the progeny under this condition resulted in a substantial rescue of the synthetic lethality: of the total progeny, 19% was homozygous *parkin* and *MTF-1* double mutant (statistical expectation, 25%) (Table 1A). Other antioxidants tested included ascorbate, zinc, and metal chelators of copper and iron (BCS and BPS, respectively) which we had shown before to positively influence the *parkin* mutant phenotype (55, 56). However, none of these was able to rescue synthetic lethality.

MTF-1 overexpression rescues the lifespan and the low eclosing frequency of *parkin* mutants

We examined the lifespan of *Drosophila* *MTF-1*-overexpressing transgenic lines (*tub-MTF-1* and *Act-Gal4; UAS-MTF-1*) in a *park*^{25/25} background. These experiments illustrated that an elevated expression of MTF-1 from the ubiquitously active *tubulin* enhancer/promoter prolonged the lifespan of *parkin* mutants significantly, from a median of 7 days for the mutants alone to 21 days (Figure 2A). In this experiment, the maximal lifespan was extended from 12 to 41 days by the *MTF-1* transgene (Figure 2A). The stronger combination with *actin-Gal4* driving *UAS-MTF-1* revealed a similar effect: in an experiment with 23 independent replicas, 10% of the mutant animals were still alive at day 34 (Figure 2B). Overexpression of MTF-1 in control and wild type flies (*park*^{25/+} or *park*^{+/+}) did not increase their normal lifespan (data not shown). Also, in an independent study, MTF-1 overexpression did not extend lifespan of flies kept on standard food (4). Elevated MTF-1 expression not only prolonged the lifespan of *parkin* mutant adult flies but also enhanced survival during development. In a genetic cross involving *parkin* heterozygous parents, only 2.5% of the eclosing progeny flies were homozygous *parkin* mutants (statistical expectation, 25%). In comparison, the same cross but also expressing an MTF-1 transgene resulted in a 15% eclosure frequency (Table IB).

Elevated expression of MTF-1 rescues female fertility and fecundity of *parkin* mutant flies

Strikingly, female fertility and fecundity was completely rescued by *MTF-1*. When crossed with wild type males, *parkin* mutant females with the tubulin-driven *MTF-1* transgene produced the same number of progeny as a cross of wild type males and

females. *Drosophila* gonad formation requires a complex morphogenetic process (35, 37). As in the majority of metazoans, *Drosophila* oogenesis occurs within the ovarian follicles in which germline cells develop in close proximity to specialized somatic cells (Figure 3A-A''). Parkin mutant females lack the proper spatio-temporal development in the germarium and thus have stunted ovaries with few mature oocytes, which fail to get fertilized (Figure 3B-B''). The restoration of female fertility by strong MTF-1 expression was also evident at the morphological level: dissected ovaries showed a normalized structure with follicles formed in the germarium and mature stages in the posterior regions of the ovariole, with several oocytes ready for fertilization (Figure 3C-C''). In contrast, the sterility phenotype of *parkin* mutant males which is due to defective spermatogenesis at the individualization step (24) was not rescued. This suggests that Parkin is particularly important for male fertility.

Improved locomotion and rescued mitochondrial/myofibrillar morphology of *parkin* mutants with enhanced MTF-1 expression

Strong MTF-1 expression dramatically improved the climbing ability of *parkin* mutant flies (Figure 4). What is more, they generally moved around fast, responded by running away when physically perturbed, jumped and occasionally displayed short flight episodes (data not shown). To investigate the effect of elevated MTF-1 expression on muscle morphology we examined the ultrastructure of the indirect flight muscle in heterozygous control flies, *parkin* mutants and the MTF-1 transgenic flies. Cross-thoracic sections of control adults analyzed by transmission electron microscopy (TEM) revealed well organized muscle fibres in parallel stripes with a regular M- and Z-line banding pattern and darkly stained, electron-dense mitochondria with regularly packed cristae (Figure 5A-C). In contrast, age-matched *parkin* mutants had abnormal muscle structure with large vacuoles, a reduced muscle content with mostly irregular arrangement and enlarged mitochondria with disintegrated cristae (Figure 5D-F). The MTF-1 overexpressing *parkin* mutant flies displayed a clear rescue effect in that muscle fibre structure was more regular with less prominent vacuoles; moreover, mitochondria had more densely packed cristae with considerably lesser signs of disintegration in comparison to *parkin* mutants (Figure 5G-I).

MTF-1 dependent expression of metallothioneins is higher in *parkin* mutant *Drosophila*

Owing to their high cysteine content, metallothioneins can act as antioxidants, in addition to their obvious role as metal chelators (33, 38). Basal and induced levels of metallothionein expression depend on the transcription factor MTF-1 (17, 51, 74). To find out whether metallothionein genes could be induced by MTF-1 in a *parkin* mutant background, we used a reporter line in which a yellow fluorescent protein (EYFP) is driven by the promoter of *MtnA*, the most highly expressed *Drosophila* metallothionein gene. Compared to heterozygous controls, even in the absence of any heavy metal load the basal expression of this reporter was increased, likely due to elevated oxidative stress in the *parkin* mutant flies (Figure 6C and D). This metal-independent upregulation of the *MtnA* promoter, was strictly dependent on MTF-1, since no trace of fluorescence could be detected in flies lacking MTF-1 (data not shown). RT-PCR of *parkin* mutants also revealed elevated transcript levels of *MTF-1* (2-fold) (Figure 7A) and the embryo-enriched metallothionein *MtnB* (6-fold) (Figure 7B) compared to *park*^{25/+} flies. MTF-1 overexpression in the *park*^{25/25} flies was achieved from the tubulin promoter or indirectly with the stronger UAS-Act-Gal4 system, both induced a more than 200-fold increase in *MtnB* transcripts (Figure 7B). Conversely, the level of *parkin* transcripts was increased in the *MTF-1* knockout flies (Figure 7C). This pattern of regulation can be explained by a partial redundancy of Parkin and MTF-1 where one is upregulated to compensate for the loss of the other. The synthetic lethality of a combined knockout of *parkin* and *MTF-1* genes mentioned above is in line with this hypothesis. Another experiment revealed that the transcript levels of the zinc importer *foi* (37) in *parkin* mutants were enhanced 3-fold by elevated MTF-1 expression, which may contribute to the normalized structure of ovaries and rescued fertility of female *parkin* mutants (Figure 7D). This hypothesis is further supported by the increased level of zinc ions found in tub-MTF-1; *park*^{25/25} flies (discussed below, Figure 8B).

Elevated MTF-1 expression confers resistance to oxidative stress and restores metal homeostasis

Parkin mutant flies display high levels of ROS indicative of intrinsic oxidative stress (23, 72). In agreement with MTF-1 having an antioxidant function, we observed a substantial decrease of ROS levels in the heads of *parkin* mutants expressing an MTF-1 transgene: ROS dropped to approximately half of the levels in *park*^{25/25} flies (Figure 8A). Previously we had observed that limiting the availability of redox-active metals, achieved by supplementing the food with chelators for copper and iron (BCS and BPS, respectively), also increased the lifespan of *parkin* mutant flies (55). Furthermore, the *w;tub-MTF1;park*^{25/25} flies raised on the metal-chelator-supplemented food displayed a somewhat lower ROS level (Figure 8A). ROS levels were not significantly changed in heterozygous control flies, either upon ubiquitous MTF-1 overexpression or following dietary intake of metal-chelators. The *MTF-1* knockout flies showed the highest levels of ROS, probably due to the reduced expression of MTF-1-dependent antioxidant genes such as metallothioneins (Figure 8A). Metals like zinc and the redox active copper and iron are required in trace amounts for several structural and biological processes in organisms (66). *Park*^{25/25} flies display not only reduced basal levels of zinc (see also (56)) but also of copper and iron in comparison to control flies (Figure 8B). Tubulin-driven MTF-1 expression in *park*^{25/25} restores the basal level of these metals (Figure 8B). Concentrations of cadmium are generally low since it is a non-essential, toxic metal. Upon supplementing fly food with metals, their levels became quite similar in all three genotypes tested (Figure 8C). This is particularly important in the case of zinc supplementation, which we had previously shown to improve the condition of *parkin* mutant *Drosophila* (56).

Human MTF-1 or elevated metallothionein expression also improve condition of *parkin* mutants

Human MTF-1 has been expressed in *Drosophila* and shown to largely, but not completely, rescue the metal sensitivity of *Drosophila* lacking its endogenous MTF-1 (6). We therefore tested the effect of *actin-Gal4*-driven hMTF-1 expression in a *parkin* mutant background. Indeed the lifespan of *parkin* mutants was increased from

a median of 7 days to 19 days, which is close to the 21 days obtained with elevated *Drosophila*-MTF-1 expression (Figure 9). Other rescue effects paralleled those observed with elevated *Drosophila* MTF-1 expression (see above) but were less pronounced (data not shown).

Metallothionein genes are the major targets of MTF-1; in *Drosophila*, metallothionein A (*MtnA*) shows the strongest expression. Thus, we also tested if overexpression of *MtnA* in a *parkin* mutant background exerted a similar beneficial effect as *MTF-1* overexpression. To this end we crossed *inter se* three independent lines with tubulin-driven *MtnA* overexpression in a *parkin* heterozygous background to raise *parkin* homozygous knockouts. The median lifespan of the *parkin* mutants was extended up to 17 days (Figure 9) but other rescue effects associated with elevated MTF-1 expression (discussed in results) were not observed.

Discussion

Here we show that metal responsive transcription factor (MTF-1) plays a crucial role in modulating the severity of a Parkin loss-of-function phenotype. On the one hand, the combined loss of Parkin and MTF-1 is not viable, i.e., displays synthetic lethality. On the other hand, elevated expression of MTF-1 dramatically improves the condition of *parkin* mutant flies: there is an overall extension of life span, females regain normal fertility, and the motoric abilities of flies improve to the point that they can walk fast and even display short episodes of flight. The latter is noteworthy since flight muscles have a high energy consumption that depends on robust mitochondrial function. At the histological level, the degenerated mitochondria characteristically seen in *parkin* mutants are rescued to a more regular, electron-dense structure following MTF-1 overexpression. Additionally, flight muscles have an improved myofibrillar arrangement. In earlier studies, mitochondrial and muscle degeneration observed in *parkin* mutant flies were proposed to be a result of excessive oxidative stress (24, 48). Mitochondrial malfunction can indeed result in an increased susceptibility to oxygen radical damage, and mitochondria-associated increase in ROS production has been implicated in Parkinson's disease (19, 26, 73). In line with this concept, the major target genes of MTF-1 are metallothioneins (MTs), which encode small, cysteine-rich proteins that can scavenge heavy metals, notably the

redox-active copper, and ROS. The elevated basal level of *MTF-1* and MT transcripts in *parkin* mutant flies can thus be seen as a compensatory attempt to counteract enhanced ROS levels (24). In accordance with such a scenario is the rescue effect seen upon food supplementation with the antioxidant N-acetylcysteine, which however falls short of the dramatic effect seen with elevated MTF-1 expression. Besides metallothioneins and a number of other stress-associated genes, *ferritin* genes are a major target of MTF-1 in *Drosophila* (74). Ferritins are well-characterized iron binding proteins which keep iron in a soluble and non-toxic form in the cell. Thus ROS production by redox-active iron might be lowered via upregulation of *ferritin* levels. The multiple targets of MTF-1 help to explain why overexpression of metallothionein alone was less effective than MTF-1 overexpression in improving the condition of *parkin* mutants. In a separate series of experiments we found that strong expression of an ortholog, the human MTF-1, also increased lifespan, rescued female fertility and improved locomotion ability, although the effects were less pronounced than with *Drosophila*'s own MTF-1, presumably due to the evolutionary distance between mammals and insects. This is in line with previous findings from our laboratory that mammalian and *Drosophila* *MTF-1* transgenes are able to largely, but not completely, compensate for each other's absence (6).

Although we observed remarkable improvements following elevated MTF-1 expression, a complete rescue of *parkin* mutants, including male fertility and sustained flight ability, was only observed with a *parkin* transgene. Furthermore, it has to be pointed out that although *MTF-1* knockout flies display high ROS levels, they show no signs of Parkinson's disease-like symptoms. Together these findings are consistent with the idea that oxidative stress is an important but not the sole culprit in PD etiology (1, 14, 49). Nevertheless, the dramatic effect of MTF-1 on a *parkin* loss-of-function mutation underscores the importance of this transcriptional regulator in cellular stress response.

Acknowledgments

We are indebted to Drs. Jessica Greene and Leo Pallanck (University of Washington) for providing the *park*^{25/25} flies, to Gery Barmettler and Theresa Bruggmann (Center for Microscopy and Image analysis) for TEM analysis of muscle sections, to Martin

Moser for real-time analysis, Dr. Eva Freisinger and Tamara Huber (Inorganic Chemistry, UZH) for FAAS experiments, Ivan Ostojic (FMI Basel) for statistical analyses, and to Dr. Ben Schuler (Biochemistry, UZH) for providing the fluorimeter for ROS analysis. We also thank Till Strassen for maintenance of fly stocks and Dr. George Hausmann for critical reading of the manuscript. This work was supported by the Swiss National Science Foundation grant 31003A-113993 and the University of Zürich.

Conflict of interest

The authors declare that they have no conflict of interest.

Figure Legends

Figure 1. Scheme of genetic cross to obtain strong MTF-1 expression in double mutant background. An MTF-1 cDNA transgene driven by the *tubulin* promoter was combined with homozygous null mutations of both *MTF-1* (*MTF-1⁻*) (17) and *parkin* (*park²⁵*) (24); progeny in red. The *Sp* marker on the second chromosome gives a uniform-length side-bristle phenotype and the *CyO* second chromosome balancer results in curly wings. *TM3* and *TM6B* are third chromosome balancers which display a serrated wing phenotype and a dense side bristle phenotype, respectively. Balancers are lethal in homozygous form; progeny marked with a cross indicate lethality of that particular genotype. Chromosome 1 is wild type.

Figure 2. Enhanced lifespan of *parkin* mutants expressing an MTF-1 transgene. (A) A cDNA transgene of *Drosophila*-MTF-1 driven by the ubiquitously active *tubulin* enhancer/promoter prolongs the lifespan of *parkin* mutants (*park^{25/25}*) up to 41 days. For survival of *park^{25/25}* vs. *tub-MTF-1*, *park^{25/25}* flies, $p < 0.001$. (B) Strong expression of a UAS-MTF-1 transgene driven by actin-Gal4 in a *parkin* mutant background also extends lifespan, as tested in multiple sets of *Act-Gal4; UAS-MTF-1*, *park^{25/25}* flies. In both (A) and (B), heterozygous *park^{25/+}* flies with MTF-1 overexpression had a similar lifespan as *park^{25/+}* controls (data not shown).

Figure 3. Elevated MTF-1 expression restores ovary structures and restores

fertility of *parkin* mutant females. (A-A'') Normal ovariole structures of fertile control female flies (*park*^{25/+}). (B-B'') Infertile female *parkin* mutants (*park*^{25/25}) have a distorted ovary structure with very few mature eggs. (C-C'') Upon expression of a tubulin-driven *MTF-1* transgene ovary structures are normalised, resulting in normal fertility. a, anterior with germarium; p, posterior with vitellarium; broken arrows indicate mature eggs; full arrows developing stages of eggs. Phalloidin (red) stains tubulin structures and Toto (blue) stains the nuclei.

Figure 4. MTF-1 transgene expression restores the climbing ability of *parkin* mutants. Tubulin enhancer/promoter-driven MTF-1 expression largely rescues the locomotion ability of *park*^{25/25} flies. In control flies, MTF-1 overexpression does not further improve climbing ability. Data shown represents mean value \pm standard deviation of each group tested every other day up to 8 days. Asterisks show highly significant differences between *parkin* control (*park*^{25/+}) and *parkin* mutant (*park*^{25/25}) flies on each day of the assay ($p < 0.0001$).

Figure 5. Strong MTF-1 expression improves muscle and mitochondrial morphology of *parkin* mutants. (A-C) Transverse sections of indirect flight muscles (IFMs) show well preserved muscle in *park*^{25/+} heterozygous controls with a regular myofibril arrangement (white arrows) and many electron-dense mitochondria (broken red arrows). (D-F) *park*^{25/25} adult IFMs show an irregular myofibrillar arrangement with diffuse Z-lines and M-bands and numerous vacuoles. Mitochondria are swollen with fragmented cristae (red arrows). (G-I) Myofibril and mitochondrial integrity of *parkin* mutants is restored by MTF-1 overexpression. Mitochondria are more dense (broken white arrows) and muscle structure is more regular (broken green arrows), although occasional vacuoles are observed (green arrows). The scale bar is for top and middle panels while the bottom panel is at a higher magnification, shown for clarity.

Figure 6. MTF-1 activity is upregulated in a *parkin* deficient background. Top: Transgenic *MtnA-EYFP* reporter gene (6). Bottom: (A-B) *MtnA-EYFP*, *park*^{25/+} and (C-D) *MtnA-EYFP*, *park*^{25/25}. Pictures of 1-2 day old adult flies were taken at 730 ms exposure with a Leica fluorescence microscope. 'a' and 'p' show anterior and posterior ends, respectively.

Figure 7. Increased *MTF-1* and metallothionein (*MtnB*) transcript levels in *parkin* mutants that also express an *MTF-1* transgene. Real-time transcript-levels of (A) *MTF-1*, (B) *MtnB*, (C) *parkin* and (D) *foi* in *park*^{25/+} and *park*^{25/25} and *w; tub-MTF-1*, *park*^{25/25} and *Act-Gal4; UAS-MTF-1*; *park*^{25/+} and *Act-Gal4; UAS-MTF-1*; *park*^{25/25} flies with *MTF-1* knockout flies as an appropriate control. All flies were raised on normal food (NF). *MTF-1* null flies show no *MtnB* and *parkin* null flies show no *parkin* transcripts. (B) *MTF-1* overexpression from the tubulin enhancer/promoter or via the UAS-Act-Gal4 system induced a 234-fold and 270-fold increase in *MtnB* transcripts, respectively. *** indicates significant difference between *parkin* mutant adult flies and *parkin* mutants with elevated *MTF-1* expression (p<0.0001).

Figure 8. *MTF-1* reduces reactive oxygen species (ROS) levels in *parkin* mutants and restores metal homeostasis. (A) *park*^{25/25} flies (NF) show high amounts of ROS. *MTF-1* transgene expression, or treatment with chelators of redox-active metals (BPS and BCS) reduce ROS levels. *parkin* heterozygous controls (*park*^{25/+}) do not show significant differences in ROS levels with or without *MTF-1*-overexpression or dietary supplementation with metal chelators. The *MTF-1* knockout shows the highest ROS level. ***, p<0.001 and **, p<0.01, chelator-supplemented compared to normal food for the same genotype (black stars), or compared to the same treatment between genotypes (red stars). (B) Elevated *MTF-1* expression in a *parkin* mutant background restores normal basal levels of the essential trace metals copper, iron and zinc when flies are raised on NF (normal food). (C) Differences in metal content between controls, *park*^{25/25} and *tub-MTF-1*, *park*^{25/25} flies are largely leveled out by metal supplementation of the food.

Figure 9. Strong expression of human *MTF-1* or of *Drosophila* metallothionein prolongs lifespan of *parkin* mutants. A human-*MTF-1* transgene driven by *actin-Gal4* extends the median lifespan of *parkin* mutant flies from 7 (red) to 19 days (light blue). Direct overexpression of metallothionein *MtnA* by tubulin promoter also enhances the median lifespan of *parkin* mutants up to 17 days (green). An even better extension up to a median of 21 days is observed with a tubulin-driven transgene of *Drosophila* *MTF-1* (dark blue). *park*^{25/+} (black) and *park*^{25/25} (red) serve as controls.

Table I. (A) Partial rescue of synthetic lethality of combined *parkin* and *MTF-1* mutants by N-acetylcysteine: *MTF-1*, *parkin* heterozygous parents were crossed *inter se* on food supplemented with 15mM NAC. **(B) Increased frequency of flies reaching adulthood among *parkin* mutants overexpressing MTF-1.** *park*^{25/25} mutants were obtained from *parkin* heterozygous parents crossed *inter se*. In both (A) and (B), egg laying was allowed for two days by equal numbers of parents and eclosing progeny flies were counted thereafter. Three independent crosses were done.

References

1. **Abbott, R. D., G. W. Ross, L. R. White, W. T. Sanderson, C. M. Burchfiel, M. Kashon, D. S. Sharp, K. H. Masaki, J. D. Curb, and H. Petrovitch.** 2003. Environmental, life-style, and physical precursors of clinical Parkinson's disease: recent findings from the Honolulu-Asia Aging Study. *J Neurol* **250 Suppl 3**:III30-9.
2. **Andrews, G. K.** 2000. Regulation of metallothionein gene expression by oxidative stress and metal ions. *Biochem Pharmacol* **59**:95-104.
3. **Auf der Maur, A., T. Belser, G. Elgar, O. Georgiev, and W. Schaffner.** 1999. Characterization of the transcription factor MTF-1 from the Japanese pufferfish (*Fugu rubripes*) reveals evolutionary conservation of heavy metal stress response. *Biol Chem* **380**:175-85.
4. **Bahadorani, S., S. Mukai, D. Egli, and A. J. Hilliker.** 2010. Overexpression of metal-responsive transcription factor (MTF-1) in *Drosophila melanogaster* ameliorates life-span reductions associated with oxidative stress and metal toxicity. *Neurobiol Aging* **31**:1215-26.
5. **Balabanova, M., L. Popova, and R. Tchipeva.** 2003. Dyes in dermatology. *Clin Dermatol* **21**:2-6.
6. **Balamurugan, K., D. Egli, A. Selvaraj, B. Zhang, O. Georgiev, and W. Schaffner.** 2004. Metal-responsive transcription factor (MTF-1) and heavy metal stress response in *Drosophila* and mammalian cells: a functional comparison. *Biol Chem* **385**:597-603.
7. **Brugnera, E., O. Georgiev, F. Radtke, R. Heuchel, E. Baker, G. R. Sutherland, and W. Schaffner.** 1994. Cloning, chromosomal mapping and

- characterization of the human metal-regulatory transcription factor MTF-1. *Nucleic Acids Res* **22**:3167-73.
8. **Bueler, H.** 2010. Mitochondrial dynamics, cell death and the pathogenesis of Parkinson's disease. *Apoptosis*.
 9. **Chen, W. Y., J. A. John, C. H. Lin, and C. Y. Chang.** 2002. Molecular cloning and developmental expression of zinc finger transcription factor MTF-1 gene in zebrafish, *Danio rerio*. *Biochem Biophys Res Commun* **291**:798-805.
 10. **Cookson, M. R., G. Xiromerisiou, and A. Singleton.** 2005. How genetics research in Parkinson's disease is enhancing understanding of the common idiopathic forms of the disease. *Curr Opin Neurol* **18**:706-11.
 11. **Culotta, V. C., M. Yang, and T. V. O'Halloran.** 2006. Activation of superoxide dismutases: putting the metal to the pedal. *Biochim Biophys Acta* **1763**:747-58.
 12. **Dalton, T. P., Q. Li, D. Bittel, L. Liang, and G. K. Andrews.** 1996. Oxidative stress activates metal-responsive transcription factor-1 binding activity. Occupancy in vivo of metal response elements in the metallothionein-I gene promoter. *J Biol Chem* **271**:26233-41.
 13. **Dawson, T. M., and V. L. Dawson.** 2003. Molecular pathways of neurodegeneration in Parkinson's disease. *Science* **302**:819-22.
 14. **Dexter, D. T., C. J. Carter, F. R. Wells, F. Javoy-Agid, Y. Agid, A. Lees, P. Jenner, and C. D. Marsden.** 1989. Basal lipid peroxidation in substantia nigra is increased in Parkinson's disease. *J Neurochem* **52**:381-9.
 15. **Ebadi, M., H. Brown-Borg, H. El Refaey, B. B. Singh, S. Garrett, S. Shavali, and S. K. Sharma.** 2005. Metallothionein-mediated neuroprotection in genetically engineered mouse models of Parkinson's disease. *Brain Res Mol Brain Res* **134**:67-75.
 16. **Egli, D., J. Domenech, A. Selvaraj, K. Balamurugan, H. Hua, M. Capdevila, O. Georgiev, W. Schaffner, and S. Atrian.** 2006b. The four members of the *Drosophila* metallothionein family exhibit distinct yet overlapping roles in heavy metal homeostasis and detoxification. *Genes Cells* **11**:647-58.
 17. **Egli, D., A. Selvaraj, H. Yepiskoposyan, B. Zhang, E. Hafen, O. Georgiev, and W. Schaffner.** 2003. Knockout of 'metal-responsive transcription factor'

- MTF-1 in *Drosophila* by homologous recombination reveals its central role in heavy metal homeostasis. *Embo J* **22**:100-8.
18. **Egli, D., H. Yepiskoposyan, A. Selvaraj, K. Balamurugan, R. Rajaram, A. Simons, G. Multhaup, S. Mettler, A. Vardanyan, O. Georgiev, and W. Schaffner.** 2006a. A family knockout of all four *Drosophila* metallothioneins reveals a central role in copper homeostasis and detoxification. *Mol Cell Biol* **26**:2286-96.
 19. **Gandhi, S., A. Wood-Kaczmar, Z. Yao, H. Plun-Favreau, E. Deas, K. Klupsch, J. Downward, D. S. Latchman, S. J. Tabrizi, N. W. Wood, M. R. Duchen, and A. Y. Abramov.** 2009. PINK1-associated Parkinson's disease is caused by neuronal vulnerability to calcium-induced cell death. *Mol Cell* **33**:627-38.
 20. **Gasser, T.** 2005. Genetics of Parkinson's disease. *Curr Opin Neurol* **18**:363-9.
 21. **Ghoshal, K., and S. Jacob.** 2000. Regulation of metallothionein gene expression. *Progress in Nucleic Acid Research and Molecular Biology* **66**:357-384.
 22. **Ghoshal, K., S. Majumder, Q. Zhu, J. Hunzeker, J. Datta, M. Shah, J. F. Sheridan, and S. T. Jacob.** 2001. Influenza virus infection induces metallothionein gene expression in the mouse liver and lung by overlapping but distinct molecular mechanisms. *Mol Cell Biol* **21**:8301-17.
 23. **Greene, J. C., A. J. Whitworth, L. A. Andrews, T. J. Parker, and L. J. Pallanck.** 2005. Genetic and genomic studies of *Drosophila* parkin mutants implicate oxidative stress and innate immune responses in pathogenesis. *Hum Mol Genet* **14**:799-811.
 24. **Greene, J. C., A. J. Whitworth, I. Kuo, L. A. Andrews, M. B. Feany, and L. J. Pallanck.** 2003. Mitochondrial pathology and apoptotic muscle degeneration in *Drosophila* parkin mutants. *Proc Natl Acad Sci U S A* **100**:4078-83.
 25. **Gunes, C., R. Heuchel, O. Georgiev, K. H. Muller, P. Lichtlen, H. Bluthmann, S. Marino, A. Aguzzi, and W. Schaffner.** 1998. Embryonic lethality and liver degeneration in mice lacking the metal-responsive transcriptional activator MTF-1. *Embo J* **17**:2846-54.
 26. **Haque, M. E., K. J. Thomas, C. D'Souza, S. Callaghan, T. Kitada, R. S. Slack, P. Fraser, M. R. Cookson, A. Tandon, and D. S. Park.** 2008.

- Cytoplasmic Pink1 activity protects neurons from dopaminergic neurotoxin MPTP. *Proc Natl Acad Sci U S A* **105**:1716-21.
27. **Heuchel, R., F. Radtke, O. Georgiev, G. Stark, M. Aguet, and W. Schaffner.** 1994. The transcription factor MTF-1 is essential for basal and heavy metal-induced metallothionein gene expression. *Embo J* **13**:2870-5.
 28. **Jenner, P., and C. W. Olanow.** 1996. Oxidative stress and the pathogenesis of Parkinson's disease. *Neurology* **47**:S161-70.
 29. **Kagi, J. H.** 1991. Overview of metallothionein. *Methods Enzymol* **205**:613-26.
 30. **Klaassen, C. D., J. Liu, and S. Choudhuri.** 1999. Metallothionein: an intracellular protein to protect against cadmium toxicity. *Annu Rev Pharmacol Toxicol* **39**:267-94.
 31. **La Fontaine, S., and J. F. Mercer.** 2007. Trafficking of the copper-ATPases, ATP7A and ATP7B: role in copper homeostasis. *Arch Biochem Biophys* **463**:149-67.
 32. **Langmade, S. J., R. Ravindra, P. J. Daniels, and G. K. Andrews.** 2000. The transcription factor MTF-1 mediates metal regulation of the mouse ZnT1 gene. *J Biol Chem* **275**:34803-9.
 33. **Levy, M. A., Y. H. Tsai, A. Reaume, and T. M. Bray.** 2001. Cellular response of antioxidant metalloproteins in Cu/Zn SOD transgenic mice exposed to hyperoxia. *Am J Physiol Lung Cell Mol Physiol* **281**:L172-82.
 34. **Lichtlen, P., and W. Schaffner.** 2001. Putting its fingers on stressful situations: the heavy metal-regulatory transcription factor MTF-1. *Bioessays* **23**:1010-7.
 35. **Lin, H.** 1998. The self-renewing mechanism of stem cells in the germline. *Curr Opin Cell Biol* **10**:687-93.
 36. **Lindert, U., M. Cramer, M. Meuli, O. Georgiev, and W. Schaffner.** 2009. Metal-responsive transcription factor 1 (MTF-1) activity is regulated by a nonconventional nuclear localization signal and a metal-responsive transactivation domain. *Mol Cell Biol* **29**:6283-93.
 37. **Mathews, W. R., F. Wang, D. J. Eide, and M. Van Doren.** 2005. *Drosophila* fear of intimacy encodes a Zrt/IRT-like protein (ZIP) family zinc transporter functionally related to mammalian ZIP proteins. *J Biol Chem* **280**:787-95.

38. **Miura, T., S. Muraoka, and T. Ogiso.** 1997. Antioxidant activity of metallothionein compared with reduced glutathione. *Life Sci* **60**:PL 301-9.
39. **Miyazaki, I., M. Asanuma, H. Hozumi, K. Miyoshi, and N. Sogawa.** 2007. Protective effects of metallothionein against dopamine quinone-induced dopaminergic neurotoxicity. *FEBS Lett* **581**:5003-8.
40. **Mokdad, R., A. Debec, and M. Wegnez.** 1987. Metallothionein genes in *Drosophila melanogaster* constitute a dual system. *Proc Natl Acad Sci U S A* **84**:2658-62.
41. **Murphy, B. J., G. K. Andrews, D. Bittel, D. J. Discher, J. McCue, C. J. Green, M. Yanovsky, A. Giaccia, R. M. Sutherland, K. R. Laderoute, and K. A. Webster.** 1999. Activation of metallothionein gene expression by hypoxia involves metal response elements and metal transcription factor-1. *Cancer Res* **59**:1315-22.
42. **Nelson, N.** 1999. Metal ion transporters and homeostasis. *Embo J* **18**:4361-71.
43. **Olanow, C. W., and W. G. Tatton.** 1999. Etiology and pathogenesis of Parkinson's disease. *Annu Rev Neurosci* **22**:123-44.
44. **Otsuka, F., I. Okugaito, M. Ohsawa, A. Iwamatsu, K. Suzuki, and S. Koizumi.** 2000. Novel responses of ZRF, a variant of human MTF-1, to in vivo treatment with heavy metals. *Biochim Biophys Acta* **1492**:330-40.
45. **Palacino, J. J., D. Sagi, M. S. Goldberg, S. Krauss, C. Motz, M. Wacker, J. Klose, and J. Shen.** 2004. Mitochondrial dysfunction and oxidative damage in parkin-deficient mice. *J Biol Chem* **279**:18614-22.
46. **Palmiter, R. D.** 1998. The elusive function of metallothioneins. *Proc Natl Acad Sci U S A* **95**:8428-30.
47. **Pendleton, R. G., F. Parvez, M. Sayed, and R. Hillman.** 2002. Effects of pharmacological agents upon a transgenic model of Parkinson's disease in *Drosophila melanogaster*. *J Pharmacol Exp Ther* **300**:91-6.
48. **Pesah, Y., T. Pham, H. Burgess, B. Middlebrooks, P. Verstreken, Y. Zhou, M. Harding, H. Bellen, and G. Mardon.** 2004. *Drosophila* parkin mutants have decreased mass and cell size and increased sensitivity to oxygen radical stress. *Development* **131**:2183-94.
49. **Petrovitch, H., G. W. Ross, R. D. Abbott, W. T. Sanderson, D. S. Sharp, C. M. Tanner, K. H. Masaki, P. L. Blanchette, J. S. Popper, D. Foley, L.**

- Launer, and L. R. White.** 2002. Plantation work and risk of Parkinson disease in a population-based longitudinal study. *Arch Neurol* **59**:1787-92.
50. **Quaife, C. J., S. D. Findley, J. C. Erickson, G. J. Froelick, E. J. Kelly, B. P. Zambrowicz, and R. D. Palmiter.** 1994. Induction of a new metallothionein isoform (MT-IV) occurs during differentiation of stratified squamous epithelia. *Biochemistry* **33**:7250-9.
 51. **Radtke, F., R. Heuchel, O. Georgiev, M. Hergersberg, M. Gariglio, Z. Dembic, and W. Schaffner.** 1993. Cloned transcription factor MTF-1 activates the mouse metallothionein I promoter. *Embo J* **12**:1355-62.
 52. **Riparbelli, M. G., and G. Callaini.** 2007. The Drosophila parkin homologue is required for normal mitochondrial dynamics during spermiogenesis. *Dev Biol* **303**:108-20.
 53. **Rossi, L., M. Arciello, C. Capo, and G. Rotilio.** 2006. Copper imbalance and oxidative stress in neurodegeneration. *Ital J Biochem* **55**:212-21.
 54. **Rouault, T. A., and S. Cooperman.** 2006. Brain iron metabolism. *Semin Pediatr Neurol* **13**:142-8.
 55. **Saini, N., S. Oelhafen, H. Hua, O. Georgiev, W. Schaffner, and H. Bueler.** 2010. Extended lifespan of Drosophila parkin mutants through sequestration of redox-active metals and enhancement of anti-oxidative pathways. *Neurobiol Dis* **40**:82-92.
 56. **Saini, N., and W. Schaffner.** 2010. Zinc supplement greatly improves the condition of parkin mutant Drosophila. *Biol Chem* **391**:513-8.
 57. **Sang, T. K., H. Y. Chang, G. M. Lawless, A. Ratnaparkhi, L. Mee, L. C. Ackerson, N. T. Maidment, D. E. Krantz, and G. R. Jackson.** 2007. A Drosophila model of mutant human parkin-induced toxicity demonstrates selective loss of dopaminergic neurons and dependence on cellular dopamine. *J Neurosci* **27**:981-92.
 58. **Saydam, N., O. Georgiev, M. Y. Nakano, U. F. Greber, and W. Schaffner.** 2001. Nucleo-cytoplasmic trafficking of metal-regulatory transcription factor 1 is regulated by diverse stress signals. *J Biol Chem* **276**:25487-95.
 59. **Schulz, J. B.** 2008. Update on the pathogenesis of Parkinson's disease. *J Neurol* **255 Suppl 5**:3-7.
 60. **Selvaraj, A., K. Balamurugan, H. Yepiskoposyan, H. Zhou, D. Egli, O. Georgiev, D. J. Thiele, and W. Schaffner.** 2005. Metal-responsive

- transcription factor (MTF-1) handles both extremes, copper load and copper starvation, by activating different genes. *Genes Dev* **19**:891-6.
61. **Silar, P., L. Theodore, R. Mokdad, N. E. Erraiss, A. Cadic, and M. Wegnez.** 1990. Metallothionein Mto gene of *Drosophila melanogaster*: structure and regulation. *J Mol Biol* **215**:217-24.
 62. **Smirnova, I. V., D. C. Bittel, R. Ravindra, H. Jiang, and G. K. Andrews.** 2000. Zinc and cadmium can promote rapid nuclear translocation of metal response element-binding transcription factor-1. *J Biol Chem* **275**:9377-84.
 63. **Southon, A., R. Burke, M. Norgate, P. Batterham, and J. Camakaris.** 2004. Copper homeostasis in *Drosophila melanogaster* S2 cells. *Biochem J* **383**:303-9.
 64. **Stuart, G. W., P. F. Searle, H. Y. Chen, R. L. Brinster, and R. D. Palmiter.** 1984. A 12-base-pair DNA motif that is repeated several times in metallothionein gene promoters confers metal regulation to a heterologous gene. *Proc Natl Acad Sci U S A* **81**:7318-22.
 65. **Tumer, Z., and N. Horn.** 1998. Menkes disease: underlying genetic defect and new diagnostic possibilities. *J Inherit Metab Dis* **21**:604-12.
 66. **Valko, M., H. Morris, and M. T. Cronin.** 2005. Metals, toxicity and oxidative stress. *Curr Med Chem* **12**:1161-208.
 67. **Vandesompele, J., K. De Preter, F. Pattyn, B. Poppe, N. Van Roy, A. De Paepe, and F. Speleman.** 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* **3**:RESEARCH0034.
 68. **Wang, Y., I. Lorenzi, O. Georgiev, and W. Schaffner.** 2004. Metal-responsive transcription factor-1 (MTF-1) selects different types of metal response elements at low vs. high zinc concentration. *Biol Chem* **385**:623-32.
 69. **Wang, Y., U. Wimmer, P. Lichtlen, D. Inderbitzin, B. Stieger, P. J. Meier, L. Hunziker, T. Stallmach, R. Forrer, T. Rulicke, O. Georgiev, and W. Schaffner.** 2004. Metal-responsive transcription factor-1 (MTF-1) is essential for embryonic liver development and heavy metal detoxification in the adult liver. *Faseb J* **18**:1071-9.
 70. **West, A. B., and N. T. Maidment.** 2004. Genetics of parkin-linked disease. *Hum Genet* **114**:327-36.

71. **West, A. K., R. Stallings, C. E. Hildebrand, R. Chiu, M. Karin, and R. I. Richards.** 1990. Human metallothionein genes: structure of the functional locus at 16q13. *Genomics* **8**:513-8.
72. **Whitworth, A. J., D. A. Theodore, J. C. Greene, H. Benes, P. D. Wes, and L. J. Pallanck.** 2005. Increased glutathione S-transferase activity rescues dopaminergic neuron loss in a *Drosophila* model of Parkinson's disease. *Proc Natl Acad Sci U S A* **102**:8024-9.
73. **Winklhofer, K. F., and C. Haass.** 2010. Mitochondrial dysfunction in Parkinson's disease. *Biochim Biophys Acta* **1802**:29-44.
74. **Yepiskoposyan, H., D. Egli, T. Fergestad, A. Selvaraj, C. Treiber, G. Multhaup, O. Georgiev, and W. Schaffner.** 2006. Transcriptome response to heavy metal stress in *Drosophila* reveals a new zinc transporter that confers resistance to zinc. *Nucleic Acids Res* **34**:4866-77.
75. **Zhang, B., D. Egli, O. Georgiev, and W. Schaffner.** 2001. The *Drosophila* homolog of mammalian zinc finger factor MTF-1 activates transcription in response to heavy metals. *Mol Cell Biol* **21**:4505-14.
76. **Zhang, B., O. Georgiev, M. Hagmann, C. Gunes, M. Cramer, P. Faller, M. Vasak, and W. Schaffner.** 2003. Activity of metal-responsive transcription factor 1 by toxic heavy metals and H₂O₂ in vitro is modulated by metallothionein. *Mol Cell Biol* **23**:8471-85.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Table 1A

15 mM NAC supplemented food	$\text{♀ } w; \text{Sp/CyO}; \text{MTF-1}^-, \text{park}^{25}/\text{TM3} \times \text{♂ } w; \text{Sp/CyO}; \text{MTF-1}^-, \text{park}^{25}/\text{TM3}$			
	<i>Parkin</i> and <i>MTF-1</i> heterozygous progeny: <i>w; Sp/CyO; MTF-1</i> ⁻ , <i>park</i> ²⁵ /TM3	<i>Parkin</i> and <i>MTF-1</i> homozygous progeny: <i>w; Sp/CyO; MTF-1</i> ^{-/-} , <i>park</i> ^{25/25}	Total number of progeny	Percentage of <i>parkin</i> and <i>MTF-1</i> homozygous progeny
Cross 1	62	14	76	18.4
Cross 2	79	19	98	19.4
Cross 3	68	16	84	19.0
Total	209	49	258	19.0

Table 1B

normal food	$\text{♀ } w; +; \text{tub-MTF-1}, \text{park}^{25/+} \times \text{♂ } w; +; \text{tub-MTF-1}, \text{park}^{25/+}$			
	<i>Parkin</i> and <i>MTF-1</i> heterozygous progeny: <i>w; +; tub-MTF-1, park</i> ^{25/+}	<i>Parkin</i> and <i>MTF-1</i> homozygous progeny: <i>w; +; tub-MTF-1, park</i> ^{25/25}	Total number of progeny	Percentage of <i>parkin</i> and <i>MTF-1</i> homozygous progeny
Cross 1	310	53	363	14.6
Cross 2	741	141	882	16.0
Cross 3	229	37	266	13.9
Total	1280	231	1511	15.3

Biol Chem. 2010 May;391(5):513-8.

Zinc supplement greatly improves the condition of parkin mutant *Drosophila*.

Nidhi Saini¹ and Walter Schaffner¹.

¹Institute of Molecular Life Sciences, University of Zürich, Switzerland.

Abstract

Parkinson's disease (PD) is a progressive neurodegenerative disorder in which oxidative stress is implicated as a major causative factor. Mutations in the gene encoding Parkin, a ubiquitin ligase, are responsible for a familial form of PD. In a *Drosophila* disease model lacking Parkin (*park(25)* null mutant), we tested the effect of zinc supplementation. Zinc is an essential trace metal and a component of many enzymes and transcriptional regulators. Unlike copper and iron, zinc is not redox-active and under most conditions serves as an antioxidant. We find that the condition of parkin mutants raised on zinc-supplemented food is greatly improved. At zinc concentrations where controls begin to show adverse effects as a result of the metal supplement, parkin mutants perform best, as manifested in a higher frequency of reaching adulthood, extended lifespan and improved motoric abilities.

Short Communication

Zinc supplement greatly improves the condition of *parkin* mutant *Drosophila*

Nidhi Saini and Walter Schaffner*

Institute of Molecular Life Sciences, University of Zürich,
Winterthurerstrasse 190, CH-8051 Zürich, Switzerland

* Corresponding author
e-mail: walter.schaffner@imls.uzh.ch

Abstract

Parkinson's disease (PD) is a progressive neurodegenerative disorder in which oxidative stress is implicated as a major causative factor. Mutations in the gene encoding Parkin, a ubiquitin ligase, are responsible for a familial form of PD. In a *Drosophila* disease model lacking Parkin (*park*²⁵ null mutant), we tested the effect of zinc supplementation. Zinc is an essential trace metal and a component of many enzymes and transcriptional regulators. Unlike copper and iron, zinc is not redox-active and under most conditions serves as an antioxidant. We find that the condition of *parkin* mutants raised on zinc-supplemented food is greatly improved. At zinc concentrations where controls begin to show adverse effects as a result of the metal supplement, *parkin* mutants perform best, as manifested in a higher frequency of reaching adulthood, extended lifespan and improved motoric abilities.

Keywords: antioxidant; metal homeostasis; metallothioneins; MTF-1; Parkinson's disease; zinc transporters.

Parkinson's disease (PD), characterized by the loss of dopaminergic (DA) neurons in the substantia nigra, is a progressive neurodegenerative disorder with the second highest incidence rate and is the most common age-related movement disorder (Olanow and Tatton, 1999; Dawson and Dawson, 2003; Greene et al., 2003). Both genetic and environmental factors contribute to its pathogenesis. Oxidative stress is considered to be a major factor in the pathogenesis of PD, as evidenced by an elevated content of redox-active iron and lipid peroxides in the diseased brain, impaired mitochondrial function, and alterations in the antioxidant defense mechanisms (Dexter et al., 1989; Jenner and Olanow, 1996; Greene et al., 2003; Pesah et al., 2004). Mutations in six genes, including *parkin* which encodes an E3 ubiquitin ligase, have been associated with rare, early-onset, familial forms of PD (West and Maidment, 2004; Gasser, 2005; Sang et al., 2007). Interestingly, some alleles of these genes might

be susceptibility factors for environmental toxins (Choi et al., 2000; Warner and Schapira, 2003; Bueler, 2009).

In our study, we used a *Drosophila melanogaster* line in which the ortholog of the human *parkin* gene is disrupted by transposition of a P-element (Greene et al., 2003; Pesah et al., 2004). *Parkin* mutant flies present with male and female sterility (Riparbelli and Callaini, 2007), mitochondrial and muscle abnormalities, locomotor defects, an inability to fly owing to degeneration of indirect flight muscles, increased sensitivity to multiple stresses, including oxidative stress, and a severely reduced lifespan (Palacino et al., 2004; Greene et al., 2005; Whitworth et al., 2005). Some of these defects arise because *parkin* mutants have dysfunctional mitochondria with disturbances in the electron transport chain. In mice, in contrast to its pivotal role in humans (Choi et al., 2000), *Parkin* function does not seem to be critical for the survival of DA neurons (Goldberg et al., 2003; Itier et al., 2003; Palacino et al., 2004). Similarly, *Drosophila* that are null mutants for *parkin* do not generally display DA neuron loss (Greene et al., 2003; Pesah et al., 2004), although a partial loss of DA neurons in the PPL1 cluster of the brain has been reported (Whitworth et al., 2005).

In the late 1980s, the antioxidant function of the redox-inert metal zinc was recognized and proposed to be mediated by the protection of protein sulfhydryl groups and/or by competing against redox-active metals (Bray and Bettger, 1990). Additionally, zinc ions can upregulate the expression of metallothioneins, which owing to their high cysteine content can serve as antioxidants. Zinc has an established antiapoptotic function that minimizes ROS-induced cellular oxidative damage (Suzuki et al., 1991). This also occurs in the central nervous system, particularly in the brain (Kocaturk et al., 1996).

More than 70 different enzymes involved in the metabolism of biomolecules require zinc as a cofactor (Parkin, 2004). Zinc is an integral part of the hundreds of transcription factors that contain zinc finger domains (Berg and Shi, 1996), and it plays a role in cellular signal transduction and in modulation of synaptic neurotransmission. Zinc is critical for the growth and regulation of cells and alterations in zinc metabolism have been implicated in causing neurological dysfunctions on the one hand, and on the other hand providing neuroprotection. Maintenance of intracellular zinc homeostasis is thus an essential requirement in all living organisms (Valiko et al., 2005).

The best characterized zinc-activated transcription factor is the metal response element-binding transcription factor-1,




Figure 1 *Drosophila parkin* mutants show an enhanced lifespan on zinc-supplemented food.

(A) Zinc supplementation (4 mM ZnCl_2) increased the longevity of the *parkin* mutant flies (*park^{25/25}*). The maximum lifespan of 11 days on normal food (NF) increased to 23 days. Control flies (*park^{25/+}*) showed an opposite effect with a reduction of lifespan on zinc food. (B) Survival of *park^{25/25}* flies and (C) of *park^{25/+}* control flies raised and maintained on NF or zinc-supplemented food. Error bars indicate standard deviation. The significance between survival curves was analyzed using the Kaplan-Meier log-rank statistical test (* $p < 0.01$). Methods: for survival assays, NF was supplemented with zinc chloride to a final concentration of 4 mM. One- to two-day-old flies (20 per vial) were maintained at 25°C on a 12:12 h light/dark cycle for each genotype in triplicate vials. Surviving flies were transferred to fresh food vials every 2 days and counted daily. As a control, flies of the same genotype were grown on food without metal supplement. In each lifespan assay testing different conditions, the controls of *park^{25/+}* and *park^{25/25}* flies raised on NF were the same. The variations in the median lifespan of control flies in different experiments can be attributed to subtle experimental variations.

also referred to as metal-responsive transcription factor-1 (MTF-1). In response to zinc, MTF-1 translocates from the cytoplasm to the nucleus where it regulates the expression of several genes, notably metallothioneins and some zinc

transporters (Andrews, 2000; Smirnova et al., 2000; Lichtlen and Schaffner, 2001; Saydam et al., 2001). MTF-1 requires elevated zinc concentration for DNA binding. This property was exploited for the activation of MTF-1-dependent pro-

Table 1 Zinc increases the frequency of *parkin* mutant *Drosophila* reaching adulthood.

Food condition	Genotype	<i>park^{25/25}</i> /total progeny	% of <i>park^{25/25}</i>
Normal food	Set 1 <i>park^{25/25}</i>	31/1116	2.8%
	Set 2 <i>park^{25/25}</i>	10/450	2.2%
4 mM Zn	Set 1 <i>park^{25/25}</i>	110/611	18%
	Set 2 <i>park^{25/25}</i>	186/956	19%

For the analysis of eclosion frequency, egg laying was allowed for 6 days and the parent population (10 males and 10 females each of *park^{25/+}*) was the same in all vials of normal food (NF) or zinc (Zn), and progeny flies were counted at the same time.




Figure 2 Gustatory assay with adult flies on normal or zinc-supplemented food with acid red dye as a marker.

(A, B) *park^{25/25}* control flies and (C, D) *park^{25/25}* flies show the same feeding behavior on normal food, exemplified by the fully red abdomens. A food supplement of 4 mM Zn (E, F) altered feeding behavior of *park^{25/25}* (only partially red abdomens) but no change in the behavior of *park^{25/25}* flies was observed (G, H) (fully red abdomens). The mutants are recognized by white eyes because of the loss of *w⁺* (red eyes) which are present in control flies.

Methods: the adult gustatory assay was essentially carried out as described by A. Hilliker and colleagues (Bahadorani, 2008). Briefly, newly eclosed flies were reared on normal food for 2–3 days, then starved for 18 h on Whatman paper soaked with distilled water. After this treatment, starved flies (20 per vial) were transferred onto zinc-supplemented food with 0.2% sulforhodamine B sodium salt (acid red) for 2 h. For control flies, culture medium was supplemented with 0.2% acid red, without metal supplement. After 2 h of feeding at optimum temperature (25°C) and relative humidity, flies were anesthetized and the degree of abdomen redness was visually inspected. Abdomen redness was used as an indicator of the amount of food taken up.

motors by zinc and other metals in a cell-free transcription system (Zhang et al., 2003).

Zinc is not redox-active, but nevertheless toxic when in excess (Beyersmann and Haase, 2001). Acute zinc toxicity is rare but has been reported (Duncan et al., 1992; Whittaker, 1998; Prasad et al., 1999). If the extracellular concentration of zinc exceeds the capacity of zinc homeostasis mechanisms, it becomes cytotoxic and an excess of free intracellular zinc can trigger apoptosis (Choi et al., 1988; Duncan et al., 1992; Kim et al., 1999; Beyersmann and Haase, 2001; Wilhelm et al., 2001; Beyersmann, 2002; Walther et al., 2003). Zinc transport in *Drosophila*, as in vertebrates, is mediated by two families of solute linked carrier proteins: zinc importers (ZIPs), which function in the uptake of zinc to the cytoplasm, and zinc exporters (ZnTs), which reduce cytoplasmic zinc concentrations by promoting zinc efflux (Liu and Cousins, 2004; Yepiskoposyan et al., 2006). More than 10 zinc transporter genes are annotated in *Drosophila melanogaster* based on their sequence similarities to vertebrate zinc transporters. The ZIP family gene *foi* (*fear of intimacy*) was characterized in *Drosophila* and shown to be a zinc importer that is critical during development (Moore et al., 1998; Mathews et al., 2005). Transcriptional responses to zinc in *Drosophila* larvae were analyzed in our laboratory (Yepiskoposyan et al., 2006). Apart from the expected upregulation of metallothioneins and the zinc exporters *ZnT35C*

and *ZnT63C*, there was also an induction of neurotransmitters, detoxification enzymes (such as glutathione S-transferase), ferritin and chaperone encoding genes.

We found that whereas *parkin* mutant flies readily feed on high-zinc food, their wild type counterparts avoid zinc-loaded food. The mutants also had an increased survival rate on zinc-supplemented food, which prompted us to investigate their response to zinc in more detail.

With standard 'normal food' (NF), *parkin* mutant flies have a median lifespan of 6 days with a maximum of 11 days (Figure 1A). Zinc supplementation in the form of zinc chloride increased the lifespan of *parkin* mutants. When maintained on supplements of 4 mM Zn, the mutant flies survived up to 23 days with a median lifespan of 8 days (Figure 1). This increase was as a result of zinc and not chloride ions, as similar survival assays on NaCl-supplemented medium did not extend lifespan (data not shown). Zinc supplementation also increases the eclosion frequency of *parkin* mutants, from 2.5% to 19%, i.e., close to 25% which is the expected frequency of *parkin* mutants in the progeny of *parkin* heterozygous parents (Table 1). In contrast, heterozygous control flies did not draw any benefit from zinc-supplemented food: if kept on food with 4 mM $ZnCl_2$, the median lifespan reduced significantly to 17 days, whereas on NF, 80% were still alive at the end of the experiment (23 days) (Figure 1A). Similar adverse effects of zinc load were

observed with wild type *yw* and *OregonR* flies (data not shown). To determine if zinc has a stronger effect during development from eggs to adults or during the adult feeding stage, we followed the lifespan of both *parkin* mutant and *parkin* heterozygous flies under four conditions: (i) development and adult maintenance in zinc supplement, (ii) raising the flies on NF until adult stage and then maintaining them on zinc-supplemented food post-eclosion, (iii) raising the flies on zinc food but maintaining them on NF after eclosion, and (iv) raising and maintaining the flies on NF. We observed that the strongest positive effect of zinc on *park^{25/25}* flies was when they were both raised and maintained on 4 mM Zn-supplemented food (Figure 1B). The strongest negative effect of zinc on *park^{25/+}* control flies was also observed under these conditions (Figure 1C). Development of *park^{25/+}* and *OregonR* wild type flies was delayed but generally less affected by zinc than the survival of adults; egg transfer (200 each) from NF to vials containing increasing concentrations of zinc resulted in an equal percentage of eclosing adults in NF, 4 and 6 mM Zn but none developed in 10 mM Zn (data not shown; see also Egli et al., 2003).

parkin mutants might sense a deficiency in and/or a rescuing effect of zinc, and in response to this eat normal amounts of zinc-supplemented food unlike heterozygous or wild type flies. This was indicated by a visual inspection of adult *Drosophila* in a gustatory test and was quantitatively confirmed by the measurement of total zinc uptake by flies (Figures 2 and 3). On NF, the zinc content in heads of *parkin* mutant flies was much lower than in controls, consistent with a zinc deficiency in the mutants. Zinc supplementation indeed resulted in an, albeit minor, increase of zinc content in mutant heads (Figure 3A), whereas in thoraces the zinc content was similar in mutant and control flies (Figure 3B). Interestingly, the abdomens of *park^{25/25}* flies fed on zinc food showed a 10-fold increase in zinc levels compared with flies fed on NF (Figure 3C). Such an effect was, however, not observed in the control flies which only showed a minor increase in zinc levels upon zinc feeding, which is in line with their tendency to avoid zinc-loaded food.

We also determined the transcript levels of *parkin*, MTF-1 and some other genes involved in zinc import/export (Figure 4). As expected, *parkin* mutants have no detectable *parkin* transcripts. In comparison to control flies, transcript levels of the metallothionein *MtnB* are higher in *parkin* mutants (Figure 4A,B). Zinc supplementation induced *MtnB* transcripts 24-fold in *parkin* mutants; in control flies the increase was 14-fold (Figure 4B). Boosting expression of metallothioneins, which act as ROS scavengers, is one way zinc could play an antioxidant role. The elevated metallothionein levels could also explain at least in part why an increased zinc concentration is not toxic to the *parkin* mutants. We also determined the transcript levels of three zinc transporters, the exporters *ZnT35C* and *ZnT63C* and the importer *foi* (Mathews et al., 2005; Yepiskoposyan et al., 2006) both in *parkin* mutants and in the heterozygous controls. The most conspicuous difference was observed with the exporter gene *ZnT35C*, a known target of MTF-1. In *parkin* mutants, expression was dramatically reduced com-





Figure 3 Quantification of Zn content in the different body parts of *park^{25/+}* control flies and *park^{25/25}* flies.

Flies were raised on normal food (NF) or zinc-supplemented food (4 mM Zn). (A) Heads, (B) thoraces, (C) abdomens.

Methods: female flies were allowed to lay eggs on zinc-supplemented (4 mM ZnCl₂) fly food for 6 days and removed afterwards. The resulting progeny was collected at regular intervals and the percentage of eclosing *parkin* homozygous mutants and heterozygous controls were calculated each time. These progeny were then frozen and stored until the required number of 200 flies was obtained in triplicate for each genotype. Frozen flies were then dissected to separate their heads, thoraces and abdomens which were collected separately. Each sample set was then subjected to homogenization using cold protein homogenization buffer (0.32 mM sucrose, 20 mM HEPES, 1 mM MgCl₂, 0.5 mM PMSF protease inhibitor at pH 7.4) and the samples were normalized for protein content. A highly sensitivity flame atomic absorption spectrophotometer (FAAS; GTA-120/PSD-120, Varian Australia Pty Ltd, Mulgrave, VIC, Australia) was used to detect the zinc content in each body part of each genotype assessed.

pared to controls and not responsive to zinc. In heterozygous controls, zinc supplement resulted in a 1.5-fold upregulation, which indicates a response to zinc overload (Figure 4C). An increased expression of zinc exporters in control flies is in



agreement with their avoidance of zinc-supplemented food. Transcripts of *ZnT63C* and, somewhat unexpectedly of *foi*, were also lower in *parkin* mutants compared to *parkin* heterozygous controls (data not shown). Taken together, these data suggest that zinc homeostasis is distorted in *parkin* mutants, with the effect that zinc supplement in food produces a strong phenotypic rescue effect. It remains to be seen whether zinc supplementation also has a positive effect in a mouse model of familial PD and ultimately in Parkinson's patients.

Acknowledgments

We thank Drs. Jessica Greene and Leo Pallanck (University of Washington) for providing the *park^{25/25}* flies, Drs. Eva Freisinger and Augusto Cabral for the metal quantification experiments, Martin Moser for the RT-PCR analysis, Ivan Ostojic (Friedrich Miescher Institute, Basel, Switzerland) for statistical analyses, Till Strassen for maintenance of fly stocks and Dr. George Hausmann for critical

Figure 4 Real-time analysis of *parkin* transcript levels.

(A) In *park^{25/25}* flies no *parkin* transcripts are detectable. In *park^{25+/+}* *parkin* transcripts are, if anything, elevated in flies raised on zinc food. (B) *park^{25/25}* flies have increased levels of metallothionein *MtnB* transcripts in NF compared to control flies. Zinc induces *MtnB* in both genotypes, but to a higher level in *park^{25/25}*. (C) The zinc exporter, *ZnT35C*, shows increased transcript levels in *park^{25+/+}* on zinc food, which is not observed in *park^{25/25}* flies.

Methods: total RNA was purified from the adult *Drosophila* tissue using the nucleospin RNA II protocol and eluted in 60 µl RNase-free water. cDNA was prepared using the transcriptor high fidelity cDNA synthesis kit from Roche (Mannheim, Germany). The cDNA obtained was further purified using the AM 1906 DNA-free kit (Ambion, Rotkreuz, Switzerland). This was then used for analysis by real-time PCR on the Tecan Genesis 200/8 robot (Tecan, Männedorf, Switzerland) using the Mesa Green qPCR Mastermix Plus (Eurogentec, Seraing, Belgium) for SYBR assays. The qPCR run was performed on an Applied Biosystems machine (ABI Prism SDS 7900 HAT; Seraing, Belgium) in a 384-well format with a reaction volume of 10 µl. All fold-change values are normalized to corresponding *park^{25+/+}* values on normal food (NF). Housekeeping genes used for reference were *actin5C*, *sec24* and *GAPDH*. The primer sequences used for the transcripts quantified were: for *parkin*, 5'-AAG ATC ATA TTT GCC GGT AAG GAA-3' and 5'-CGC TTT GCT GAC CCA AGT C-3' amplify a 73-bp fragment only from the *parkin* heterozygous control flies; for *MtnB*, 5'-TTG CAA GGG TTG TGG AAC AAA-3' and 5'-TGC AGG CGC AGT TGT CC-3' amplify a 65-bp fragment, and for *ZnT35C*, 5'-GCT CTC GCC GAT CTG AAA GT-3' and 5'-CGC ACC GAC AAG TGT TTC TTA TA-3' amplify a 75-bp fragment.

reading of the manuscript. This work was supported by the Kanton Zürich and the Swiss National Science Foundation.

References

- Andrews, G.K. (2000). Regulation of metallothionein gene expression by oxidative stress and metal ions. *Biochem. Pharmacol.* 59, 95–104.
- Berg, J.M. and Shi, Y. (1996). The galvanization of biology: a growing appreciation for the roles of zinc. *Science* 271, 1081–1085.
- Beyersmann, D. (2002). Effects of carcinogenic metals on gene expression. *Toxicol. Lett.* 127, 63–68.
- Beyersmann, D. and Haase, H. (2001). Functions of zinc in signaling, proliferation and differentiation of mammalian cells. *Bio-metals* 14, 331–341.
- Bray, T.M. and Bettger, W.J. (1990). The physiological role of zinc as an antioxidant. *Free Radic. Biol. Med.* 8, 281–291.
- Bueler, H. (2009). Impaired mitochondrial dynamics and function in the pathogenesis of Parkinson's disease. *Exp. Neurol.* 218, 235–246.
- Choi, D.W., Yokoyama, M., and Koh, J. (1988). Zinc neurotoxicity in cortical cell culture. *Neuroscience* 24, 67–79.
- Choi, H.Y., Song, J.H., Park, D.K., and Ross, G.M. (2000). The effects of ascorbic acid on dopamine-induced death of PC12 cells are dependent on exposure kinetics. *Neurosci. Lett.* 296, 81–84.
- Dawson, T.M. and Dawson, V.L. (2003). Molecular pathways of neurodegeneration in Parkinson's disease. *Science* 302, 819–822.

- Dexter, D.T., Carter, C.J., Wells, F.R., Javoy-Agid, F., Agid, Y., Lees, A., Jenner, P., and Marsden, C.D. (1989). Basal lipid peroxidation in substantia nigra is increased in Parkinson's disease. *J. Neurochem.* 52, 381–389.
- Duncan, M.W., Marini, A.M., Watters, R., Kopin, I.J., and Markey, S.P. (1992). Zinc, a neurotoxin to cultured neurons, contaminates cycad flour prepared by traditional guamanian methods. *J. Neurosci.* 12, 1523–1537.
- Egli, D., Selvaraj, A., Yepiskoposyan, H., Zhang, B., Hafen, E., Georgiev, O., and Schaffner, W. (2003). Knockout of 'metal-responsive transcription factor' MTF-1 in *Drosophila* by homologous recombination reveals its central role in heavy metal homeostasis. *EMBO J.* 22, 100–108.
- Gasser, T. (2005). Genetics of Parkinson's disease. *Curr. Opin. Neurol.* 18, 363–369.
- Goldberg, M.S., Fleming, S.M., Palacino, J.J., Cepeda, C., Lam, H.A., Bhatnagar, A., Meloni, E.G., Wu, N., Ackerson, L.C., Klapstein, G.J., et al. (2003). Parkin-deficient mice exhibit nigrostriatal deficits but not loss of dopaminergic neurons. *J. Biol. Chem.* 278, 43628–43635.
- Greene, J.C., Whitworth, A.J., Kuo, L., Andrews, L.A., Feany, M.B., and Pallanck, L.J. (2003). Mitochondrial pathology and apoptotic muscle degeneration in *Drosophila* parkin mutants. *Proc. Natl. Acad. Sci. USA* 100, 4078–4083.
- Greene, J.C., Whitworth, A.J., Andrews, L.A., Parker, T.J., and Pallanck, L.J. (2005). Genetic and genomic studies of *Drosophila* parkin mutants implicate oxidative stress and innate immune responses in pathogenesis. *Hum. Mol. Genet.* 14, 799–811.
- Ilier, J.M., Ibanez, P., Mena, M.A., Abbas, N., Cohen-Salmon, C., Bohme, G.A., Laville, M., Pratt, J., Corti, O., Pradier, L., et al. (2003). Parkin gene inactivation alters behaviour and dopamine neurotransmission in the mouse. *Hum. Mol. Genet.* 12, 2277–2291.
- Jenner, P. and Olanow, C.W. (1996). Oxidative stress and the pathogenesis of Parkinson's disease. *Neurology* 47, S161–S170.
- Kim, E.Y., Koh, J.Y., Kim, Y.H., Sohn, S., Joe, E., and Gwag, B.J. (1999). Zn^{2+} entry produces oxidative neuronal necrosis in cortical cell cultures. *Eur. J. Neurosci.* 11, 327–334.
- Kocaturk, S., Kocaturk, P.A., Kavas, G.O., and Mutluer, N. (1996). Antioxidant defence system in a patient with cerebrovascular accident. *J. Int. Med. Res.* 24, 376–380.
- Lichtlen, P. and Schaffner, W. (2001). Putting its fingers on stressful situations: the heavy metal-regulatory transcription factor MTF-1. *Bioessays* 23, 1010–1017.
- Liuzzi, J.P. and Cousins, R.J. (2004). Mammalian zinc transporters. *Annu. Rev. Nutr.* 24, 151–172.
- Mathews, W.R., Wang, F., Eide, D.J., and Van Doren, M. (2005). *Drosophila* fear of intimacy encodes a Zrt/IRT-like protein (ZIP) family zinc transporter functionally related to mammalian ZIP proteins. *J. Biol. Chem.* 280, 787–795.
- Moore, L.A., Brothier, H.T., Van Doren, M., and Lehmann, R. (1998). Gonadal mesoderm and fat body initially follow a common developmental path in *Drosophila*. *Development* 125, 837–844.
- Olanow, C.W. and Tatton, W.G. (1999). Etiology and pathogenesis of Parkinson's disease. *Annu. Rev. Neurosci.* 22, 123–144.
- Palacino, J.J., Sagi, D., Goldberg, M.S., Krauss, S., Motz, C., Wacker, M., Klose, J., and Shen, J. (2004). Mitochondrial dysfunction and oxidative damage in parkin-deficient mice. *J. Biol. Chem.* 279, 18614–18622.
- Parkin, G. (2004). Synthetic analogues relevant to the structure and function of zinc enzymes. *Chem. Rev.* 104, 699–767.
- Pesah, Y., Pham, T., Burgess, H., Middlebrooks, B., Verstreken, P., Zhou, Y., Harding, M., Bellen, H., and Mardon, G. (2004). *Drosophila* parkin mutants have decreased mass and cell size and increased sensitivity to oxygen radical stress. *Development* 131, 2183–2194.
- Prasad, A.S., Beck, E.W., Kaplan, J., Chandrasekar, P.H., Ortega, J., Fitzgerald, J.T., and Swerdlow, P. (1999). Effect of zinc supplementation on incidence of infections and hospital admissions in sickle cell disease (SCD). *Am. J. Hematol.* 61, 194–202.
- Riparbelli, M.G. and Callaini, G. (2007). The *Drosophila* parkin homologue is required for normal mitochondrial dynamics during spermiogenesis. *Dev. Biol.* 303, 108–120.
- Sang, T.K., Chang, H.Y., Lawless, G.M., Ratnaparkhi, A., Mee, L., Ackerson, L.C., Maidment, N.T., Krantz, D.E., and Jackson, G.R. (2007). A *Drosophila* model of mutant human parkin-induced toxicity demonstrates selective loss of dopaminergic neurons and dependence on cellular dopamine. *J. Neurosci.* 27, 981–992.
- Saydam, N., Georgiev, O., Nakano, M.Y., Greber, U.E., and Schaffner, W. (2001). Nucleo-cytoplasmic trafficking of metal-regulatory transcription factor 1 is regulated by diverse stress signals. *J. Biol. Chem.* 276, 25487–25495.
- Smirnova, I.V., Bittel, D.C., Ravindra, R., Jiang, H., and Andrews, G.K. (2000). Zinc and cadmium can promote rapid nuclear translocation of metal response element-binding transcription factor-1. *J. Biol. Chem.* 275, 9377–9384.
- Suzuki, T., Uneyama, T., Ohma, C., Yamanaka, H., Suzuki, K., Nakajima, K., and Kimura, M. (1991). Immunohistochemical study of metallothionein in normal and benign prostatic hyperplasia of human prostate. *Prostate* 19, 35–42.
- Valko, M., Morris, H., and Cronin, M.T. (2005). Metals, toxicity and oxidative stress. *Curr. Med. Chem.* 12, 1161–1208.
- Walther, U.L., Czernak, A., Muckter, H., Walther, S.C., and Fichtl, B. (2003). Decreased GSSG reductase activity enhances cellular zinc toxicity in three human lung cell lines. *Arch. Toxicol.* 77, 131–137.
- Warner, T.T. and Schapira, A.H. (2003). Genetic and environmental factors in the cause of Parkinson's disease. *Ann. Neurol.* 53 (Suppl. 3), S16–S23; discussion S23–S25.
- West, A.B. and Maidment, N.T. (2004). Genetics of parkin-linked disease. *Hum. Genet.* 114, 327–336.
- Whittaker, P. (1998). Iron and zinc interactions in humans. *Am. J. Clin. Nutr.* 68, 442S–446S.
- Whitworth, A.J., Theodore, D.A., Greene, J.C., Benes, H., Wes, P.D., and Pallanck, L.J. (2005). Increased glutathione S-transferase activity rescues dopaminergic neuron loss in a *Drosophila* model of Parkinson's disease. *Proc. Natl. Acad. Sci. USA* 102, 8024–8029.
- Wilhelm, B., Walther, U.L., and Fichtl, B. (2001). Effects of zinc chloride on glutathione and glutathione synthesis rates in various lung cell lines. *Arch. Toxicol.* 75, 388–394.
- Yepiskoposyan, H., Egli, D., Fergestad, T., Selvaraj, A., Treiber, C., Multhaup, G., Georgiev, O., and Schaffner, W. (2006). Transcriptome response to heavy metal stress in *Drosophila* reveals a new zinc transporter that confers resistance to zinc. *Nucleic Acids Res.* 34, 4866–4877.
- Zhang, B., Georgiev, O., Hagmann, M., Gunes, C., Cramer, M., Faller, P., Vasak, M., and Schaffner, W. (2003). Activity of metal-responsive transcription factor 1 by toxic heavy metals and H_2O_2 *in vitro* is modulated by metallothionein. *Mol. Cell. Biol.* 23, 8471–8485.

Received January 15, 2010; accepted January 26, 2010

Conclusion and outlook

This work presents several new insights into the interplay between *parkin* and MTF-1 gene products and the beneficial effect of increased zinc in *parkin* mutant flies. The three major advancements outlined here are: (i) the extension of lifespan of *parkin* mutant flies by chelators of redox-active metals; (ii) characterization of the dramatic rescue effect of MTF-1 in the *parkin* deficient flies and (iii) the beneficial effect of zinc in ameliorating the effect of a *parkin* null mutation in *Drosophila*.

Since it was previously known that *parkin* mutant *Drosophila* suffer from high oxidative stress, we tested the effect of chelating redox-active metals (copper/iron) from the flies. Interestingly, as discussed before, we observed a dramatic rescue of the lifespan of the sick flies which demonstrated reduced ROS levels. Contrary to our expectations, *parkin* mutant flies fed with food containing high micromolar concentrations of redox-active metals like iron or copper did not display any obvious detrimental effects beyond their severe condition already observed with normal food.

To determine the exact reason why the *parkin* null *Drosophila* benefit from metal chelation we proceeded to investigate the effects of modulating the dosage of MTF-1 which is required for restoring homeostasis in a physiologically disturbed cellular system. Further, while testing the effect of the essential trace element, zinc on *parkin* mutants, we found the unexpected but nevertheless consistent result that zinc supplement ameliorated the *parkin* mutant phenotype, whereby the *parkin*

mutants seem to sense that high zinc levels help improve the quality of their life.

Upon MTF-1 overexpression, we were able to rescue, largely or completely, a number of phenotypic traits of a *Drosophila* model of Parkinson's disease, namely; locomotion disability associated with distorted muscle/mitochondrial morphology; significantly reduced lifespan; female infertility, and low eclosing frequency of *parkin* mutant adults. The metal responsive transcription factor MTF-1 plays a pivotal role in modulating the *parkin* phenotype: flies lacking both MTF-1 and *parkin* are not viable ("synthetic lethality"), while overexpression of MTF-1 has strong beneficial effects.

Zinc deprivation by chelation or starvation is known to cause cells to die, in many cell types by programmed cell death (Truong-Tran et al., 2001). Our results also indicate an imbalance of zinc homeostasis in the *parkin* mutant flies which can be counteracted by zinc supplement. By providing food with a high zinc concentration to the *parkin* mutant flies, we were able to rescue a few major phenotypic traits to a significant extent. These included an increase in the eclosing frequency of *parkin* mutant adults, a reduced developmental delay and an increased lifespan of mutant adults. However, a high zinc concentration in the food, while extending the lifespan of *parkin* mutants, was harmful for the *parkin* heterozygous control flies, as evidenced by a developmental delay and a reduced lifespan as well as a decreased propensity of flies to egg-laying. This seeming zinc paradox prompted us to investigate the behavioural response of both *parkin* mutants and control flies towards higher zinc levels. If the extracellular concentration of zinc exceeds the capacity of cellular zinc homeostasis, it becomes cytotoxic and the subsequent

increased concentration of intracellular zinc triggers apoptosis (Beyersmann, 2002; Beyersmann and Haase, 2001; Choi et al., 1988; Duncan et al., 1992; Kim et al., 1999; Walther et al., 2003; Wilhelm et al., 2001). Acute zinc toxicity is rare but has been reported (Duncan et al., 1992) and has been associated with reduced iron absorption and impaired immune function (Whittaker, 1998). Excess zinc may also affect the cellular status of copper and iron (Prasad et al., 1999). How the zinc leads to the improvements in several phenotypes of the mutants and how the flies perceive its potential rescue effects are still open questions. Several mechanisms are feasible. We propose that extra zinc helps to make up for the intrinsic zinc depletion faced by the *parkin* mutants during development, helping them to cope better, most likely by restoring several zinc-dependent functions.

Taken together, these data suggest the importance of metal homeostasis and antioxidant and stress relieving genes induced by MTF-1 in rescuing *parkin* mutant *Drosophila*. Hence, therapeutic approaches that modulate metal bio-availability might also ameliorate the symptoms of Parkinson's disease. This situation may not even be restricted to Parkinson's disease and might inspire research on novel therapeutic approaches to the other neurodegenerative disorders.

ACKNOWLEDGEMENTS

To begin with I would like to express my profound gratitude to my supervisor Prof. Walter Schaffner. He provided me with this exciting and highly motivating topic as my Ph.D. topic. I am sincerely thankful to him for his excellent scientific advices and thorough support both at the work and personal front. It was a pleasure to spend the time in the Lab of such a motivated mentor who extended his enthusiasm not only to experiments but to fun activities as well. The entire group is like a family thanks to his sincere efforts.

I am thankful to all the ex- and present Lab members. Sandra who was the initiator of my project, Haiqing who helped me a lot in the setting up of the project. Viola, Uschi, Lilit, Till, Dominik, Dominique, Alla, Michael and Kurt have all provided nice inputs in the Lab meetings and a warm and friendly environment at work. A special thanks goes to Dr. Oleg Georgiev and his wife Dr. Antonia Manova. They have been like my guardians here, always ready with best scientific and personal advices. It was very encouraging to work in such a stimulating and jovial environment. I would also like to thank all the people from the University who helped me with diverse experiments during my Ph.D. In addition I am grateful to my Ph.D. program (MLS Life Science Zürich) and Dr. Susanna Bachmann, the coordinator of the program.

Finally, my deepest gratitude is extended towards my Parents, sisters (Geetanjali and Swati) and my dear dog, Julie (who is no more). They have been my constant strength without whom I cannot imagine myself. I extend a special thanks to my close friend Mayank who has been a big support throughout. My friends Drs. Amrita and Kanupriya also deserve a

special mention. I thank my friends from work and home (particularly Isha, Rajesh, Ravi, Ankita, Kalina, Rebecca, Quy, Ivan, Estepni, Satinder, Garima, Sonia, Gautam, Prasuna, Vinod, Tanushree, Shradha, Shabda, Shaveta, Abhishek, Bose, Nessy, Nitika and Richa) for believing in my capabilities and for all the fun and leisure activities which motivated me during my stressful work days.

I will always be thankful to God for giving me a beautiful family and such nice friends who provide me with encouragement, support and blessings.

CURRICULUM VITAE

Surname: Saini
Name: Nidhi
Date of Birth: 05.02.1984
Nationality: Indian

Education:
Since August 2006 Ph.D. student
University of Zürich, Switzerland
Institute of Molecular Life Sciences
Supervisor: Prof. Walter Schaffner

07/2004-05/2006 Masters degree
Genetics
University of Delhi, South Campus
New Delhi, India

07/2001-05/2004 Bachelors degree
Biochemistry
University of Delhi
New Delhi, India

Title of Ph.D. thesis:

Relationship between Parkin function, heavy metal homeostasis and oxidative stress in a *Drosophila* model of Parkinson's disease

Publications:

1. Extended lifespan and improved condition of *Drosophila parkin* mutants fed with chelators of redox-active metals.
2. *Parkin* mutant in the fly is largely rescued by metal-responsive transcription factor (MTF-1)
3. Zinc supplement greatly improves the condition of *parkin* mutant *Drosophila*.

